abcam

ab193359

Anti-CHST3 antibody [EPR15789]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR15789
Purification technique	Affinity purification Protein A
Concentration	1.636 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Mouse
Dilution info	1/1000 - 1/2000
Notes	-

Species	Human

Dilution info	1/1000 - 1/2000
Notes	-

Predicted

Species	Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/350
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Rat
Dilution info	-
Notes	-

Target data

See full target information CHST3

Function	Sulfotransferase that utilizes 3'-phospho-5'-adenylyl sulfate (PAPS) as sulfonate donor to catalyze the transfer of sulfate to position 6 of the N-acetylgalactosamine (GalNAc) residue of chondroitin (PubMed:15215498, PubMed:9714738, PubMed:9883891). Chondroitin sulfate constitutes the predominant proteoglycan

present in cartilage and is distributed on the surfaces of many cells and extracellular matrices (PubMed:9714738). Catalyzes with a lower efficiency the sulfation of Gal residues of keratan sulfate, another glycosaminoglycan (PubMed:9714738). Can also catalyze the sulfation of the Gal residues in sialyl N-acetyllactosamine (sialyl LacNAc) oligosaccharides (By similarity). May play a role in the maintenance of naive T-lymphocytes in the spleen (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


3 product images



Flow Cytometry (Intracellular) - Anti-CHST3 antibody [EPR15789] (ab193359)

CHST3 Flow Cytometry (Intracellular) staining using rabbit Anti-CHST3 antibody

Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed Jurkat cells labeling CHST3 with ab193359 at 1/350 dilution (red) compared with a rabbit monoclonal IgG isotype control (black) and a unlabelled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (FITC) at 1/150 dilution was used as the secondary antibody.



Western blot - Anti-CHST3 antibody [EPR15789] (ab193359)

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-CHST3 antibody [EPR15789] (ab193359) at 1/1000 dilution

Lane 1:
293 cell lysate at 20 µg

Lane 2:
HeLa cell lysate at 20 µg

Lane 3:
HepG2 cell lysate at 20 µg

Lane 4:
Jurkat cell lysate at 20 µg


Secondary

All lanes:

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 54 kDa

Observed band size: 55 kDa



Western blot - Anti-CHST3 antibody [EPR15789] (ab193359)

CHST3 Western blot staining of Raw264.7 cell lysate using rabbit Anti-CHST3 antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-CHST3 antibody [EPR15789] (ab193359) at 1/1000 dilution

All lanes:
Raw264.7 cell lysate at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 54 kDa

Observed band size: 55 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.