abcam

ab194982

Alexa Fluor® 647 Anti-E-Cadherin/CD324 antibody [EP700Y] - Intercellular Junction Marker

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	Alexa Fluor® 647
Excitation/Emission	Ex: 650nm, Em: 665nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EP700Y
Purification technique	Affinity purification Protein A
Dissociation constant	2.8 x 10^{-11} M
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100
Notes	This product gave a positive signal in CaCo2 cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min).

Not recommended

Species	Mouse
Dilution info	-
Notes	This product gave a positive signal in CaCo2 cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min).

Species	Rat
Dilution info	-
Notes	This product gave a positive signal in CaCo2 cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min).

Flow Cyt (Intra)

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.

Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information E-Cadherin/CD324

Function	Cadherins are calcium-dependent cell adhesion proteins (PubMed:11976333). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells (PubMed:11976333). Promotes organization of radial actin fiber structure and cellular response to contractile forces, via its interaction with AMOTL2 which facilitates anchoring of radial actin fibers to CDH1 junction complexes at the cell membrane (By similarity). Plays a role in the early stages of desmosome cell-cell junction formation via facilitating the recruitment of DSG2 and DSP to desmosome plaques (PubMed:29999492). Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7. E-Cad/CTF2 promotes non-amyloidogenic degradation of Abeta precursors. Has a strong inhibitory effect on APP C99 and C83 production. (Microbial infection) Serves as a receptor for Listeria monocytogenes; internalin A (InlA) binds to this protein and promotes uptake of the bacteria.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	E-Cadherin sometimes called CDH1 or Cadherin-1 is a protein that plays a role in cell-to-cell adhesion. This transmembrane protein has a molecular weight of approximately 120 kDa. E-Cadherin is mainly expressed in epithelial tissues of various organs including the skin and gut. Its adhesive function is made possible by its extracellular domain that facilitates homophilic binding between cells contributing to the maintenance of tissue architecture and cellular integrity.
Biological function summary	E-Cadherin participates in establishing and maintaining adherens junctions which are vital for tissue structure. E-Cadherin operates as a core component of the cadherin-catenin complex which links the protein to the actin cytoskeleton. Through this linkage E-Cadherin plays an important role in signaling pathways that influence cellular growth and differentiation. The protein's ability to mediate intercellular connections also regulates cellular motility and supports basic aspects of cell behavior in epithelial tissues.
Pathways	E-Cadherin influences both the Wnt signaling pathway and the epithelial-to-mesenchymal transition (EMT). Within the Wnt signaling pathway E-Cadherin partners with β-catenin a significant player in transcription regulation and cell signaling. Disruption in E-Cadherin's adhesive functionality can lead to increased β-catenin availability affecting downstream transcriptional control. In the EMT process E-Cadherin loss characterizes an important step in which cells gain migratory and invasive properties typically seen during metastasis in cancer progression.
Associated diseases and disorders	E-Cadherin's role is prominent in cancer particularly in the context of gastric and breast cancers. Mutations or altered expression of E-Cadherin lead to diminished cell adhesion promoting tumorigenesis and metastatic spread. In gastric cancer for instance mutated E-Cadherin often associates with loss of epithelial function

facilitating cancer cell dissemination. Additionally its loss or dysfunction may correlate with proteins such as β-catenin further impacting cancer progression and pathology.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

7 product images




Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-E-Cadherin/CD324 antibody [EP700Y] - Intercellular Junction Marker (ab194982)

E-Cadherin/CD324 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-E-Cadherin/CD324 antibody

ab194982 staining E Cadherin in CaCo2 cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab194982 at 1/100 dilution (shown in red) and ab195887, Mouse monoclonal to alpha Tubulin (Alexa Fluor® 488), at 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).
Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).


This product also gave a positive signal under the same testing conditions in CaCo2 cells fixed with 4% formaldehyde (10 min).



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-E-Cadherin/CD324 antibody [EP700Y] - Intercellular Junction Marker (ab194982)

E-Cadherin/CD324 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-E-Cadherin/CD324 antibody

ab194982 staining E Cadherin in A431 cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab194982 at 1/100 dilution (shown in red) and ab195887, Mouse monoclonal to alpha Tubulin (Alexa Fluor® 488), at 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).
Image was taken with an Operetta High Content Analysis System (Perkin Elmer).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Alexa Fluor® 647 Anti-E-Cadherin/CD324 antibody [EP700Y] - Intercellular Junction Marker (ab194982)

E-Cadherin/CD324 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of human cerebrum tissue using rabbit Anti-E-Cadherin/CD324 antibody


Immunohistochemical analysis of paraffin-embedded human cerebrum tissue labeling E Cadherin with ab194982 at 1/100 dilution.
Negative control: no staining on human cerebrum.

The primary antibody was incubated for 60 mins at room temperature (shown in magenta). Nuclear DNA was labeled with DAPI (shown in blue). The section was then mounted using Fluoromount®.

The immunostaining was performed on a Leica Biosystems BOND RX instrument.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 40 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Alexa Fluor® 647 Anti-E-Cadherin/CD324 antibody [EP700Y] - Intercellular Junction Marker (ab194982)

E-Cadherin/CD324 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of human colon cancer tissue using rabbit Anti-E-Cadherin/CD324 antibody

Immunohistochemical analysis of paraffin-embedded human colon cancer tissue labeling E Cadherin with ab194982 at 1/100 dilution.
Positive staining on human colon cancer.


The primary antibody was incubated for 60 mins at room temperature (shown in magenta). Nuclear DNA was labeled with DAPI (shown in blue). The section was then mounted using

Fluoromount®.

The immunostaining was performed on a Leica Biosystems BOND RX instrument.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 40 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Alexa Fluor® 647 Anti-E-Cadherin/CD324 antibody [EP700Y] - Intercellular Junction Marker (ab194982)

E-Cadherin/CD324 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of human colon tissue using rabbit Anti-E-Cadherin/CD324 antibody


Immunohistochemical analysis of paraffin-embedded human colon tissue labeling E Cadherin with ab194982 at 1/100 dilution.
Positive staining on human colon.

The primary antibody was incubated for 60 mins at room temperature (shown in magenta). Nuclear DNA was labeled with DAPI (shown in blue). The section was then mounted using Fluoromount®.

The immunostaining was performed on a Leica Biosystems BOND RX instrument.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 40 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Alexa Fluor® 647 Anti-E-Cadherin/CD324 antibody [EP700Y] - Intercellular Junction Marker (ab194982)

E-Cadherin/CD324 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of human cerebrum tissue using rabbit Anti-E-Cadherin/CD324 antibody


Immunohistochemical analysis of paraffin-embedded human cerebrum tissue labeling E Cadherin with ab206878 at 1/100 dilution.
Negative control: no staining on human cerebrum.

The primary antibody was incubated for 60 mins at room temperature (shown in green). Nuclear DNA was labeled with DAPI (shown in blue). The section was then mounted using Fluoromount®.

The immunostaining was performed on a Leica Biosystems BOND RX instrument.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 40 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Alexa Fluor® 647 Anti-E-Cadherin/CD324 antibody [EP700Y] - Intercellular Junction Marker (ab194982)

E-Cadherin/CD324 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of human colon cancer tissue using rabbit Anti-E-Cadherin/CD324 antibody

Immunohistochemical analysis of paraffin-embedded human colon cancer tissue labeling E Cadherin with ab206878 at 1/100 dilution.
Positive staining on human colon cancer.

The primary antibody was incubated for 60 mins at room temperature (shown in green). Nuclear DNA was labeled with DAPI (shown in blue). The section was then mounted using Fluoromount®.

The immunostaining was performed on a Leica Biosystems BOND RX instrument.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 40 mins.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.