abcam

ab195255

Alexa Fluor® 647 Anti-ALDH1A1 antibody [EP1933Y]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	Alexa Fluor® 647
Excitation/Emission	Ex: 650nm, Em: 665nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EP1933Y
Purification technique	Affinity purification Protein A
Concentration	Concentration data is not available right now, please try again later Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human

Dilution info	1/50
Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/50
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Expected

Species	Human, Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Rat
Dilution info	-
Notes	-

Target data

See full target information ALDH1A1

Function

Cytosolic dehydrogenase that catalyzes the irreversible oxidation of a wide range of aldehydes to their corresponding carboxylic acid (PubMed:12941160, PubMed:15623782, PubMed:17175089, PubMed:19296407, PubMed:25450233, PubMed:26373694). Functions downstream of retinol dehydrogenases and catalyzes the oxidation of retinaldehyde into retinoic acid, the second step in the oxidation of retinol/vitamin A into retinoic acid (By similarity). This pathway is crucial to control the levels of retinol and retinoic acid, two important molecules which excess can be teratogenic and cytotoxic (By similarity). Also oxidizes aldehydes resulting from lipid peroxidation like (E)-4-hydroxynon-2-enal/HNE, malonaldehyde and hexanal that form protein adducts and are highly cytotoxic. By participating for instance to the clearance of (E)-4-hydroxynon-2-enal/HNE in the lens epithelium prevents the formation of HNE-protein adducts and lens opacification (PubMed:12941160, PubMed:15623782, PubMed:19296407). Also functions downstream of fructosamine-3-kinase in the fructosamine degradation pathway by catalyzing the oxidation of 3-deoxyglucosone, the carbohydrate product of fructosamine 3-phosphate decomposition, which is itself a potent glycating agent that may react with lysine and arginine side-chains of proteins (PubMed:17175089). Also has an aminobutyraldehyde dehydrogenase activity and is probably part of an alternative pathway for the biosynthesis of GABA/4-aminobutanoate in midbrain, thereby playing a role in GABAergic synaptic transmission (By similarity).

Storage

Shipped at conditions	Blue Ice

Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	ALDH1A1 also known as aldehyde dehydrogenase 1 family member A1 is an enzyme with a molecular weight of approximately 54 kDa. It functions by catalyzing the irreversible oxidation of aldehydes to carboxylic acids which is an important step in detoxifying alcohol-derived acetaldehyde and other aldehydes. ALDH1A1 is expressed in various tissues including liver kidney and cornea. Its activity allows for maintaining cellular homeostasis by processing endogenous and exogenous aldehydes.
Biological function summary	ALDH1A1 plays significant roles in the biosynthesis of retinoic acid an important regulator of gene expression in developmental and adult processes. It does not form part of a larger enzyme complex but functions independently. The ALDH1A1 proteins contribute to the regulation of cell proliferation differentiation and survival. Researchers often measure its expression using methods like the ALDH1A1 ELISA which quantifies the protein levels in biological samples.

Pathways	The ALDH1A1 enzyme is notably implicated in the retinoid metabolism pathway and the detoxification of alcohol-derived aldehydes within the body. It is closely related to other members of the aldehyde dehydrogenase family including ALDH2 which shares similar detoxification functions. ALDH1A1's role in retinoic acid production links it to gene expression pathways associated with cell differentiation and growth.
Associated diseases and disorders	ALDH1A1 has associations with cancer and alcohol-induced liver disease. Its elevated expression in cancerous tissues often correlates with stem-cell like characteristics and chemoresistance. A connection exists between ALDH1A1 and ALDH2 in the context of alcohol metabolism where ALDH1A1's role in detoxifying acetaldehyde contributes to the lower risk of alcohol-induced damage. Studies explore ALDH1A1 antibodies and inhibitors as potential therapeutic approaches for these conditions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


3 product images



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-ALDH1A1 antibody [EP1933Y] (ab195255)

ALDH1A1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ALDH1A1 antibody

ab195255 staining ALDH1A1 in MCF7 cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized in 0.1% Triton X-100 for 5 minutes and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab195255 at a working dilution of 1 in 50 (shown in red) and ab195887, Mouse monoclonal [DM1A] to alpha Tubulin (Alexa Fluor® 488, shown in green) at 2µg/ml overnight at +4°C. Nuclear DNA was labelled in blue with DAPI.
Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Alexa Fluor® 647 Anti-ALDH1A1 antibody [EP1933Y] (ab195255)

ALDH1A1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human colon cancer tissue using rabbit Anti-ALDH1A1 antibody

Immunohistochemical analysis of paraffin-embedded Human colon cancer tissue labelling ALDH1A1 with ab195255 at 1/50 (10.0 ug/ml) dilution.


Positive staining on human colon cancer.

The primary antibody was incubated for 60 mins at room temperature (shown in magenta). Nuclear DNA was labeled with DAPI (shown in blue). The section was then mounted using Fluoromount®.

The immunostaining was performed on a Leica Biosystems BOND RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Counterstained with DAPI.

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 40 mins.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Alexa Fluor® 647 Anti-ALDH1A1 antibody [EP1933Y] (ab195255)

ALDH1A1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human hepatocellular cancer tissue using rabbit Anti-ALDH1A1 antibody

Immunohistochemical analysis of paraffin-embedded human hepatocellular cancer tissue labelling ALDH1A1 with ab195255 at 1/50 (10.0 ug/ml) dilution.

Positive staining on human hepatocellular cancer.

The primary antibody was incubated for 60 mins at room temperature (shown in magenta). Nuclear DNA was labeled with DAPI (shown in blue). The section was then mounted using Fluoromount®.

The immunostaining was performed on a Leica Biosystems BOND RX instrument. Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Counterstained with DAPI.

Heat mediated antigen retrieval was performed with Tris-EDTA buffer (pH 9.0, Epitope Retrieval Solution2) for 40 mins.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.