abcam

ab196465

Alexa Fluor® 488 Anti-NQO1 antibody [EPR3309]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	Alexa Fluor® 488
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR3309
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human

Dilution info	1/100
Notes	-

Expected

Species	Mouse
Dilution info	-
Notes	-

Species	Rat
Dilution info	1/100
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	ab199091 - Rabbit monoclonal IgG (Alexa FluorÂ® 488), is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse
Dilution info	-
Notes	-

Species	Rat
Dilution info	1/500

Notes	ab199091 - Rabbit monoclonal IgG (Alexa FluorÂ® 488), is suitable for use as an isotype control with this antibody.

Not recommended

Species	Rat
Dilution info	-
Notes	ab199091 - Rabbit monoclonal IgG (Alexa FluorÂ® 488), is suitable for use as an isotype control with this antibody.

Target data

See full target information NQO1

Function

Flavin-containing quinone reductase that catalyzes two-electron reduction of quinones to hydroquinones using either NADH or NADPH as electron donors. In a ping-pong kinetic mechanism, the electrons are sequentially transferred from NAD(P)H to flavin cofactor and then from reduced flavin to the quinone, bypassing the formation of semiquinone and reactive oxygen species (By similarity) (PubMed:8999809, PubMed:9271353). Regulates cellular redox state primarily through quinone detoxification. Reduces components of plasma membrane redox system such as coenzyme Q and vitamin quinones, producing antioxidant hydroquinone forms. In the process may function as superoxide scavenger to prevent hydroquinone oxidation and facilitate excretion (PubMed:15102952, PubMed:8999809, PubMed:9271353). Alternatively, can activate quinones and their derivatives by generating redox reactive hydroquinones with DNA cross-linking antitumor potential (PubMed:8999809). Acts as a gatekeeper of the core 20S proteasome known to degrade proteins with unstructured regions. Upon oxidative stress, interacts with tumor suppressors TP53 and TP73 in a NADH-dependent way and inhibits their ubiquitin-independent degradation by the 20S proteasome (PubMed:15687255, PubMed:28291250).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C

Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	NQO1 also known as NAD(P)H:quinone oxidoreductase 1 is a cytosolic enzyme involved in two-electron reduction processes. This protein plays a role in detoxification transforming quinones into less reactive and harmful hydroquinones. NQO1 has a molecular weight of approximately 31 kDa and is present in various tissues with high expression in the liver and lungs. Sometimes called DT-diaphorase it contributes to antioxidant protection within cells and assists in the stabilization of other proteins such as p53.
Biological function summary	NQO1 serves a protective function against oxidative stress by reducing quinones and preventing redox cycling that generates reactive oxygen species. Part of a critical network its function integrates with the cellular defense mechanism assisting in maintaining cellular homeostasis. NQO1 helps metabolize xenobiotics

	and is associated with phase II detoxification working alongside enzymes like glutathione S-transferases but is not part of a complex.
Pathways	NQO1 is an important component of the antioxidant defense pathway participating in the direct enzymatic reduction of quinones protecting cells from oxidative damage. It also interfaces with the KEAP1-NRF2 pathway where the NQO1 gene is regulated by the NRF2 transcription factor that upregulates its expression in response to oxidative stress. NQO1 interlinks with proteins such as p53 through pathways related to apoptotic regulation and cellular stress responses.
Associated diseases and disorders	Mutations or altered expression of NQO1 have correlations with cancer development and progression notably in liver and lung cancers. This protein's stability influences cancer cell survival particularly under oxidative stress or chemotherapeutic treatments. NQO1 also exhibits links to Alzheimer's disease where its potential role in neuroprotection against oxidative damage draws investigation relating it to proteins like tau and amyloid-beta.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

2 product images




Flow Cytometry (Intracellular) - Alexa Fluor® 488 Anti-NQO1 antibody [EPR3309] (ab196465)

NQO1 Flow Cytometry (Intracellular) staining using rabbit Anti-NQO1 antibody

Overlay histogram showing A549 cells stained with ab196465 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab196465, 1/500 dilution) for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) Alexa Fluor® 488 used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.
Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.

This antibody gave a positive signal in A549 fixed with 4% formaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-NQO1 antibody [EPR3309] (ab196465)

NQO1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-NQO1 antibody

ab196465 staining NQO1 A549 cells. The cells were fixed with 100% methanol (5 min), permeabilized in 0.1% Triton X-100 for 5 minutes and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab196465 at a working dilution of 1 in 100 (shown in green) and ab195889, Mouse monoclonal [DM1A] to alpha Tubulin (Alexa Fluor® 594, shown in red) at 2µg/ml overnight at +4°C. Nuclear DNA was labelled in blue with DAPI.
Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.