abcam

ab198996

Anti-RCN1/RCN antibody [EPR17163] - C-terminal

Recombinant | RabMAb | KO Validated

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR17163
Purification technique	Affinity purification Protein A
Concentration	1.876 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/100
Notes	-

Expected

Species	Mouse
Dilution info	1/100
Notes	-

Species	Rat
Dilution info	1/100
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/250
Notes	-

Expected

Species	Mouse
Dilution info	1/250

Notes	-

Species	Rat
Dilution info	1/250
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/70
Notes	-

Expected

Species	Mouse
Dilution info	1/70
Notes	-

Species	Rat
Dilution info	1/70
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/800
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Expected

Species	Mouse
Dilution info	1/800

Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/800
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information RCN1

Function	May regulate calcium-dependent activities in the endoplasmic reticulum lumen or post-ER compartment.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	RCN1 also known as reticulocalbin-1 or RCN is a calcium-binding protein located in the lumen of the endoplasmic reticulum (ER). It belongs to the CREC family of proteins which are characterized by their EF-hand motifs that mediate calcium binding. RCN1 has a molecular mass of around 37 kDa. This protein is expressed in various tissues including the pancreas heart and brain suggesting its involvement in diverse cellular functions. Within the ER RCN1 interacts with other proteins aiding in their proper folding and assembly.
Biological function summary	RCN1 plays a role in cellular calcium homeostasis and may influence processes such as protein folding and trafficking. It is not known to be a part of a larger complex but it interacts with other ER-resident proteins. Maintaining calcium levels is important within the ER where RCN1 helps buffer calcium concentrations. This function supports processes that are sensitive to calcium fluctuations such as signal transduction and stress responses.
Pathways	The role of RCN1 in cellular mechanisms connects it to important pathways like the calcium signaling pathway and the unfolded protein response pathway. In the calcium signaling pathway RCN1 aids in regulating calcium levels within the ER affecting downstream signaling events. Meanwhile in the unfolded protein response pathway RCN1 participates by ensuring the proper folding of proteins particularly during times of stress when misfolded proteins accumulate. RCN1's interaction with calumenin another calcium-binding protein highlights its involvement in these pathways.
Associated diseases and disorders	RCN1 has associations with cardiovascular disease and some forms of cancer. In cardiovascular disease altered expression of RCN1 may impact calcium signaling and lead to dysfunction in heart cells. Similarly in certain cancers changes in RCN1 expression levels could affect cellular proliferation via calcium-regulated pathways. The interaction between RCN1 and calreticulin a multifunctional protein involved in calcium homeostasis and immune response further connects RCN1's influence on disease progression.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


7 product images



Immunocytochemistry/ Immunofluorescence - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996)

RCN1/RCN Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-RCN1/RCN antibody

Immunocytochemistry/Immunofluorescence analysis of HeLa cells labelling RCN1/RCN with ab198996 at 1/500. Cells were fixed with 100% methanol. ab150077, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody.
Control: PBS only.
Nuclear counter stain: DAPI.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996)

RCN1/RCN Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-RCN1/RCN antibody

Immunohistochemical analysis of paraffin-embedded Human cerebral cortex tissue labeling RCN1/RCN with ab198996 at 1/1600, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500. Cytoplasm staining on Human cerebral cortex tissue is observed. Subcellular location Endoplasmic reticulum lumen [UniProt]. Counter stained with Hematoxylin.
Negative control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996)

RCN1/RCN Flow Cytometry (Intracellular) staining using rabbit Anti-RCN1/RCN antibody

Intracellular Flow Cytometry analysis of HeLa cells labelling RCN1/RCN (red) with purified ab198996 at dilution of 1/70. The secondary antibody used was Alexa Fluorr® 488 goat-anti-rabbit IgG (1/2000). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Isotype control antibody was Rabbit monoclonal IgG (black). The blue line shows cells without incubation with primary antibody and secondary antibody.



Immunoprecipitation - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996)

RCN1/RCN was immunoprecipitated from 1mg of HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell extract with ab198996 at 1/100. Western blot was performed from the immunoprecipitate using ab198996 at 1/1000. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500.
Lane 1: HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell extract. 10ug (Input.

Lane 2: HeLa whole cell extract.

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab198996 in HeLa whole cell extract.


The expression profile observed is consistent with what has been described in the literature PMID: 8416973.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

All lanes:
Immunoprecipitation - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996)

Predicted band size: 39 kDa

Observed band size: 44 kDa, 46 kDa



Western blot - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996)

RCN1/RCN Western blot staining using rabbit Anti-RCN1/RCN antibody

Western blot: Anti-RCN1/RCN antibody [EPR17163] - C-terminal ab198996 staining at 1/1000 dilution, shown in green; Mouse anti-CANX ab238078 loading control staining at 1/20000 dilution, shown in magenta. A band was observed at 45 kDa in Wild-type HEK-293T cell lysates with no signal observed at this size in RCN1 knockout HEK-293T cell line. To generate this image, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 5% Milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit 800CW and Goat anti-Mouse 680RD at 1/20,000 dilution.
All lanes:
Western blot - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996) at 1/1000 dilution

Lane 1:
Wild-type HEK-293T cell lysates at 20 µg

Lane 2:
Western blot - Human RCN1 (RCN) knockout HEK-293T cell line (ab266098) at 20 µg

Lane 3:
HeLa cell lysates at 20 µg

Lane 4:
Daudi cell lysates at 20 µg


Secondary

All lanes:
Goat anti-Rabbit 800CW & Goat anti-Mouse 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 45 kDa

Observed band size: 45 kDa



Western blot - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996)

Blocking and diluting buffer was 5% NFDM/TBST.
The expression profile observed is consistent with what has been described in the literature PMID: 8416973.

All lanes:
Western blot - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996) at 1/10000 dilution

Lane 1:
293 (Human epithelial cells from embryonic kidney) cell lysate at 10 µg

Lane 2:
HeLa (Human epithelial cells from cervix adenocarcinoma) cell lysate at 10 µg

Secondary


All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Developed using the ECL technique.

Predicted band size: 39 kDa

Observed band size: 44 kDa, 46 kDa

Exposure time: 1min



Western blot - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996)

Blocking and diluting buffer was 5% NFDM/TBST.
The expression profile observed is consistent with what has been described in the literature PMID: 8416973.
All lanes:
Western blot - Anti-RCN1/RCN antibody [EPR17163] - C-terminal (ab198996) at 1/1000 dilution

Lane 1:
Raw264.7 (Mouse macrophage cells transformed with Abelson murine leukemia virus) cell lysate at 10 µg

Lane 2:
Rat heart tissue lysate at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Developed using the ECL technique.

Predicted band size: 39 kDa

Observed band size: 44 kDa, 46 kDa

Exposure time: 1min

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.