abcam

ab199733

Alexa Fluor® 488 Anti-Superoxide Dismutase 1 antibody [EP1727Y]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	Alexa Fluor® 488
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EP1727Y
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	1/100
Notes	This product gave a positive signal in HeLa cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min).

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Expected

Species	Human, Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information SOD1

Function	Destroys radicals which are normally produced within the cells and which are toxic to biological systems (PubMed:24140062). Catalyzes the oxidation of hydrogen sulfide (H2S) to sulfate, playing an important role in detoxifying H2S and limiting the accumulation of reactive sulfur species (RSS) such as persulfides and polysulfides (PubMed:36630448).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Superoxide Dismutase 1 (SOD1) also known as Cu/Zn SOD or simply dismutase is an enzyme important for the detoxification of superoxide radicals. SOD1 catalyzes the conversion of two superoxide molecules into oxygen and hydrogen peroxide maintaining cellular redox balance. This enzyme typically exhibits a mass of approximately 32 kDa. SOD1 is expressed in the cytoplasm of cells throughout the body including tissues like liver kidney and brain.
Biological function summary	The enzyme functions as a homodimer with each subunit containing a copper and zinc ion. These metal ions are essential for the catalytic activity of SOD1 as the copper ion participates in electron transfer while the zinc ion provides structural stability. The enzyme protects cells from oxidative stress by neutralizing excess reactive oxygen species ensuring cellular health and functioning.
Pathways	SOD1 plays an important role in the cellular antioxidant defense system and is a part of the reactive oxygen species (ROS) metabolic pathway. It works in conjunction with catalase and glutathione peroxidase to limit oxidative damage within cells. The close interaction between these enzymes highlights the interdependence within the antioxidant defense network emphasizing their role in maintaining cellular homeostasis.
Associated diseases and disorders	Mutations in the SOD1 gene are linked to familial Amyotrophic Lateral Sclerosis (ALS) a neurodegenerative condition. In ALS aberrant SOD1 proteins may lead to increased oxidative stress and motor neuron damage. Research also connects SOD1 to the pathogenesis of Alzheimer's disease where oxidative stress is a contributing factor. These associations underline SOD1's significance in neurodegenerative diseases highlighting potential therapeutic targets for interventions focused on oxidative stress management.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

2 product images




Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-Superoxide Dismutase 1 antibody [EP1727Y] (ab199733)

Superoxide Dismutase 1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Superoxide Dismutase 1 antibody

ab199733 staining Superoxide Dismutase 1 in HeLa cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab at a 1/100 dilution (shown in green) and ab195889, Mouse monoclonal to alpha Tubulin (Alexa Fluor® 594), at a 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).
Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in HeLa cells fixed with 4% formaldehyde (10 min).



Flow Cytometry (Intracellular) - Alexa Fluor® 488 Anti-Superoxide Dismutase 1 antibody [EP1727Y] (ab199733)

Superoxide Dismutase 1 Flow Cytometry (Intracellular) staining using rabbit Anti-Superoxide Dismutase 1 antibody

Overlay histogram showing HeLa cells stained with ab199733 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab199733, 1/500 dilution) for 30 min at 22°C. Isotype control antibody (black line) was rabbit monoclonal IgG [EPR25A] Alexa Fluor® 488 (ab199091) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in HeLa cells fixed with 4% formaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.