abcam

ab20143

Anti-Integrin alpha V beta 3 antibody [23C6]

Key facts

Isotype	IgG1
Host species	Mouse
Storage buffer	Preservative: 0.02% Sodium azide Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	Cell preparation containing ITGAV protein. The exact immunogen used to generate this antibody is proprietary information. Database link P06756
Clone number	23C6
Purification technique	Affinity purification Protein A/G
Specificity	Reacts with Integrin Alpha V Beta 3. 23C6 may be useful for bone resorption modulation, osteoclast identification, receptor purification, malignant melanoma identification and treatment.
Light chain type	unknown
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Expected

Species	Human

Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt

Expected

Species	Human
Dilution info	-
Notes	ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.

FuncS (Neut/Block)

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-Fr

Expected

Species	Human
Dilution info	-
Notes	Samples should be acetone fixed.

Target data

See full target information ITGAV

Function The alpha-V (ITGAV) integrins are receptors for vitronectin, cytotactin, fibronectin, fibrinogen, laminin, matrix metalloproteinase-2, osteopontin, osteomodulin, prothrombin, thrombospondin and vWF. They recognize the sequence R-G-D in a wide array of ligands. ITGAV:ITGB3 binds to fractalkine (CX3CL1) and may act as its coreceptor in CX3CR1-dependent fractalkine signaling (PubMed:23125415). ITGAV:ITGB3 binds to NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling (PubMed:20682778). ITGAV:ITGB3 binds to FGF1 and this binding is essential for FGF1 signaling (PubMed:18441324). ITGAV:ITGB3 binds to FGF2 and this binding is essential for FGF2 signaling (PubMed:28302677).

ITGAV:ITGB3 binds to IGF1 and this binding is essential for IGF1 signaling (PubMed:19578119). ITGAV:ITGB3 binds to IGF2 and this binding is essential for IGF2 signaling (PubMed:28873464). ITGAV:ITGB3 binds to IL1B and this binding is essential for IL1B signaling (PubMed:29030430). ITGAV:ITGB3 binds to PLA2G2A via a site (site 2) which is distinct from the classical ligand-binding site (site 1) and this induces integrin conformational changes and enhanced ligand binding to site 1 (PubMed:18635536, PubMed:25398877). ITGAV:ITGB3 and ITGAV:ITGB6 act as receptors for fibrillin-1 (FBN1) and mediate R-G-D-dependent cell adhesion to FBN1 (PubMed:12807887, PubMed:17158881). Integrin alpha-V/beta-6 or alpha-V/beta-8 (ITGAV:ITGB6 or ITGAV:ITGB8) mediates R-G-D-dependent release of transforming growth factor beta-1 (TGF-beta-1) from regulatory Latency-associated peptide (LAP), thereby playing a key role in TGF-beta-1 activation (PubMed:15184403, PubMed:22278742, PubMed:28117447). ITGAV:ITGB3 acts as a receptor for CD40LG (PubMed:31331973). ITGAV:ITGB3 acts as a receptor for IBSP and promotes cell adhesion and migration to IBSP (PubMed:10640428). (Microbial infection) Integrin ITGAV:ITGB5 acts as a receptor for Adenovirus type C. (Microbial infection) Integrin ITGAV:ITGB5 and ITGAV:ITGB3 act as receptors for Coxsackievirus A9 and B1. (Microbial infection) Integrin ITGAV:ITGB3 acts as a receptor for Herpes virus 8/HHV-8. (Microbial infection) Integrin ITGAV:ITGB6 acts as a receptor for herpes simplex 1/HHV-1. (Microbial infection) Integrin ITGAV:ITGB3 acts as a receptor for Human parechovirus 1. (Microbial infection) Integrin ITGAV:ITGB3 acts as a receptor for West nile virus. (Microbial infection) In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Integrin alpha V beta 3 also known as integrin αvβ3 or avb3 integrin is a transmembrane receptor with a molecular weight of about 140 kDa. This protein is composed of the alpha V (αv) and beta 3 (β3) subunits. Integrin αvβ3 is widely expressed in various cell types including endothelial cells osteoclasts and some

	tumor cells. Its function involves mechanical interaction with the extracellular matrix (ECM) allowing cells to adhere to ECM proteins such as vitronectin and fibronectin. This integrin plays an important role in cell adhesion migration and survival.
Biological function summary	Integrin alpha V beta 3 engages in various cellular processes by serving as a part of integrin complexes. It mediates signal transduction from the ECM to the cell interior influencing cellular responses such as shape changes survival and proliferation. The receptor regulates angiogenesis which is the formation of new blood vessels and osteoclast-mediated bone resorption. These functions are essential for tissue remodeling and repair. The integrin can also modulate immune responses by interacting with immune cells affecting inflammation and wound healing processes.
Pathways	Integrin alpha V beta 3 is an important component in signaling pathways such as the PI3K/AKT and MAPK pathways. These pathways facilitate the integrin's role in cell migration growth and survival. It interacts with proteins like focal adhesion kinase (FAK) and SRC kinase bridging signals from the ECM to intracellular signaling cascades. The integrin's ability to bind ligands and initiate these pathways makes it an important player in cellular responses to the extracellular environment influencing how cells align mechanically and biochemically.
Associated diseases and disorders	Integrin alpha V beta 3 has significant associations with cancer progression and osteoporosis. In various cancers including melanoma and breast cancer overexpression of this integrin supports tumor angiogenesis and metastasis connecting its role with angiogenic growth factors and matrix metalloproteinases. In osteoporosis it collaborates with proteins like osteopontin promoting osteoclast attachment to the bone matrix thereby influencing bone degradation. Understanding its role in these disorders highlights potential therapeutic targets for inhibiting unwanted angiogenesis and excessive bone resorption.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.