abcam

ab201684

Anti-PRPF4 antibody [EPR17206(B)]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR17206(B)
Purification technique	Affinity purification Protein A
Concentration	0.576 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/40
Notes	-

WB

Tested

Species	Human
Dilution info	1/5000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/1200
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/50
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/3000
Notes	-

Target data

See full target information PRPF4

Function Plays a role in pre-mRNA splicing as component of the U4/U6-U5 tri-snRNP complex that is involved in spliceosome assembly, and as component of the precatalytic spliceosome (spliceosome B complex).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


7 product images



Flow Cytometry (Intracellular) - Anti-PRPF4 antibody [EPR17206(B)] (ab201684)

PRPF4 Flow Cytometry (Intracellular) staining using rabbit Anti-PRPF4 antibody


Intracellular Flow Cytometry analysis of HeLa cells labelling PRPF4 (red) with purified ab201684 at dilution of 1/50. The secondary antibody used was Alexa Fluorr® 488 goat-anti-rabbit IgG (1/2000). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Isotype control antibody was Rabbit monoclonal IgG (black). The blue line shows cells without incubation with primary antibody and secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-PRPF4 antibody [EPR17206(B)] (ab201684)

PRPF4 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-PRPF4 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed 0.1% Triton X-100 permeabilized HeLa (Human epithelial cells from cervix adenocarcinoma) cells labeling PRPF4 with ab201684 at 1/1200 dilution followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/500 dilution (green). Confocal image showing nuclear staining on HeLa cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).
The negative controls are as follows:
-ve control 1: ab201684 at 1/1200 dilution followed by ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.
-ve control 2: ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution followed by ab150077 (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.




Immunocytochemistry/ Immunofluorescence - Anti-PRPF4 antibody [EPR17206(B)] (ab201684)

PRPF4 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-PRPF4 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed 0.1% Triton X-100 permeabilized MCF7 (Human breast adenocarcinoma cell line) cells labeling PRPF4 with ab201684 at 1/1200 dilution followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/500 dilution (green). Confocal image showing nuclear staining on MCF7 cell line. The nuclear counter

stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).
The negative controls are as follows:
-ve control 1: ab201684 at 1/1200 dilution followed by ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.
-ve control 2: ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution followed by ab150077 (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-PRPF4 antibody [EPR17206(B)] (ab201684)

PRPF4 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-PRPF4 antibody

Immunohistochemical analysis of paraffin-embedded Human cervix carcinoma tissue labeling PRPF4 with ab201684 at 1/3000 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) secondary antibody at 1/500 dilution. Nuclear staining on Human cervix carcinoma tissue is observed. Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunoprecipitation - Anti-PRPF4 antibody [EPR17206(B)] (ab201684)

PRPF4 was immunoprecipitated from 1mg of HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell lysate with ab201684 at 1/40 dilution. Western blot was performed from the immunoprecipitate using ab201684 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used for detection at 1/1500 dilution.
Lane 1: HeLa whole cell lysate 10 µg (Input). Lane 2: ab201684 IP in HeLa whole cell lysate. Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab201684 in HeLa whole cell lysate.
Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 5 seconds.

All lanes:
Immunoprecipitation - Anti-PRPF4 antibody [EPR17206(B)] (ab201684)

Predicted band size: 58 kDa



Western blot - Anti-PRPF4 antibody [EPR17206(B)] (ab201684)

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-PRPF4 antibody [EPR17206(B)] (ab201684) at 1/5000 dilution

Lane 1:
HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell lysate at 20 µg

Lane 2:
SW480 (Human colon adenocarcinoma cell line) whole cell lysate at 20 µg

Lane 3:
Raji (Human Burkitt's lymphoma cell line) whole cell lysate at 20 µg

Lane 4:
Jurkat (Human T cell leukemia cells from peripheral blood) whole cell lysate at 20 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 58 kDa

Observed band size: 58 kDa

Exposure time: 3min



Western blot - Anti-PRPF4 antibody [EPR17206(B)] (ab201684)

PRPF4 Western blot staining of Human fetal spleen lysate using rabbit Anti-PRPF4 antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-PRPF4 antibody [EPR17206(B)] (ab201684) at 1/5000 dilution

All lanes:
Human fetal spleen lysate at 10 µg

Secondary

All lanes:
Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG at 1/1000 dilution

Predicted band size: 58 kDa

Observed band size: 58 kDa

Exposure time: 3min

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.