abcam

ab202504

Alexa Fluor® 568 Anti-Vimentin antibody [EPR3776] - Cytoskeleton Marker

Recombinant | RabMAb | KO Validated

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	Alexa Fluor® 568
Excitation/Emission	Ex: 578nm, Em: 603nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR3776
Purification technique	Affinity purification Protein A
Dissociation constant	1.1 x 10^{-10} M
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/1000
Notes	This product gave a positive signal in NIH3T3 cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min) and HeLa cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min)

Species	Human
Dilution info	1/1000
Notes	This product gave a positive signal in NIH3T3 cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min) and HeLa cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min)

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	0.1 µg/mL
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Expected

Species	Human, Mouse, Rat

Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information VIM

Function	Vimentins are class-III intermediate filaments found in various non-epithelial cells, especially mesenchymal cells. Vimentin is attached to the nucleus, endoplasmic reticulum, and mitochondria, either laterally or terminally. Plays a role in cell directional movement, orientation, cell sheet organization and Golgi complex polarization at the cell migration front (By similarity). Protects SCRIB from proteasomal degradation and facilitates its localization to intermediate filaments in a cell contact-mediated manner (By similarity). May promote axon outgrowth and motor fiber repair via DSP-mediated recruitment to outgrowth tips (By similarity). Involved with LARP6 in the stabilization of type I collagen mRNAs for CO1A1 and CO1A2.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by

the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

4 product images




Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 568 Anti-Vimentin antibody [EPR3776] - Cytoskeleton Marker (ab202504)

Vimentin Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Vimentin antibody

ab202504 staining Vimentin in wild-type HAP1 cells (top panel) and Vimentin knockout HAP1 cells (bottom panel). The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab202504 at a 1/1000

dilution (shown in green) and ab195884, Rat monoclonal [YOL1/34] to Tubulin (Alexa Fluor® 647), at a 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).
Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).




Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 568 Anti-Vimentin antibody [EPR3776] - Cytoskeleton Marker (ab202504)

Vimentin Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Vimentin antibody

ab202504 staining Vimentin in HeLa cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab202504 at 1/1000 dilution (**pseudocolored in green**) and ab195884, Rat monoclonal to Tubulin (Alexa Fluor® 647), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).
Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in HeLa cells fixed with 100% methanol (5 min).



Flow Cytometry (Intracellular) - Alexa Fluor® 568 Anti-Vimentin antibody [EPR3776] - Cytoskeleton Marker (ab202504)


Vimentin Flow Cytometry (Intracellular) staining using rabbit Anti-Vimentin antibody

Overlay histogram showing HAP1 wildtype (green line) and HAP1-VIM knockout cells (red line) stained with ab202504. The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS / 10% normal goat serum to block non-specific protein-protein interactions followed by the antibody (ab202504, 0.1µg/ml dilution) for 30 min at 22°C.

A rabbit monoclonal IgG isotype control antibody (ab209613) was used at the same concentration and conditions as the primary antibody (HAP1 wildtype - black line, HAP1-VIM knockout - grey line). Unlabelled sample was also used as a control (this line is not shown for the purpose of simplicity).

Acquisition of >5,000 events were collected using a 50 mW Yellow/Green laser (561nm) and 586/15 bandpass filter.

This antibody can also be used in HAP1 cells fixed with 4% formaldehyde (10 min), permeabilized with 0.1% PBS-Triton X-100 for 15 min under the same conditions.



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 568 Anti-Vimentin antibody [EPR3776] - Cytoskeleton Marker (ab202504)

Vimentin Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Vimentin antibody

ab202504 staining Vimentin in NIH3T3 cells. The cells were fixed with 4% formaldehyde (10 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab202504 at a 1/1000 dilution (shown in pseudocolor green) and ab195884, Rat

monoclonal to alpha Tubulin (Alexa Fluor® 647), at a 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).
Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in NIH3T3 cells fixed with 100% methanol (5 min).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.