abcam

ab202896

Anti-MYBBP1A antibody [EPR7205]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR7205
Purification technique	Affinity purification Protein A
Concentration	0.718 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/70
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	-

Target data

See full target information MYBBP1A

Function May activate or repress transcription via interactions with sequence specific DNA-binding proteins (By similarity). Repression may be mediated at least in part by histone deacetylase activity (HDAC activity) (By similarity). Acts as a corepressor and in concert with CRY1, represses the transcription of the core circadian clock component PER2 (By similarity). Preferentially binds to dimethylated histone H3 'Lys-9' (H3K9me2) on the PER2 promoter (By similarity). Has a role in rRNA biogenesis together with PWP1 (PubMed:29065309).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks

Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


8 product images



Flow Cytometry (Intracellular) - Anti-MYBBP1A antibody [EPR7205] (ab202896)

MYBBP1A Flow Cytometry (Intracellular) staining using rabbit Anti-MYBBP1A antibody

Intracellular Flow Cytometry analysis of HeLa cells labelling MYBBP1A (red) with purified ab202896 at dilution of 1/70. The secondary antibody used was Alexa Fluorr® 488 goat-anti-rabbit IgG (1/2000). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Isotype control antibody used was Rabbit monoclonal IgG (black). The blue line shows cells without incubation with primary antibody and secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MYBBP1A antibody [EPR7205] (ab202896)

MYBBP1A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MYBBP1A antibody

Immunohistochemical analysis of paraffin-embedded Human cerebral cortex tissue labeling MYBBP1A with ab202896 at 1/100 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution. Cytoplasmic and nuclear staining on Human cerebral cortex tissue is observed. Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunocytochemistry/ Immunofluorescence - Anti-MYBBP1A antibody [EPR7205] (ab202896)

MYBBP1A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-MYBBP1A antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed 0.1% Triton X-100 permeabilized HeLa (Human epithelial cells from cervix adenocarcinoma) cells labeling MYBBP1A with ab202896 at 1/500 dilution followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution (green). Confocal image showing nuclear and weakly cytoplasmic staining on HeLa cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution (red).
The negative controls are as follows:
-ve control 1: ab202896 at 1/500 dilution followed by ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution.


-ve control 2: ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution followed by ab150077 (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/1000 dilution.



Immunocytochemistry/ Immunofluorescence - Anti-MYBBP1A antibody [EPR7205] (ab202896)

MYBBP1A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-MYBBP1A antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed 0.1% Triton X-100 permeabilized SH-SY5Y (Human neuroblastoma from bone marrow cells) cells labeling MYBBP1A with ab202896 at 1/500 dilution followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution (green). Confocal image showing nuclear and weakly cytoplasmic staining on SH-SY5Y cell line. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution (red).
The negative controls are as follows:
-ve control 1: ab202896 at 1/500 dilution followed by ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/1000 dilution.
-ve control 2: ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution followed by ab150077 (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MYBBP1A antibody [EPR7205] (ab202896)

MYBBP1A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MYBBP1A antibody

Immunohistochemical analysis of paraffin-embedded human cervix carcinoma tissue labeling MYBBP1A with ab202896 at 1/100 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution. Cytoplasmic and nuclear staining on Human cervix carcinoma tissue is observed. Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Western blot - Anti-MYBBP1A antibody [EPR7205] (ab202896)

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-MYBBP1A antibody [EPR7205] (ab202896) at 1/1000 dilution

All lanes:
HeLa (Human epithelial cells from cervix adenocarcinoma) cell lysate at 10 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 148 kDa

Observed band size: 149 kDa

Exposure time: 5s



Western blot - Anti-MYBBP1A antibody [EPR7205] (ab202896)

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-MYBBP1A antibody [EPR7205] (ab202896) at 1/1000 dilution

All lanes:
HEK-293 (Human epithelial cells from embryonic kidney) cell lysate at 10 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 148 kDa

Observed band size: 149 kDa

Exposure time: 15s



Western blot - Anti-MYBBP1A antibody [EPR7205] (ab202896)

MYBBP1A Western blot staining of Jurkat (Human T cell leukemia cells from peripheral blood) cell lysate using rabbit Anti-MYBBP1A antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-MYBBP1A antibody [EPR7205] (ab202896) at 1/1000 dilution

All lanes:
Jurkat (Human T cell leukemia cells from peripheral blood) cell lysate at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 148 kDa

Observed band size: 149 kDa

Exposure time: 3min

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.