abcam

ab203085

Anti-Hsp90 beta antibody [EPR16621]

Anti-Hsp90 beta antibody [EPR16621] (ab203085) is a rabbit monoclonal antibody detecting Hsp90 beta in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat,**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 10 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR16621
Purification technique	Affinity purification Protein A
Concentration	1.729 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Recombinant fragment
Dilution info	1/70
Notes	-

Species	Human
Dilution info	1/70
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Recombinant fragment
Dilution info	1/5000
Notes	-

Species	Mouse
Dilution info	1/5000
Notes	-

Species	Rat
Dilution info	1/5000
Notes	-

Species	Human
Dilution info	1/5000
Notes	-

ICC/IF

Tested

Species	Recombinant fragment
Dilution info	1/250
Notes	-

Species	Human
Dilution info	1/250
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Recombinant fragment
Dilution info	1/100
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Species	Human
Dilution info	1/100
Notes	ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Recombinant fragment
Dilution info	1/500
Notes	-

Species	Mouse
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Human
Dilution info	1/500
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information HSP90AB1

Function

Molecular chaperone that promotes the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. Undergoes a functional cycle linked to its ATPase activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. Interacts dynamically with various co-chaperones that modulate its substrate recognition, ATPase cycle and chaperone function (PubMed:16478993, PubMed:19696785). Engages with a range of client protein classes via its interaction with various co-chaperone proteins or complexes, that act as adapters, simultaneously able to interact with the specific client and the central chaperone itself. Recruitment of ATP and co-chaperone followed by client protein forms a functional chaperone. After the completion of the chaperoning process, properly folded client protein and co-chaperone leave HSP90 in an ADP-bound partially open conformation and finally, ADP is released from HSP90 which acquires an open conformation for the next cycle (PubMed:26991466, PubMed:27295069). Apart from its chaperone activity, it also plays a role in the regulation of the transcription machinery. HSP90 and its co-chaperones modulate transcription at least at three different levels. They first alter the steady-state levels of certain transcription factors in response to various physiological cues.

Second, they modulate the activity of certain epigenetic modifiers, such as histone deacetylases or DNA methyl transferases, and thereby respond to the change in the environment. Third, they participate in the eviction of histones from the promoter region of certain genes and thereby turn on gene expression (PubMed:25973397). Antagonizes STUB1-mediated inhibition of TGF-beta signaling via inhibition of STUB1-mediated SMAD3 ubiquitination and degradation (PubMed:24613385). Promotes cell differentiation by chaperoning BIRC2 and thereby protecting from auto-ubiquitination and degradation by the proteasomal machinery (PubMed:18239673). Main chaperone involved in the phosphorylation/activation of the STAT1 by chaperoning both JAK2 and PRKCE under heat shock and in turn, activates its own transcription (PubMed:20353823). Involved in the translocation into ERGIC (endoplasmic reticulum-Golgi intermediate compartment) of leaderless cargos (lacking the secretion signal sequence) such as the interleukin 1/IL-1; the translocation process is mediated by the cargo receptor TMED10 (PubMed:32272059). (Microbial infection) Binding to N.meningitidis NadA stimulates monocytes (PubMed:21949862). Seems to interfere with N.meningitidis NadA-mediated invasion of human cells (Probable).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-Hsp90 beta antibody [EPR16621] (ab203085) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat, samples.

What is the molecular weight of Hsp90 beta?
Anti-Hsp90 beta [EPR16621] (ab203085) specifically detects a band for Hsp90 beta (UniProt: P08238) at a molecular weight of 85kDa.

Trusted by the scientific community
Anti-Hsp90 beta [EPR16621] (ab203085) was first used in a scientific publication in 2015 and has been cited over 10 times in peer-reviewed journals.

Specificity confirmed
The specificity of Anti-Hsp90 beta antibody [EPR16621] (ab203085) has been confirmed by Western blot testing in HSP90AB1 Knockout HEK293T cell line, ab266117.

Other related products
We have a range of other formats of antibody clone [EPR16621] also available for your convenience: ab32568, Carrier free - ab247273, Alexa Fluor® 488 - ab202822, Alexa Fluor® 647 - ab202823

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Hsp90 beta also known as Hsp90AB1 or Hsp90 protein is a heat shock protein of approximately 90 kDa. It is a molecular chaperone found in most eukaryotic cells. Hsp90 beta helps in the proper folding stabilization and degradation of many proteins. Unlike its isoform Hsp90 alpha Hsp90 beta has a more stable expression and is not typically induced by stress. This protein is primarily localized in the cytosol but can also be present in other cellular compartments depending on the cellular state.
Biological function summary	Hsp90 beta functions to maintain protein homeostasis and cellular integrity. It forms part of a multi-protein chaperone complex which includes cochaperones such as Hop Hsp70 and p23 necessary for its full functionality. Hsp90 beta supports the maturation of steroid hormone receptors kinases and other client proteins. It plays an important role in the cell cycle regulation through its interaction with various proteins ensuring proper cell division and growth.
Pathways	Hsp90 beta is deeply involved in signal transduction and cellular stress response pathways. Its interaction with the Akt pathway is significant for cell survival signals. Hsp90 beta also participates in the MAP kinase pathway affecting cell growth and differentiation. The protein associates with multiple kinases including RAF and Src which are important for downstream signaling.

Associated diseases and disorders	The dysregulation of Hsp90 beta is associated with cancer and neurodegenerative diseases. In cancer Hsp90 beta stabilizes many oncoproteins making it a potential therapeutic target for inhibiting tumor growth. It interacts with client proteins like the proto-oncogene c-Src promoting tumorigenesis. In neurodegenerative disorders such as Alzheimer's disease improper interaction between Hsp90 beta and tau proteins can contribute to disease progression impacting neuronal function.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

11 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Hsp90 beta Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Hsp90 beta antibody

Immunohistochemical analysis of paraffin-embedded Human cerebral cortex tissue labeling Hsp90 beta with ab203085 at 1/500 dilution followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasm and nucleus staining on neuron of cerebral cortex is observed. This protein is synthesized in cytoplasm, but it can shuttle between cytoplasm and nucleus depend on the cell status. Counter stained with hematoxylin. Negative control also shown using PBS instead of primary antibody and secondary as above.


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Hsp90 beta Flow Cytometry (Intracellular) staining using rabbit Anti-Hsp90 beta antibody


Intracellular Flow Cytometry analysis of 2% paraformaldehyde fixedHeLa (Human epithelial cells from cervix adenocarcinoma) cells using ab203085 at a 1/100 dilution (red) or a Rabbit monoclonal IgG (negative) (black). Goat anti rabbit IgG (FITC) secondary used at a 1/150 dilution.



Immunocytochemistry/ Immunofluorescence - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Hsp90 beta Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Hsp90 beta antibody

Immunofluorescence analysis of 4% paraformaldehyde fixed HeLa cells (Human epithelial cells from cervix adenocarcinoma) labeling Hsp90 beta with ab203085 at a 1/250 dilution showing cytoplasmic staining. Permeabilisation using 0.1% tritonX-100.
Secondary ab: Anti-Rabbit Alexa Fluor® 488 (ab150077) at 1/200 dilution. Counter stain is labeling tubulin (red) with ab7291 at 1/500 dilution with secondary antibody Anti-Mouse AlexaFluor® 594 (ab150120) at 1/400 dilution. DAPI stains the nucleus in blue. -ve control 1 is ab178854 at 1/250 dilution, Anti-Mouse AlexaFluor® 594 (ab150120) at 1/400 dilution. -ve control 2 is ab7291 at 1/500 dilution, Anti-Rabbit Alexa Fluor® 488 (ab150077) at 1/200 dilution.



Immunoprecipitation - Anti-Hsp90 beta antibody [EPR16621] (ab203085)


Hsp90 beta was immunoprecipitated from 1mg of Hela (Human epithelial cells from cervix adenocarcinoma) whole cell extract with ab203085 at 1/70 dilution. Western blot was performed of the immunoprecipitate using ab203085 at 1/5000 dilution. Goat Anti-Rabbit IgG, (H+L), peroxidase conjugated, was used as secondary antibody at 1:1000 dilution.
Blocking buffer and concentration: 5% NFDM/TBST.

Diluting buffer and concentration: 5% NFDM/TBST.

All lanes:
Immunoprecipitation - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Predicted band size: 83 kDa


Observed band size: 90 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Hsp90 beta Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Hsp90 beta antibody


Immunohistochemical analysis of paraffin-embedded Mouse cerebral cortex tissue labeling Hsp90 beta with ab203085 at 1/500 dilution followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasm and nucleus staining on neuron of cerebral cortex is observed. This protein is synthesized in cytoplasm, but it can shuttle between cytoplasm and nucleus depending on the cell status. Counter stained with hematoxylin. Negative control also shown using PBS instead of primary antibody and secondary as above.
Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Hsp90 beta Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Hsp90 beta antibody

Immunohistochemical analysis of paraffin-embedded Rat cerebral cortex tissue labeling Hsp90 beta with ab203085 at 1/500 dilution followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasm and nucleus staining on neuron of cerebral cortex is observed. This protein is synthesized in cytoplasm, but it can shuttle between cytoplasm and nucleus depending on the cell status. Counter stained with hematoxylin. Negative control also shown using PBS instead of primary antibody and secondary as above.
Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Western blot - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Hsp90 beta Western blot staining using rabbit Anti-Hsp90 beta antibody

Blocking and Diluting buffer and concentration: 5%
NFDM/TBST.
ab203085 only recognizes Hsp90 beta based on recombinant protein WB result.

All lanes:
Western blot - Anti-Hsp90 beta antibody [EPR16621] (ab203085) at 1/10000 dilution

Lane 1:
Hsp90 alpha recombinant protein fragment (GST-tag): aa533-732 at 0.01 µg


Lane 2:
Hsp90 beta recombinant protein fragment (His-Tag®): aa525-724 at 0.01 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/1000 dilution

Predicted band size: 83 kDa

Exposure time: 3min



Western blot - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Hsp90 beta Western blot staining using rabbit Anti-Hsp90 beta antibody

Lanes 1-4: Merged signal (red and green). Green - ab203085 observed at 90 kDa. Red - loading control ab8245 observed at 36 kDa.
ab203085 Anti-Hsp90 beta antibody [EPR16621] was shown to specifically react with Hsp90 beta in wild-type HEK293T cells. Loss of signal was observed when knockout cell line ab266117 (knockout cell lysate ab257190) was used. Wild-type and Hsp90 beta knockout samples were subjected to SDS-PAGE. ab203085 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) were incubated at room temperature for 2.5 hours at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-Hsp90 beta antibody [EPR16621] (ab203085) at 1/1000 dilution

Lane 1:
Wild-type HEK293T cell lysate at 20 µg

Lane 2:
Western blot - Human HSP90AB1 (Hsp90 beta) knockout HEK-293T cell lysate (ab257190)

Lane 3:
Jurkat cell lysate at 20 µg

Lane 4:
SH-SY5Y cell lysate at 20 µg


Secondary

All lanes:

Western blot - Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) at 1/10000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa



Western blot - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Hsp90 beta Western blot staining using rabbit Anti-Hsp90 beta antibody

Lanes 1 - 4: Merged signal (red and green). Green - ab203085 observed at 85 kDa. Red - loading control ab8245 (Mouse anti-GAPDH antibody [6C5]) observed at 37kDa.
ab203085 was shown to react with Hsp90 beta in wild-type HEK-293T cells in western blot with loss of signal observed in HSP90AB1 knockout cell line ab266117 (HSP90AB1 knockout cell lysate ab257190). Wild-type and HSP90AB1 knockout HEK-293T cell lysates were subjected to SDS-PAGE. Membranes were blocked in 3% milk in TBS-T (0.1% Tween®) before incubation with ab203085 and ab8245 (Mouse anti-GAPDH antibody [6C5]) overnight at 4°C at a 1 in 5000 dilution and a 1 in 20000 dilution respectively. Blots were incubated with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-Hsp90 beta antibody [EPR16621] (ab203085) at 1/5000 dilution

Lane 1:
Wild-type HEK-293T cell lysate at 20 µg

Lane 2:
Western blot - Human HSP90AB1 (Hsp90 beta) knockout HEK-293T cell lysate (ab257190) at 20 µg


Lane 3:
Saos-2 cell lysate at 20 µg

Lane 4:
HL-60 cell lysate at 20 µg

Performed under reducing conditions.

Predicted band size: 83 kDa

Observed band size: 85 kDa



Western blot - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Hsp90 beta Western blot staining using rabbit Anti-Hsp90 beta antibody

Blocking buffer and concentration: 5% NFDM/TBST
Diluting buffer and concentration: 5% NFDM /TBST

All lanes:
Western blot - Anti-Hsp90 beta antibody [EPR16621] (ab203085) at 1/20000 dilution

Lane 1:
HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell lysates at 20 µg

Lane 2:
293T (Human epithelial cells from embryonic kidney) whole cell lysates at 20 µg

Lane 3:
K562 (Human chronic myelogenous leukemia cells from bone marrow) whole cell lysates at 20 µg


Lane 4:
Jurkat (Human T cell leukemia cells from peripheral blood) whole cell lysates at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa



Western blot - Anti-Hsp90 beta antibody [EPR16621] (ab203085)

Hsp90 beta Western blot staining using rabbit Anti-Hsp90 beta antibody

Blocking buffer and concentration: 5% NFDM/TBST
Diluting buffer and concentration: 5% NFDM /TBST

All lanes:
Western blot - Anti-Hsp90 beta antibody [EPR16621] (ab203085) at 1/5000 dilution

Lane 1:
Mouse brain at 10 µg

Lane 2:
Mouse heart at 10 µg

Lane 3:
Mouse kidney at 10 µg

Lane 4:
Mouse spleen at 10 µg

Lane 5:
Rat brain at 10 µg

Lane 6:
Rat heart at 10 µg

Lane 7:
Rat kidney at 10 µg

Lane 8:
Rat spleen at 10 µg

Lane 9:
C6 (Rat glial tumor cells) whole cell lysate at 10 µg

Lane 10:
RAW 264.7 (Mouse macrophage cells transformed with Abelson murine leukemia virus) whole cell lysate at 10 µg

Lane 11:
PC-12 (Rat adrenal gland pheochromocytoma) whole cell lysate at 10 µg

Lane 12:
NIH/3T3 (Mouse embyro fibroblast cells) whole cell lysate at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 83 kDa

Observed band size: 90 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.