abcam

ab208020

Anti-Mre11 antibody [EPR21027] - ChIP Grade

Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020) is a rabbit monoclonal antibody detecting Mre11 in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ChIP**. Suitable for **Human, Mouse, Rat**.

- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR21027
Purification technique	Affinity purification Protein A
Concentration	0.634 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Mouse
Dilution info	1/2000

Notes	Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

Species	Human
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval before commencing with IHC staining protocol.

ChIP

Tested

Species	Human
Dilution info	-
Notes	5µg

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Rat
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000

Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information MRE11

Function

Core component of the MRN complex, which plays a central role in double-strand break (DSB) repair, DNA recombination, maintenance of telomere integrity and meiosis (PubMed:11741547, PubMed:14657032, PubMed:22078559, PubMed:23080121, PubMed:24316220, PubMed:26240375, PubMed:27889449, PubMed:28867292, PubMed:29670289, PubMed:30464262, PubMed:30612738, PubMed:31353207, PubMed:37696958, PubMed:38128537, PubMed:9590181, PubMed:9651580, PubMed:9705271). The MRN complex is involved in the repair of DNA double-strand breaks (DSBs) via homologous recombination (HR), an error-free mechanism which primarily occurs during S and G2 phases (PubMed:24316220, PubMed:28867292, PubMed:31353207, PubMed:38128537). The complex (1) mediates the end resection of damaged DNA, which generates proper single-stranded DNA, a key initial steps in HR, and is (2) required for the recruitment of other repair factors and efficient activation of ATM and ATR upon DNA damage (PubMed:24316220, PubMed:27889449, PubMed:28867292, PubMed:36050397, PubMed:38128537). Within the MRN complex, MRE11 possesses both single-strand endonuclease activity and double-strand-specific 3'-5'

exonuclease activity (PubMed:11741547, PubMed:22078559, PubMed:24316220, PubMed:26240375, PubMed:27889449, PubMed:29670289, PubMed:31353207, PubMed:36563124, PubMed:9590181, PubMed:9651580, PubMed:9705271). After DSBs, MRE11 is loaded onto DSBs sites and cleaves DNA by cooperating with RBBP8/CtIP to initiate end resection (PubMed:27814491, PubMed:27889449, PubMed:30787182). MRE11 first endonucleolytically cleaves the 5' strand at DNA DSB ends to prevent non-homologous end joining (NHEJ) and licence HR (PubMed:24316220). It then generates a single-stranded DNA gap via 3' to 5' exonucleolytic degradation to create entry sites for EXO1- and DNA2-mediated 5' to 3' long-range resection, which is required for single-strand invasion and recombination (PubMed:24316220, PubMed:28867292). RBBP8/CtIP specifically promotes the endonuclease activity of MRE11 to clear protein-DNA adducts and generate clean double-strand break ends (PubMed:27814491, PubMed:27889449, PubMed:30787182). MRE11 endonuclease activity is also enhanced by AGER/RAGE (By similarity). The MRN complex is also required for DNA damage signaling via activation of the ATM and ATR kinases: the nuclease activity of MRE11 is not required to activate ATM and ATR (PubMed:14657032, PubMed:15064416, PubMed:15790808, PubMed:16622404). The MRN complex is also required for the processing of R-loops (PubMed:31537797). The MRN complex is involved in the activation of the cGAS-STING pathway induced by DNA damage during tumorigenesis: the MRN complex acts by displacing CGAS from nucleosome sequestration, thereby activating it (By similarity). In telomeres the MRN complex may modulate t-loop formation (PubMed:10888888). MRE11 contains two DNA-binding domains (DBDs), enabling it to bind both single-stranded DNA (ssDNA) and double-stranded DNA (dsDNA).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?

Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), ChIP in Human, Mouse, Rat samples.

What is the molecular weight of Mre11?
Anti-Mre11 [EPR21027] - ChIP Grade (ab208020) specifically detects a band for Mre11 (UniProt: P49959) at a molecular weight of 81kDa.

Other related products
We have a range of other formats of antibody clone [EPR21027] also available for your convenience: ab208020, Carrier free - ab230381

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The Mre11 protein also known as MRE11A serves as an integral part of the DNA repair machinery in cells. It forms a component of the MRE11/RAD50/NBS1 (MRN) complex which is essential for the detection and repair of DNA double-strand breaks (DSBs). The molecular weight of the Mre11 protein is approximately 81 kDa. It is broadly expressed in various human tissues highlighting its extensive role in maintaining genomic stability.
Biological function summary	This protein acts in the repair of DSBs by initiating homologous recombination and non-homologous end joining pathways. Together with the RAD50 and NBS1 proteins Mre11 forms the MRN complex which processes DNA ends and signals to other repair mechanisms. Additionally Mre11's exonuclease and endonuclease activities are important for the resection of DNA at break sites facilitating subsequent repair synthesis.
Pathways	Mre11 is instrumental in the DNA damage response and maintenance of genomic integrity. It operates within the ATM (ataxia-telangiectasia mutated) signaling pathway which activates upon DNA damage and regulates cell cycle checkpoints. Mre11 interacts with the ATM protein modifying the cellular response to DNA damage. The complex also collaborates closely with BRCA1 an important regulator of the repair process and associated with preventing breast cancer development.

Associated diseases and disorders	Mutations or dysfunction in the MRE11A gene can be linked to several conditions including ataxia-telangiectasia-like disorder (ATLD) and Nijmegen breakage syndrome (NBS). These disorders result from impaired DNA repair mechanisms leading to increased sensitivity to radiation and predisposition to cancer. The NBS1 protein as part of the MRN complex works closely with Mre11 in these conditions. Both disorders highlight the critical role of Mre11 in safeguarding genomic stability and preventing disease.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


10 product images



Flow Cytometry (Intracellular) - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020)

Mre11 Flow Cytometry (Intracellular) staining using rabbit Anti-Mre11 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol permeabilized K562 (Human human chronic myelogenous leukemia cell line from bone marrow) cell line labeling Mre11 with ab208020 at 1/500 dilution (red) compared with a Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) (black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020)

Mre11 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Mre11 antibody


Immunohistochemical analysis of paraffin-embedded human endometrial carcinoma tissue labeling Mre11 with ab208020 at 1/2000 dilution, followed by Rabbit specific IHC polymer detection kit HRP/DAB (ab209101), ready to use. Nuclear staining in human endometrial carcinoma is observed (PMID: 24927325). Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Rabbit specific IHC polymer detection kit HRP/DAB (ab209101), ready to use.

Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020)

Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue labeling Mre11 with ab208020 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP polymer) (ab214880) Ready to use. Counter stained with Hematoxylin. Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody Goat Anti-Rabbit IgG H&L (HRP polymer) (ab214880) Heat mediated antigen retrieval was performed using ab93684 (Tris/EDTA buffer, pH 9.0) Positive staining on human breast carcinoma. The section was incubated with ab208020 at 4°C overnight.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020)

Mre11 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Mre11 antibody


Immunohistochemical analysis of paraffin-embedded human colon tissue labeling Mre11 with ab208020 at 1/2000 dilution, followed by Rabbit specific IHC polymer detection kit HRP/DAB (ab209101), ready to use. Nuclear staining in human colon tissue is observed (PMID: 25447316). Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Rabbit specific IHC polymer detection kit HRP/DAB (ab209101), ready to use.

Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



ChIP - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020)

Chromatin was prepared from non-treated HeLa cells or HeLa cells treated with 1% Methyl methanesulfonate for 1 hour according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 25 µg of chromatin, 5 µg of ab208020 (blue), and 20 µL of protein A/G sepharose beads slurry (10 µL of sepharose A beads + 10 µL of sepharose G beads). 5 µg of rabbit normal IgG was added to the beads control (yellow). The immunoprecipitated DNA was quantified by real time PCR (Sybr green approach).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020)

Mre11 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Mre11 antibody

Immunohistochemical analysis of paraffin-embedded rat colon tissue labeling Mre11 with ab208020 at 1/2000 dilution, followed by Rabbit specific IHC polymer detection kit HRP/DAB (ab209101), ready to use. Nuclear staining in rat colon tissue is observed (PMID: 28357369). Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Rabbit specific IHC polymer detection kit HRP/DAB (ab209101), ready to use.

Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020)

Mre11 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Mre11 antibody

Immunohistochemical analysis of paraffin-embedded mouse testis tissue labeling Mre11 with ab208020 at 1/2000 dilution, followed by Rabbit specific IHC polymer detection kit HRP/DAB (ab209101), ready to use. Nuclear staining in mouse testis is observed (PMID: 10508516). Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Rabbit specific IHC polymer detection kit HRP/DAB (ab209101), ready to use.

Heat mediated antigen retrieval was performed before commencing with IHC staining protocol.



Western blot - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020)

Blocking/Dilution: 5% NFDM/TBST.
Exposure time: 3 minutes.

All lanes:
Western blot - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020) at 1/1000 dilution

All lanes:
Human fetal kidney tissue lysate at 10 µg


Secondary

All lanes:
Western blot - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/4000 dilution

Developed using the ECL technique.

Predicted band size: 80 kDa

Observed band size: 81 kDa



Western blot - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020)

Blocking/Dilution: 5% NFDM/TBST.
Exposure time: 3 minutes.

All lanes:
Western blot - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020) at 1/5000 dilution

Lane 1:
K562 (human chronic myelogenous leukemia cell line from bone marrow) whole cell lysate at 20 µg

Lane 2:
Jurkat (human T cell leukemia cell line from peripheral blood) whole cell lysate at 20 µg

Lane 3:
HEK-293T (human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate at 20 µg


Lane 4:
D3 (mouse embryo stem cell) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 80 kDa

Observed band size: 81 kDa



Western blot - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020)

Mre11 Western blot staining using rabbit Anti-Mre11 antibody

Blocking/Dilution: 5% NFDM/TBST.
Exposure time: 3 minutes.

All lanes:
Western blot - Anti-Mre11 antibody [EPR21027] - ChIP Grade (ab208020) at 1/1000 dilution

Lane 1:
C6 (rat glial tumor cell line) whole cell lysate at 10 µg

Lane 2:
RAW 264.7 (mouse macrophage cell line transformed with Abelson murine leukemia virus) whole cell lysate at 10 µg

Lane 3:
PC-12 (rat adrenal gland pheochromocytoma cell line) whole cell lysate at 10 µg

Lane 4:
NIH/3T3 (mouse embryo fibroblast cell line) whole cell lysate at 10 µg

Lane 5:
Mouse brain tissue lysate at 10 µg

Lane 6:
Rat brain tissue lysate at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 80 kDa

Observed band size: 81 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.