abcam

ab208035

Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR18841-95
Purification technique	Affinity purification Protein A
Concentration	0.675 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/40
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

WB

Tested

Species	Recombinant full length protein - Human
Dilution info	1/2000
Notes	-

Species	Mouse
Dilution info	1/2000
Notes	-

Species	Rat
Dilution info	1/2000
Notes	-

Species	Human
Dilution info	1/2000
Notes	-

ICC/IF

Tested

Species	Mouse
Dilution info	1/1000

Notes	-

Species	Human
Dilution info	1/1000
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-

Notes	-

Target data

See full target information MAPK8

Function	Serine/threonine-protein kinase involved in various processes such as cell proliferation, differentiation, migration, transformation and programmed cell death. Extracellular stimuli such as pro-inflammatory cytokines or physical stress stimulate the stress-activated protein kinase/c-Jun N-terminal kinase (SAP/JNK) signaling pathway (PubMed:28943315). In this cascade, two dual specificity kinases MAP2K4/MKK4 and MAP2K7/MKK7 phosphorylate and activate MAPK8/JNK1. In turn, MAPK8/JNK1 phosphorylates a number of transcription factors, primarily components of AP-1 such as JUN, JDP2 and ATF2 and thus regulates AP-1 transcriptional activity (PubMed:18307971). Phosphorylates the replication licensing factor CDT1, inhibiting the interaction between CDT1 and the histone H4 acetylase HBO1 to replication origins (PubMed:21856198). Loss of this interaction abrogates the acetylation required for replication initiation (PubMed:21856198). Promotes stressed cell apoptosis by phosphorylating key regulatory factors including p53/TP53 and Yes-associates protein YAP1 (PubMed:21364637). In T-cells, MAPK8 and MAPK9 are required for polarized differentiation of T-helper cells into Th1 cells. Contributes to the survival of erythroid cells by phosphorylating the antagonist of cell death BAD upon EPO stimulation (PubMed:21095239). Mediates starvation-induced BCL2 phosphorylation, BCL2 dissociation from BECN1, and thus activation of autophagy (PubMed:18570871). Phosphorylates STMN2 and hence regulates microtubule dynamics, controlling neurite elongation in cortical neurons (By similarity). In the developing brain, through its cytoplasmic activity on STMN2, negatively regulates the rate of exit from multipolar stage and of radial migration from the ventricular zone (By similarity). Phosphorylates several other substrates including heat shock factor protein 4 (HSF4), the deacetylase SIRT1, ELK1, or the E3 ligase ITCH (PubMed:16581800, PubMed:17296730, PubMed:20027304). Phosphorylates the CLOCK-BMAL1 heterodimer and plays a role in the regulation of the circadian clock (PubMed:22441692). Phosphorylates the heat shock transcription factor HSF1, suppressing HSF1-induced transcriptional activity (PubMed:10747973). Phosphorylates POU5F1, which results in the inhibition of POU5F1's transcriptional activity and enhances its proteasomal degradation (By similarity). Phosphorylates JUND and this phosphorylation is inhibited in the presence of MEN1 (PubMed:22327296). In neurons, phosphorylates SYT4 which captures neuronal dense core vesicles at synapses (By similarity). Phosphorylates EIF4ENIF1/4-ET in response to oxidative stress, promoting P-body assembly (PubMed:22966201). Phosphorylates SIRT6 in response to oxidative stress, stimulating its mono-ADP-ribosyltransferase activity (PubMed:27568560). Phosphorylates NLRP3, promoting assembly of the NLRP3 inflammasome (PubMed:28943315). Phosphorylates ALKBH5 in response to reactive oxygen species (ROS), promoting ALKBH5 sumoylation and inactivation (PubMed:34048572). JNK1 isoforms display different binding patterns: beta-1 preferentially binds to c-Jun, whereas alpha-1, alpha-2, and beta-2 have a similar low level of binding to both c-Jun or ATF2. However, there is no correlation between binding and phosphorylation, which is achieved at about the same efficiency by all isoforms.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	JNK1 JNK2 and JNK3 collectively known as c-Jun N-terminal kinases are important members of the mitogen-activated protein kinase (MAPK) family. JNK1 JNK2 and JNK3 have molecular weights of approximately 46 to 55 kDa with JNK1 being around 46 kDa JNK2 about 54 kDa and JNK3 also around 54 kDa. They are expressed differently across tissues: JNK1 and JNK2 are widely present in many tissues while JNK3 is mostly in brain heart and testis. These kinases primarily phosphorylate serine and threonine residues acting on various transcription factors including c-Jun to regulate cellular processes.
Biological function summary	These kinases function as significant regulators of cellular responses to stress and cytokines. They do not act alone but often form part of larger signaling complexes that include other MAPKs and various scaffold proteins. Their roles are substantial

	in cell differentiation proliferation apoptosis and migration. Precise regulation by JNK molecules influences these critical cellular events which play a role in maintaining homeostasis and response to external stresses.
Pathways	JNK proteins are central to the MAPK signaling pathway and are tightly linked to the MAPK/ERK pathway. Activation of the JNK pathways leads to the regulation of various other proteins including those in the apoptosis pathway such as Bcl-2 and Bax. These interactions focus on cellular stress response regulation and contribute to processes like inflammation stress-induced apoptosis and some metabolic processes.
Associated diseases and disorders	JNKs have important roles in inflammatory diseases and neurodegenerative disorders such as Alzheimer's disease and Parkinson’s disease. Their activity links to the regulation of pro-inflammatory cytokines and neuronal apoptosis. JNK1 and JNK2 are particularly connected to inflammatory responses whereas JNK3 with its expression in the brain has associations with neuronal death in neurodegenerative diseases. Modulating JNK activity may offer therapeutic approaches for these conditions targeting pathways involving proteins like TNF-alpha and Aβ-amyloid.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


10 product images



Flow Cytometry (Intracellular) - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

JNK1 + JNK2 + JNK3 Flow Cytometry (Intracellular) staining using rabbit Anti-JNK1 + JNK2 + JNK3 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed Jurkat (Human T cell leukemia cell line from peripheral blood) cell linelabeling JNK1+JNK2+JNK3 with ab208035 at 1/100 dilution (red) compared with a Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) (black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (blue).
Goat anti-Rabbit IgG (Alexa Fluor® 488) at 1/500 dilution was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

JNK1 + JNK2 + JNK3 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-JNK1 + JNK2 + JNK3 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cell line from cervix adenocarcinoma) cell line labeling JNK1+JNK2+JNK3 with ab208035 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution (green).
Confocal image showing cytoplasm and weak nuclear staining on HeLa cell line.


The nuclear counterstain is DAPI (blue).

Tubulin is detected with Anti-alpha Tubulin antibody [DM1A] - Loading Control (ab7291) at 1/1000 dilution and Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) preadsorbed (ab150120) at 1/1000 dilution (red).

The negative controls are as follows:-

-ve control 1: ab208035 at 1/1000 dilution followed by ab150120 at 1/1000 dilution.


-ve control 2: ab7291 at 1/1000 dilution followed by ab150077 at 1/1000 dilution.



Flow Cytometry (Intracellular) - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

JNK1 + JNK2 + JNK3 Flow Cytometry (Intracellular) staining using rabbit Anti-JNK1 + JNK2 + JNK3 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed HeLa (Human epithelial cell line from cervix adenocarcinoma) cell linelabeling JNK1+JNK2+JNK3with ab208035 at 1/100 dilution (red) compared with a Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) (black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (blue).
Goat anti-Rabbit IgG (Alexa Fluor® 488) at 1/500 dilution was used as the secondary antibody.



Immunoprecipitation - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

JNK1+JNK2+JNK3 was immunoprecipitated from 1mg of HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate with ab208035 at 1/40 dilution.
Western blot was performed from the immunoprecipitate using ab208035 at 1/1000 dilution.

VeriBlot for IP Detection Reagent (HRP) (ab131366), was used for detection at 1/10000 dilution.

Lane 1: HeLa whole cell lysate, 10ug (Input).

Lane 2: ab208035 IP in HeLa whole cell lysate.

Lane 3: Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) instead of ab208035 in HeLa whole cell lysate.


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 10 seconds.

All lanes:
Immunoprecipitation - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

Predicted band size: 48 kDa

Observed band size: 46-54 kDa



Immunocytochemistry/ Immunofluorescence - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

JNK1 + JNK2 + JNK3 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-JNK1 + JNK2 + JNK3 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (Mouse embryonic fibroblast cell line) cell line labeling JNK1+JNK2+JNK3 with ab208035 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution (green).
Confocal image showing cytoplasm and nuclear staining on NIH/3T3 cell line.


The nuclear counterstain is DAPI (blue).

Tubulin is detected with Anti-alpha Tubulin antibody [DM1A] - Loading Control (ab7291) at 1/1000 dilution and Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) preadsorbed (ab150120) at 1/1000 dilution (red).

The negative controls are as follows:-

-ve control 1: ab208035 at 1/1000 dilution followed by ab150120 at 1/1000 dilution.

-ve control 2: ab7291 at 1/1000 dilution followed by ab150077 at 1/1000 dilution.



Western blot - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

JNK1 + JNK2 + JNK3 Western blot staining using rabbit Anti-JNK1 + JNK2 + JNK3 antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035) at 1/2000 dilution

Lane 1:
Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate at 20 µg

Lane 2:
HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 20 µg

Lane 3:
K562 (Human chronic myelogenous leukemia cell line from bone marrow) whole cell lysate at 20 µg

Lane 4:
MCF7 (Human breast adenocarcinoma cell line) whole cell lysate at 20 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 48 kDa

Observed band size: 46-54 kDa

Exposure time: 3min



Western blot - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

JNK1 + JNK2 + JNK3 Western blot staining using rabbit Anti-JNK1 + JNK2 + JNK3 antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
Exposure time: Lane 1 & 2: 5 seconds; Lane 3: 15 seconds.

Human JNK1 full length recombinant protein contains aa1-427 with a His-Tag®. Human JNK2 full length recombinant protein contains aa1-424 with a His-Tag®. Human JNK3 full length recombinant protein contains aa1-464 with a His-Tag®.

All three recombinant human full length proteins were made in-house.

All lanes:
Western blot - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035) at 1/2000 dilution

Lane 1:
Human JNK1 full length recombinant protein at 0.01 µg


Lane 2:
Human JNK2 full length recombinant protein at 0.01 µg

Lane 3:
Human JNK3 full length recombinant protein at 0.01 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 48 kDa



Western blot - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

Blocking/Dilution buffer: 5% NFDM/TBST.
All lanes:
Western blot - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035) at 1/2000 dilution

Lane 1:
Human fetal liver lysate at 10 µg

Lane 2:
Human fetal heart lysate at 10 µg

Lane 3:
Human fetal kidney lysate at 10 µg


Secondary

All lanes:
Goat Anti-Rabbit IgG Peroxidase Conjugate, specific to the non-reduced form of IgG at 1/10000 dilution

Predicted band size: 48 kDa

Observed band size: 46-54 kDa

Exposure time: 30s



Western blot - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

Blocking and Diluting buffer and concentration: 5% NFDM/TBST
All lanes:
Western blot - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035) at 1/2000 dilution

Lane 1:
C6 (Rat glial tumor cell line) whole cell lysate at 10 µg

Lane 2:
RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) whole cell lysate at 10 µg

Lane 3:
PC-12 (Rat adrenal gland pheochromocytoma cell line) whole cell lysate at 10 µg

Lane 4:
NIH/3T3 (Mouse embryonic fibroblast cell line) whole cell lysate at 10 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 48 kDa

Observed band size: 46-54 kDa

Exposure time: 8s



Western blot - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035)

Blocking/Dilution buffer: 5% NFDM/TBST.
Exposure time: Lane 1: 4 seconds; Lane 2,3 & 4: 30 seconds; Lane 5: 15 seconds; Lane 6 & 7: 3 minutes.

All lanes:
Western blot - Anti-JNK1 + JNK2 + JNK3 antibody [EPR18841-95] (ab208035) at 1/2000 dilution

Lane 1:
Mouse brain lysate at 10 µg

Lane 2:
Mouse heart lysate at 10 µg

Lane 3:
Mouse kidney lysate at 10 µg

Lane 4:
Mouse spleen lysate at 10 µg

Lane 5:
Rat brain lysate at 10 µg

Lane 6:
Rat kidney lysate at 10 µg

Lane 7:
Rat spleen lysate at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 48 kDa

Observed band size: 46-54 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.