abcam

ab209665

Anti-Brachyury / Bry antibody [EPR18113]

Anti-Brachyury / Bry antibody [EPR18113] (ab209665) is a rabbit monoclonal antibody detecting Brachyury / Bry in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, IHC- Wmt, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- Clone EPR18113 is the most cited clone to Brachyury / Bry
- Biophysical QC for unrivalled batch-batch consistency
- Over 30 publications

Recombinant | RabMAb | Advanced Validation | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR18113
Purification technique	Affinity purification Protein A
Concentration	0.405 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/1000
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Primates, Common marmoset
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Primates, Common marmoset
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/1000
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Primates, Common marmoset
Dilution info	-
Notes	-

IHC - Wmt

Tested

Species	Mouse
Dilution info	1/100
Notes	-

Expected

Species	Human, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Primates, Common marmoset

Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/400
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Primates, Common marmoset
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/8000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/8000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Human
Dilution info	1/8000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Predicted

Species	Primates, Common marmoset
Dilution info	-
Notes	-

Target data

See full target information TBXT

Function	Involved in the transcriptional regulation of genes required for mesoderm formation and differentiation. Binds to a palindromic T site 5'-TTCACACCTAGGTGTGAA-3' DNA sequence and activates gene transcription when bound to such a site.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Anti-Brachyury / Bry antibody [EPR18113] (ab209665) was developed by Abcam using patented rabbit monoclonal antibody technology and is validated for use in Flow Cyt (Intra), ICC/IF, IHC - Wmt, IHC-P, IP, WB in human, mouse, rat samples.

Anti-Brachyury / Bry antibody [EPR18113] (ab209665) has been cited over 38 times in peer reviewed journals and is trusted by the scientific community.

Abcam's high quality manufacturing and validation processes ensure Anti-Brachyury / Bry antibody [EPR18113] (ab209665) has high sensitivity and specificity alongside high lot-to-lot consistency and reproducibility.

Anti-Brachyury / Bry antibody [EPR18113] (ab209665) has 9 independent reviews from customers.

Anti-Brachyury / Bry antibody [EPR18113] (ab209665) specifically detects Brachyury / Bry (UniProt ID: O15178; Molecular weight: 47kDa) and is sold in 100 µL and 1 mL selling sizes.

Conjugation-ready, carrier free format available for antibody clone EPR18113 - ab236023.

Antibody clone EPR18113 is also available pre-conjugated to a variety of labels for your convenience - Alexa Fluor® 488, Alexa Fluor® 555 (ab314674, ab314675).

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The Brachyury protein also known as Bry or T acts as a transcription factor and belongs to the T-box family of genes. It consists of about 435 amino acids giving it an estimated mass of around 47 kDa. Brachyury plays a critical role in the development of the posterior mesoderm and nucleus pulposus during embryogenesis. It is strongly expressed in the notochord mesoderm and some tumor tissues. The protein's presence can be detected using specific assays such as Brachyury IHC (immunohistochemistry) and Brachyury stain which help identify Brachyury marker expression in both normal and pathological states.
Biological function summary	The Brachyury protein influences the morphogenetic movement in mesoderm development. It directly interacts with specific DNA sequences to regulate the transcription of target genes. Brachyury often operates as part of a larger transcriptional complex that manages the differentiation and migration of cells during embryogenesis. The gene's expression indicates mesodermal differentiation and is important for cell fate decisions within developing embryos.

Pathways	Brachyury plays a significant role in the Wnt signaling and the TGF-beta pathways. These pathways are essential in controlling cell proliferation and differentiation. Brachyury directly impacts the regulation of these pathways influencing other proteins like beta-catenin in Wnt signaling. This interconnectedness suggests its broader role in cellular communication and development processes potentially affecting gene expression across these pathways.
Associated diseases and disorders	Brachyury has been associated with various cancers such as chordoma and other related tumors. Chordoma specifically involves the notochord where Brachyury marks the disease's presence. Other proteins like E-cadherin interact with Brachyury in tumor progression and metastasis. Understanding Brachyury's function helps in diagnosing and potentially targeting treatment options for these ailments.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

12 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Brachyury / Bry Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Brachyury / Bry antibody

Immunohistochemical analysis of paraffin-embedded Human chordoma tissue labeling Brachyury / Bry with ab209665 at 1/8000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution. Nucleus staining on Human chordoma is observed. Counter stained with Hematoxylin
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Brachyury / Bry Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Brachyury / Bry antibody

Immunohistochemical analysis of paraffin-embedded Human chondrosarcoma tissue labeling Brachyury / Bry with ab209665 at 1/8000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution. Negative staining on Human chondrosarcoma. Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Brachyury / Bry Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Brachyury / Bry antibody

Immunohistochemical analysis of paraffin-embedded Human meningioma tissue labeling Brachyury / Bry with ab209665 at 1/8000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution. Negative staining on Human meningioma. Counter stained with Hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Brachyury / Bry Immunocytochemistry/ Immunofluorescence staining of MUG-Chor1 (human sacral bone chordoma) cell using rabbit Anti-Brachyury / Bry antibody


Confocal image showing nucleus staining in MUG-Chor1 cell line (shown in green). The counterstain was observed in red. Nuclear DNA was labelled with DAPI (shown in blue). ab209665 staining Brachyury/Bry in MUG-Chor1 (human sacral bone chordoma) cells by ICC/IF (Immunocytochemistry/Immunofluorescence). Cells were fixed 4% paraformaldehyde and permeabilized with 0.1% tritonX-100. Samples were incubated with primary antibody at a 1/1000 dilution. An AlexaFluor®488 Goat anti-Rabbit (ab150077) was used as the secondary antibody at a 1/1000 dilution. An Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) (ab195889) was used as a counterstain antibody. DAPI was used as nuclear counterstain. Nuclear staining on MUG-Chor1 cells.



Flow Cytometry (Intracellular) - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Brachyury / Bry Flow Cytometry (Intracellular) staining using rabbit Anti-Brachyury / Bry antibody

ab209665 staining Brachyury/Bry in MUG-Chor1 (Human sacral bone chordoma) cell line by intracellular flow cytometry. Cells were fixed with 4% paraformaldehyde and 90% methanol. Sample was incubated with primary antibody at 1/400 dilution (red). A Goat anti-rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution. Rabbit monoclonal IgG (ab172730) (Black) was used as an isotype control. Cell without incubation with primary antibody and secondary antibody (Blue).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Brachyury / Bry Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Brachyury / Bry antibody

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling Brachyury / Bry with ab209665 at 1/8000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution. Negative staining on Human kidney. Counter stained with Hematoxylin. Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunoprecipitation - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Lane 1 (input): MUG-Chor1 (human sacral bone chordoma) whole cell lysate, 10 µg
Lane 2 (+): MUG-Chor1 whole cell lysate
Lane 3 (-): Rabbit monoclonal IgG (ab172730) instead of ab209665 in MUG-Chor1 whole cell lysate ab209665 immunoprecipitating Brachyury/Bry in MUG-Chor1 lysates. For western blotting, primary antibody used was ab209665 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)

(ab131366), was used for detection at 1/10,000 dilution. Blocking and diluting buffer used was 5% NFDM/TBST.
All lanes:
Immunoprecipitation - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Predicted band size: 47 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Immunohistochemical analysis of formalin fixed paraffin embedded mouse embryo E14 labelling Brachyury/Bry with ab209665 at a concentration of 4µg/ml. The immunostaining was performed on a Ventana DISCOVERY ULTRA (Roche Tissue Diagnostics) instrument with a ChromoMap DAB IHC Kit with anti-rabbit HQ, anti-HQ HRP detection. Heat mediated antigen retrieval was performed with DISCOVERY cell conditioning solution (CC1) 100°C, pH8.5 for 64mins.

Anti-Brachyury/Bry antibody [EPR18113] ab209665 was incubated for 16mins at 37°C. Sections were counterstained with Hematoxylin II. Image inset shows absence of staining in secondary antibody only control.


Customers are encouraged to optimise antigen retrieval conditions, antibody concentration, incubation times and temperature for best results in their own IHC assay workflow (automated and manual).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Brachyury / Bry Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Brachyury / Bry antibody


ab209665 staining Brachyury/Bry in Rat E14.5 embryo tissue sections by Immunohistochemistry (IHC-P). Tissue was fixed paraffin and antigen retrieval was performed by heat mediation using ab93684 (Tris/EDTA buffer, pH 9.0). Samples were incubated with primary antibody at a 1/4000 dilution. A ready to use Goat anti-rabbit IgG H&L (HRP) was used as the secondary antibody. Hematoxylin was used as a counter-stain. Nuclear staining on notochord of rat E14.5 embryo.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Brachyury / Bry Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Brachyury / Bry antibody

ab209665 staining Brachyury/Bry in mouse E14.5 embryo tissue sections by Immunohistochemistry (IHC-P). Tissue was fixed paraffin and antigen retrieval was performed by heat mediation using ab93684 (Tris/EDTA buffer, pH 9.0). Samples were incubated with primary antibody at a 1/4000 dilution. A ready to use Goat anti-rabbit IgG H&L (HRP) was used as the secondary antibody. Hematoxylin was used as a counter-stain. Nuclear staining on notochord of mouse E14.5 embryo.



Western blot - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

Brachyury / Bry Western blot staining of MUG-Chor1 (human sacral bone chordoma), whole cell lysate using rabbit Anti-Brachyury / Bry antibody

Blocking and diluting buffer: 5% NFDM/TBST
All lanes:
Western blot - Anti-Brachyury / Bry antibody [EPR18113] (ab209665) at 1/1000 dilution

All lanes:
MUG-Chor1 (human sacral bone chordoma), whole cell lysate at 10 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 47 kDa

Observed band size: 43 kDa, 49 kDa

Exposure time: 5s



This image is courtesy of a customer review by Dr Sophie Morgani

IHC - Wholemount - Anti-Brachyury / Bry antibody [EPR18113] (ab209665)

IHC - Wholemount of E6.5 wholemount mouse embryo labelling Brachyury / Bry with ab209665. Sample was incubated with primary antibody (1/100 in PBS/0.1% Triton-X/5% donkey serum/1% BSA) for 18 hours at 4°C. An Alexa Fluor® 488-conjugated donkey anti-rabbit IgG monoclonal was used as the secondary antibody (1/500).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.