abcam

ab210322

FITC Anti-CD4 antibody [OKT4]

Anti-CD4 antibody [OKT4] conjugated to FITC - suitable for flow cytometry, and reacts with human samples. Clone [OKT4] is a mouse monoclonal antibody and one of the most commonly used clones for flow cytometry.

Design your panel with our interactive flow cytometry panel builder. Click here.

Lab Essentials

Key facts

Isotype	IgG2b
Host species	Mouse
Conjugation	FITC
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	pH: 7.2 Preservative: 0.09% Sodium azide Constituents: 0.87% Sodium chloride, 0.12% Sodium dihydrogen phosphate, 0.1% Gelatin
Form	Liquid
Clonality	Monoclonal
Clone number	OKT4
Purification technique	Affinity purification Protein A
Light chain type	kappa
Purification notes	ab210322 was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation.

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	5 µL for 10^5-10^8 Cells
Notes	The antibody has been diluted for use at 5 µl (0.25 µg) per test, defined as the amount of antibody that will stain a cell sample in a final volume of approximately 100 µl.

Predicted

Species	Chimpanzee, Cynomolgus monkey, Rhesus monkey
Dilution info	-
Notes	-

Target data

See full target information CD4

Function

Integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class II molecule:peptide complex. The antigens presented by class II peptides are derived from extracellular proteins while class I peptides are derived from cytosolic proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class II presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion and activation of T-helper cells. In other cells such as macrophages or NK cells, plays a role in differentiation/activation, cytokine expression and cell migration in a TCR/LCK-independent pathway. Participates in the development of T-helper cells in the thymus and triggers the differentiation of monocytes into functional mature macrophages. (Microbial infection) Primary receptor for human immunodeficiency virus-1 (HIV-1) (PubMed:12089508, PubMed:16331979, PubMed:2214026, PubMed:9641677). Down-regulated by HIV-1 Vpu (PubMed:17346169). Acts as a receptor for Human Herpes virus 7/HHV-7 (PubMed:7909607).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	+4°C
Storage information	Store in the dark

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Flow Cytometry - FITC Anti-CD4 antibody [OKT4] (ab210322)

CD4 Flow Cytometry staining using mouse Anti-CD4 antibody

Flow cytometric analysis of Human peripheral blood lymphocytes labeling CD4 with 5 µl (0.25 µg) ab210322 (solid line), compared with 0.5 µg FITC Mouse IgG2b isotype control (dashed line).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.