abcam

ab211493

Anti-MEF2C antibody [EPR19089-202] - ChIP Grade

Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493) is a rabbit monoclonal antibody detecting MEF2C in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF, ChIP**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 10 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR19089-202
Purification technique	Affinity purification Protein A
Concentration	0.535 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ChIP

Tested

Species	Human
Dilution info	5 µg for 25 µg chromatin
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Common marmoset
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Rat
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

Predicted

Species	Common marmoset
Dilution info	-

Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Common marmoset
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Common marmoset
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	-
Notes	Use at 1:500 (human), 1:1000 (mouse, rat) dilution. Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	-
Notes	Use at 1:500 (human), 1:1000 (mouse, rat) dilution. Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Human
Dilution info	-
Notes	Use at 1:500 (human), 1:1000 (mouse, rat) dilution. Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Predicted

Species	Common marmoset
Dilution info	-
Notes	-

Target data

See full target information MEF2C

Function	Transcription activator which binds specifically to the MEF2 element present in the regulatory regions of many muscle-specific genes. Controls cardiac

morphogenesis and myogenesis, and is also involved in vascular development. Enhances transcriptional activation mediated by SOX18. Plays an essential role in hippocampal-dependent learning and memory by suppressing the number of excitatory synapses and thus regulating basal and evoked synaptic transmission. Crucial for normal neuronal development, distribution, and electrical activity in the neocortex. Necessary for proper development of megakaryocytes and platelets and for bone marrow B-lymphopoiesis. Required for B-cell survival and proliferation in response to BCR stimulation, efficient IgG1 antibody responses to T-cell-dependent antigens and for normal induction of germinal center B-cells. May also be involved in neurogenesis and in the development of cortical architecture (By similarity). Isoforms that lack the repressor domain are more active than isoform 1.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF), ChIP in Human, Mouse, Rat samples.

What is the molecular weight of MEF2C?
Anti-MEF2C [EPR19089-202] - ChIP Grade (ab211493) specifically detects a band for MEF2C (UniProt: Q06413) at a molecular weight of 51kDa.

Trusted by the scientific community
Anti-MEF2C [EPR19089-202] - ChIP Grade (ab211493) was first used in a scientific publication in 2018 and has been cited over 10 times in peer-reviewed journals.

Reviewed by scientists
Anti-MEF2C [EPR19089-202] - ChIP Grade (ab211493) has over 5 independent reviews from customers.

Specificity confirmed

The specificity of Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493) has been confirmed by Western blot testing in MEF2C Knockout THP-1 cell line, ab313739.

Other related products
We have a range of other formats of antibody clone [EPR19089-202] also available for your convenience: ab211493, Carrier free - ab231859, ab309199, Alexa Fluor® 488 - ab319010

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The MEF2C protein also known as MADS box transcription enhancer factor 2 polypeptide C plays an important role in gene expression regulation. This protein has a mass of approximately 52 kDa and is mainly found in the heart skeletal muscle brain and endothelial cells. It is part of the MEF2 family which includes other members like MEF2A MEF2B and MEF2D. MEF2C acts as a transcription factor binding to specific DNA sequences to activate the transcription of target genes involved in various biological processes.
Biological function summary	The MEF2C protein is essential in cellular differentiation and development. MEF2C is not part of a complex but interacts with other cofactors and regulatory proteins to fulfill its roles. It is important in the development of cardiac and skeletal muscle cells and plays a significant role in neurogenesis and neuronal connectivity in the central nervous system. This protein's precise regulation ensures proper development and function of these tissues and organs.
Pathways	MEF2C influences cellular signaling through the MAPK/ERK and calcium/calmodulin-dependent protein kinase pathways. In the MAPK/ERK pathway MEF2C modulates gene expression in response to external stimuli interacting with proteins like ERK1/2. In the calcium/calmodulin-dependent kinase pathway MEF2C plays a role in calcium signaling networks impacting neuronal plasticity and survival. These pathways highlight MEF2C's involvement in muscle and brain function.

Associated diseases and disorders	MEF2C mutations or dysregulation link to conditions like dilated cardiomyopathy and neurodevelopmental disorders such as MEF2C haploinsufficiency syndrome. In cardiac disease abnormal MEF2C activity can affect heart muscle cell function often involving proteins like actin and myosin. In neurodevelopmental disorders MEF2C's interaction with various transcriptional regulators impacts neuronal networks contributing to symptoms like impaired cognitive development and autism spectrum-like behaviors.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

11 product images




Immunocytochemistry/ Immunofluorescence - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)

MEF2C Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-MEF2C antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilised Raji (human Burkitt's lymphoma B lymphocyte) cells labelling MEF2C with ab211493 at 1/500 dilution, followed by AlexaFluor®488 Goat anti-Rabbit secondary (ab150077) at 1/1000 dilution (green). Confocal image showing nuclear staining in Raji cell line. **Negative control:** Jurkat (PMID: 27876533). DAPI was used as the nuclear counterstain, and the Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) (ab195889 antibody was used as a counterstain at 1/200 dilution.


The negetive controls are as follows:
-ve control 1: ab197070 on jurkat (human T cell leukemia cell line from peripheral blood) cells.
-ve control 2: Jurkat cells stained with DAPI.
-ve control 3: Merged negetive contol images.



Flow Cytometry (Intracellular) - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)

MEF2C Flow Cytometry (Intracellular) staining using rabbit Anti-MEF2C antibody


Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol permeabilsed Jurkat (Human T cell leukemia T lymphocyte, Left) / Raji (Human Burkitt's lymphoma B lymphocyte, Right) cell lines labelling MEF2C with ab211493 at 1/500 dilution (Red) compared with the isotype control Rabbit monoclonal IgG (ab172730) (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG Alexa Fluor® 488 (ab150077), at 1/2000 dilution was used as the secondary antibody.
Negative control: Jurkat (PMID: 27876533).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)

MEF2C Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MEF2C antibody


Immunohistochemical analysis of paraffin-embedded human skeletal muscle tissue labelling MEF2C with ab211493 at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Nuclear staining in human skeletal muscle cells is observed(PMID: 8506376, PMID 15703219). Counterstained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP)ready to use.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)


MEF2C Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MEF2C antibody

Immunohistochemical analysis of paraffin-embedded human endometrial carcinoma tissue labelling MEF2C with ab211493 at 1/500 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Nuclear staining in leukocytes but not in tumor cells of human endometrial carcinoma is observed (PMID: 8506376, PMID 15703219). Counterstained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP)ready to use.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



ChIP - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)


Chromatin was prepared from HUVEC cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with: 25 µg of chromatin, 5 µg of ab211493 (red), and 20 µl of protein A/G sepharose beads slurry (10 µl of sepharose A beads + 10 µl of sepharose G beads). 5 µg of rabbit normal IgG was added to the beads control (grey). The immunoprecipitated DNA was quantified by real time PCR (SYBR green chemistry).
ChIP was performed according to the literature (PMID: 26923194).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)


MEF2C Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-MEF2C antibody

Immunohistochemical analysis of paraffin-embedded mouse spleen tissue labelling MEF2C with ab211493 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Nuclear staining in B lymphocytes of mouse spleen but not in T cells in the periarterial lymphatic sheath is observed (PMID: 8506376, PMID 15703219). Counterstained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP)ready to use.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)

Immunohistochemical analysis of paraffin-embedded rat spleen tissue labelling MEF2C with ab211493 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Nuclear staining in B lymphocytes of rat spleen but not in T cells in the periarterial lymphatic sheath is observed (PMID: 8506376, PMID 15703219). Counterstained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP)ready to use.
Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.



Western blot - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)

False colour image of Western blot: Anti-MEF2C antibody [EPR19089-202] - ChIP Grade staining at 1/1000 dilution, shown in green; Mouse anti-GAPDH antibody [6C5] (ab8245) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab211493 was shown to bind specifically to MEF2C. A band was observed at 55/60 kDa in wild-type THP-1 cell lysates with no signal observed at this size in MEF2C knockout cell line (ab313739). To generate this image, wild-type and MEF2C knockout THP-1 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room

temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.
All lanes:
Western blot - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493) at 1/1000 dilution

Lane 1:
Wild-type THP-1 cell lysate at 20 µg

Lane 2:
MEF2C knockout THP-1 cell lysate at 20 µg

Lane 2:
Western blot - Human MEF2C knockout THP-1 cell line (ab313739)

Lane 3:
Daudi cell lysate at 20 µg

Lane 4:
Jurkat cell lysate at 20 µg


Secondary

All lanes:
Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 51 kDa

Observed band size: 55/60 kDa, 37 kDa



Western blot - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time: 92 seconds.

The expression profile observed is consistent with the literature (PMID: 18450586). **Negative control**: Jurkat (PMID: 27876533)

All lanes:
Western blot - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493) at 1/1000 dilution

Lane 1:
Raji (human Burkitt's lymphoma B lymphocyte), whole cell lysate 20 µg

Lane 2:
Jurkat (human T cell leukemia T lymphocyte), whole cell lysate 20 µg

Lane 3:
Ramos (human Burkitt's lymphoma B lymphocyte), whole cell lysate 20µg


Lane 4:
Daudi (human Burkitt's lymphoma lymphoblast), whole cell lysate 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 51 kDa

Observed band size: 50-60 kDa



Western blot - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)

MEF2C Western blot staining using rabbit Anti-MEF2C antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time:

Lanes 1-2,4: 5 minutes

Lane 3: 15 seconds

Lane 5: 26 seconds

All lanes:
Western blot - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493) at 1/1000 dilution

Lane 1:
RAW264.7 (mouse Abelson murine leukemia virus-induced tumor macrophage), whole cell lysate at 10 µg

Lane 2:
NIH/3T3 (mouse embryonic fibroblast), whole cell lysate at 10 µg

Lane 3:
C6 (rat glial tumor glial cell), whole cell lysate at 10 µg

Lane 4:
Mouse brain lysate at 10 µg

Lane 5:
Rat brain lysate at 10 µg


Secondary

Lanes 1, 2, 3 and 5:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Lane 4:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 51 kDa

Observed band size: 50-60 kDa



Western blot - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time: 3 seconds.
All lanes:
Western blot - Anti-MEF2C antibody [EPR19089-202] - ChIP Grade (ab211493) at 1/1000 dilution

Lane 1:
Human MEF2C recombinant protein (aa271-493) 10 ng

Lane 2:
Human MEF2A recombinant protein (aa1-499, ab204772) 10 ng

Lane 2:
Western blot - Human MEF2C knockout THP-1 cell line (ab288702)

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 51 kDa

Observed band size: 26 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.