abcam

ab213362

Anti-CD3D antibody [EPR20544]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR20544
Purification technique	Affinity purification Protein A
Concentration	0.634 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/600
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

IHC-Fr

Tested

Species	Mouse
Dilution info	1/100
Notes	Antigen retrieval: Heated citrate solution (10mM citrate pH 6.0 + 0.05% Tween-20).

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	-

Target data

See full target information Cd3d

Function Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD247/CD3Z. All CD3 chains

contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways. In addition of this role of signal transduction in T-cell activation, CD3D plays an essential role in thymocyte differentiation. Indeed, participates in correct intracellular TCR-CD3 complex assembly and surface expression. In absence of a functional TCR-CD3 complex, thymocytes are unable to differentiate properly (PubMed:10935641). Interacts with CD4 and CD8 and thus serves to establish a functional link between the TCR and coreceptors CD4 and CD8, which is needed for activation and positive selection of CD4 or CD8 T-cells.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

6 product images




Immunohistochemistry (Frozen sections) - Anti-CD3D antibody [EPR20544] (ab213362)

CD3D Immunohistochemistry (Frozen sections) staining using rabbit Anti-CD3D antibody

Immunohistochemical analysis of 4% paraformaldehyde-fixed 0.2% Triton X-100 permeabilized frozen mouse spleen tissue labeling CD3D with ab213362 at 1/100 dilution followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution (green). Cytoplasmic and membranous staining on mouse splenic periarterial lymphatic sheath is observed.

The nuclear counterstain is DAPI (blue).

Secondary antibody only control: Used PBS instead of primary antibody secondary antibody is Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) at 1/1000 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD3D antibody [EPR20544] (ab213362)


CD3D Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD3D antibody

Immunohistochemical analysis of paraffin-embedded mouse thymus tissue labeling CD3D with ab213362 at 1/2000 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Cytoplasmic and membranous staining on T cells of mouse thymus medulla is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry - Anti-CD3D antibody [EPR20544] (ab213362)

CD3D Flow Cytometry staining using rabbit Anti-CD3D antibody

Flow cytometric analysis of mouse spleen cells labeling CD3D with ab213362 at 1/600 dilution (right panel) compared with a rabbit monoclonal IgG isotype control (ab172730) (left panel). Goat anti rabbit IgG (Alexa Fluor® 488) at 1/2000 dilution was used as the secondary antibody.
Firstly cells surface was stained with anti-mouse CD3 Alexa Fluor 647 (Y axis) then cells were fixed with 4% PFA followed by intracellular staining with ab213362. The same population of cells were stained by both: anti-mouse CD3 and anti-mouse CD3D (ab213362) antibodies.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD3D antibody [EPR20544] (ab213362)


CD3D Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD3D antibody

Immunohistochemical analysis of paraffin-embedded mouse spleen tissue labeling CD3D with ab213362 at 1/2000 dilution followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use.
Cytoplasmic staining on T cells of mouse splenic periarterial lymphatic sheath is observed.

Counter stained with Hematoxylin.

Secondary antibody only control: Used PBS instead of primary antibody secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunoprecipitation - Anti-CD3D antibody [EPR20544] (ab213362)

CD3D was immunoprecipitated from 0.35 mg of mouse thymus lysate with ab213362 at 1/30 dilution.
Western blot was performed from the immunoprecipitate using ab213362 at 1/1000 dilution.

VeriBlot for IP Detection Reagent (HRP) (ab131366), was used for detection at 1/10,000 dilution.

Lane 1: Mouse thymus lysate, 10 µg (Input).

Lane 2: ab213362 IP in mouse thymus lysate.

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab213362 in mouse thymus lysate .

Blocking and dilution buffer and concentration: 5% NFDM/TBST.


Exposure time: 10 seconds.

All lanes:

Immunoprecipitation - Anti-CD3D antibody [EPR20544] (ab213362)

Predicted band size: 19 kDa

Observed band size: 24 kDa



Western blot - Anti-CD3D antibody [EPR20544] (ab213362)

CD3D Western blot staining using rabbit Anti-CD3D antibody

Blocking/Dilution buffer: 5% NFDM/TBST.
Exposure time: Lane 1: 15 seconds; Lane 2: 3 minutes.

All lanes:
Western blot - Anti-CD3D antibody [EPR20544] (ab213362) at 1/1000 dilution

Lane 1:
Mouse thymus tissue lysate at 20 µg

Lane 2:
Mouse lymph node tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 19 kDa

Observed band size: 24 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.