abcam

ab223200

Anti-CD16+CD32 antibody [EPR23501-203]

Anti-CD16+CD32 antibody [EPR23501-203] (ab223200) is a rabbit monoclonal antibody detecting CD16+CD32 in **Western Blot, Flow Cytometry, IP, IHC-P, ICC/IF**. Suitable for **Mouse**.

- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23501-203
Purification technique	Affinity purification Protein A
Concentration	0.483 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/500

Notes	-

Not recommended

Species	Recombinant fragment - Mouse, Human, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Not recommended

Species	Recombinant fragment - Mouse, Human, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Not recommended

Species	Recombinant fragment - Mouse, Human, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Recombinant fragment - Mouse
Dilution info	1/1000
Notes	-

Species	Mouse
Dilution info	1/1000
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/1000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Human
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Recombinant fragment - Mouse
Dilution info	-
Notes	-

IHC-Fr

Not recommended

Species	Mouse, Recombinant fragment - Mouse, Human, Rat
Dilution info	-
Notes	-

Target data

See full target information Fcgr3

Function	Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor which binds to IgG1, IgG2a and IgG2b (PubMed:17558411). Mediates neutrophil activation by IgG complexes redundantly with Fcgr4 (PubMed:18097064).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?

Anti-CD16+CD32 antibody [EPR23501-203] (ab223200) is a rabbit recombinant monoclonal antibody and is validated

for use in Western Blot (WB), Flow Cytometry (Flow Cyt), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Mouse samples.

Other related products
We have a range of other formats of antibody clone [EPR23501-203] also available for your convenience: ab223200, Carrier free - ab275095, Alexa Fluor® 647 - ab303478

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD16 and CD32 also referred to as FCγRIII and FCγRII are proteins expressed on the surface of various immune cells like natural killer cells macrophages and some subsets of T cells. CD16 has a mass ranging between 50-80 kDa while CD32 exhibits variability in size between 32-40 kDa due to different isoforms. These proteins play pivotal roles in immune response by binding to the Fc region of immunoglobulins influencing cell activation and signaling.
Biological function summary	CD16 and CD32 serve important roles in mediating immune cell interactions. They work as part of immune synapses facilitating antibody-dependent cellular cytotoxicity (ADCC) and phagocytosis processes. Due to their involvement in these complex activities they are included in immunological synapse formation and modulation extending their influence over overall immune regulation and response. These proteins are known to contribute to both innate and adaptive immune mechanisms as receptors for the Fc portion of immunoglobulin G.
Pathways	CD16 and CD32 interact within signaling pathways like the Fc gamma receptor-mediated phagocytosis pathway and the immune response-regulating signaling pathway. These receptors work in coordination with other immunoglobulin family members to initiate targeted cellular reactions necessary for immune challenge responses. CD32 often partners with CD19 and CD21 within pathways to modulate B cell receptor-mediated signaling.

Associated diseases and disorders	CD16 and CD32 have connections to autoimmune diseases and infectious diseases. Aberrant expression or dysfunction in these proteins can contribute to the development of conditions such as systemic lupus erythematosus where improper immune complex clearance occurs and chronic inflammation results. CD16's involvement with NK cells can also play a role in viral infection control highlighting their importance in pathogen defense mechanisms.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

11 product images




Flow Cytometry - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

CD16+CD32 Flow Cytometry staining using rabbit Anti-CD16+CD32 antibody

Flow cytometric analysis of mouse splenocytes cells labelling CD16+CD32 with ab223200 at 1/500 dilution (Right) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Left). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.


Gated on viable cells.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

Immunohistochemical analysis of paraffin-embedded mouse colon carcinoma tissue labeling CD16+CD32 with ab223200 at 1/1000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on the stroma in mouse colon carcinoma. The section was incubated with ab223200 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Flow Cytometry - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

CD16+CD32 Flow Cytometry staining using rabbit Anti-CD16+CD32 antibody

Flow cytometric analysis of bEnd.3 (mouse brain endothelioma, Left) / J774A.1 (mouse reticulum cell sarcoma monocyte macrophage, Right) cells labelling CD16+CD32 with ab223200 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue).
Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.

Negative control: bEnd.3 (PMID: 23911392).

Gated on viable cells.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

CD16+CD32 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16+CD32 antibody

Immunohistochemical analysis of paraffin-embedded mouse spleen tissue labeling CD16+CD32 with ab223200 at 1/1000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Cytoplasmic and membranouse staining on mouse spleen (PMID: 22618994). The section was incubated with ab223200 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunocytochemistry/ Immunofluorescence - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

CD16+CD32 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-CD16+CD32 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized J774A.1 (mouse reticulum cell sarcoma monocyte macrophage) cells labelling CD16+CD32 with ab223200 at 1/500 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing strong membranous and weak cytoplasmic staining in J774A.1 cells. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.

Negative control: bEnd.3 cell (PMID: 23911392).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

CD16+CD32 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16+CD32 antibody

Immunohistochemical analysis of paraffin-embedded mouse liver tissue labeling CD16+CD32 with ab223200 at 1/1000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on mouse liver. The section was incubated with ab223200 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunoprecipitation - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

CD16+CD32 was immunoprecipitated from 0.35 mg J774A.1 (mouse reticulum cell sarcoma monocyte macrophage) whole cell lysate with ab223200 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab223200 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: J774A.1 (mouse reticulum cell sarcoma monocyte macrophage) whole cell lysate 10 ug

Lane 2: ab223200 IP in J774A.1 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab223200 in J774A.1 whole cell lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 8 seconds.

All lanes:
Immunoprecipitation - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

Predicted band size: 26 kDa

Observed band size: 40-60 kDa



Immunoprecipitation - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

CD16+CD32 was immunoprecipitated from 0.35 mg Mouse spleen tissue lysate with ab223200 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab223200 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: Mouse spleen tissue lysate 10 ug

Lane 2: ab223200 IP in Mouse spleen tissue lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab223200 in mouse spleen tissue lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes.

All lanes:
Immunoprecipitation - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

Predicted band size: 26 kDa

Observed band size: 40-60 kDa



Western blot - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
The molecular weight observed is consistent with what has been described in the literature (PMID: 19503602, 31172847).

Exposure time: 3 minutes.

All lanes:
Western blot - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200) at 1/1000 dilution

Lane 1:
Mouse spleen tissue lysate at 20 µg


Lane 2:
Mouse liver tissue lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/50000 dilution

Predicted band size: 26 kDa

Observed band size: 40-60 kDa



Western blot - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

Blocking/Diluting buffer and concentration: 5% NFDM/TBST The molecular weight observed is consistent with what has been described in the literature (PMID: 19503602, 31172847). Negative control: bEnd.3 (PMID: 23911392). Exposure time: Lanes 1-2: 5.5 seconds
Lanes 1 - 2:
Western blot - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200) at 1/1000 dilution

Lanes 1 - 2:
Western blot - Anti-CD16+CD32 antibody [EPR23501-203] - BSA and Azide free (ab275095)

Lane 1:
J774A.1 (mouse reticulum cell sarcoma monocyte macrophage) whole cell lysate at 20 µg


Lane 2:
bEnd.3 (mouse brain endothelioma) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Observed band size: 40-60 kDa

Exposure time: 5.5s



Western blot - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
This product can recognize both CD16 and CD32.

The loading samples are E.coil extracts containing CD16 or CD32 recombinant protein respectively.

Exposure time: 70 seconds.

All lanes:
Western blot - Anti-CD16+CD32 antibody [EPR23501-203] (ab223200) at 1/1000 dilution

Lane 1:
His-tagged mouse CD16 recombinant protein (aa 31-215), 5 uL

Lane 2:
His-tagged mouse CD32 recombinant protein (aa 30-210), 25 uL

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/50000 dilution

Predicted band size: 26 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.