abcam

ab224803

Anti-beta Catenin antibody [SP328]

Lab Essentials | Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.6 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	SP328
Purification technique	Affinity purification Protein A/G
Purification notes	Purified from TCS by protein A/G.

Reactivity data

IHC-P

Tested

Species	Mouse
Dilution info	1/100
Notes	Primary incubation for 10 minutes at room temperature.

Species	Rat
Dilution info	1/100

Notes	Primary incubation for 10 minutes at room temperature.

Species	Human
Dilution info	1/100
Notes	Primary incubation for 10 minutes at room temperature.

Predicted

Species	Cow, Dog
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/400
Notes	Primary incubation for 30 minutes at 4°C.

Species	Rat
Dilution info	1/400
Notes	Primary incubation for 30 minutes at 4°C.

Species	Human
Dilution info	1/400
Notes	Primary incubation for 30 minutes at 4°C.

Predicted

Species	Cow, Dog
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/400
Notes	Primary incubation for 1 hour at room temperature.

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow, Dog
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Rat
Dilution info	1/10
Notes	-

Species	Human
Dilution info	1/10
Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Cow, Dog
Dilution info	-
Notes	-

IP

Tested

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Predicted

Species	Cow, Dog
Dilution info	-
Notes	-

Target data

See full target information CTNNB1

Function

Key downstream component of the canonical Wnt signaling pathway (PubMed:17524503, PubMed:18077326, PubMed:18086858, PubMed:18957423, PubMed:21262353, PubMed:22155184, PubMed:22647378, PubMed:22699938). In the absence of Wnt, forms a complex with AXIN1, AXIN2, APC, CSNK1A1 and GSK3B that promotes phosphorylation on N-terminal Ser and Thr residues and ubiquitination of CTNNB1 via BTRC and its subsequent degradation by the proteasome (PubMed:17524503, PubMed:18077326, PubMed:18086858, PubMed:18957423, PubMed:21262353, PubMed:22155184, PubMed:22647378, PubMed:22699938). In the presence of Wnt ligand, CTNNB1 is not ubiquitinated and accumulates in the nucleus, where it acts as a coactivator for transcription factors of the TCF/LEF family, leading to activate Wnt responsive genes (PubMed:17524503, PubMed:18077326, PubMed:18086858, PubMed:18957423, PubMed:21262353, PubMed:22155184, PubMed:22647378, PubMed:22699938). Also acts as a coactivator for other transcription factors, such as NR5A2 (PubMed:22187462). Promotes epithelial to mesenchymal transition/mesenchymal to epithelial transition (EMT/MET) via driving transcription of CTNNB1/TCF-target genes (PubMed:29910125). Involved in the regulation of cell adhesion, as component of an E-cadherin:catenin adhesion complex (By similarity). Acts as a negative regulator of centrosome cohesion (PubMed:18086858). Involved in the CDK2/PTPN6/CTNNB1/CEACAM1 pathway of insulin internalization (PubMed:21262353). Blocks anoikis of malignant kidney and intestinal epithelial

cells and promotes their anchorage-independent growth by down-regulating DAPK2 (PubMed:18957423). Disrupts PML function and PML-NB formation by inhibiting RANBP2-mediated sumoylation of PML (PubMed:22155184). Promotes neurogenesis by maintaining sympathetic neuroblasts within the cell cycle (By similarity). Involved in chondrocyte differentiation via interaction with SOX9: SOX9-binding competes with the binding sites of TCF/LEF within CTNNB1, thereby inhibiting the Wnt signaling (By similarity). Acts as a positive regulator of odontoblast differentiation during mesenchymal tooth germ formation, via promoting the transcription of differentiation factors such as LEF1, BMP2 and BMP4 (By similarity). Activity is repressed in a MSX1-mediated manner at the bell stage of mesenchymal tooth germ formation which prevents premature differentiation of odontoblasts (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Beta Catenin also known by names such as CTNNB1 or beta-chip is an important protein involved in cell signaling and adhesion. This protein has a molecular

	weight of around 88 kDa. Beta Catenin is expressed in many cell types and tissues indicating its widespread role in various biological processes. It functions mechanically by mediating the linkage between cadherins and the actin cytoskeleton facilitating cell-cell adhesion. Beta Catenin is also a central part of transcription regulation processes in the nucleus.
Biological function summary	This protein plays roles in both cell adhesion and the regulation of gene expression. Beta Catenin is a critical component of the Wnt signaling pathway where it can form complexes with other proteins to influence gene transcription. In the absence of Wnt signaling beta Catenin levels are low due to its degradation. However when the pathway is active it accumulates in the cytoplasm and eventually translocates to the nucleus where it interacts with TCF/LEF transcription factors to regulate the expression of target genes.
Pathways	Beta Catenin plays a central role in the Wnt signaling pathway and influences cell fate decisions and cellular proliferation. It acts in concert with proteins such as Dishevelled (DVL) and Axin to coordinate these important biological processes. In the absence of Wnt signaling proteins such as APC and GSK-3β are responsible for beta Catenin degradation keeping its cellular levels in check. Beta Catenin's interaction with transcription factors in the nucleus makes it pivotal in the regulation of cell and tissue homeostasis.
Associated diseases and disorders	Beta Catenin has associations with colorectal cancer and hepatocellular carcinoma. Its dysregulation can lead to unchecked cell proliferation and tumorigenesis. Often mutations in the beta Catenin gene (CTNNB1) or components of the Wnt pathway like APC are implicated in the development of these cancers. Its interplay with E-cadherin is important for maintaining tissue architecture and disruptions can lead to invasive cancer phenotypes. Understanding beta Catenin's role provides insights into therapeutic strategies for these cancers.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


34 product images



Flow Cytometry (Intracellular) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Flow Cytometry (Intracellular) staining using rabbit Anti-beta Catenin antibody


Flow cytometric analysis of HEK-293 (human epithelial cell line from embryonic kidney) cell line labeling beta Catenin with ab224803 at 1/400 dilution (green) compared to a negative control of rabbit IgG (blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human hepatocellular carcinoma tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human lung tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human colon tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human stomach adenocarcinoma tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human bladder tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human ovary adenocarcinoma tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human ovary tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human prostate adenocarcinoma tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human lung adenocarcinoma tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human colon adenocarcinoma tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human prostate tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human bladder transitional carcinoma tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human kidney tissue stained for beta Catenin withab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human breast carcinoma tissue sections labeling beta Catenin with ab224803 at 1/100 dilution (1.20 µg/ml). Heat mediated antigen retrieval with sodium Citrate buffer (pH 6.0, epitope retrieval solution 1) for 10mins. Goat Anti-Rabbit & Mouse IgG (HRP) was used as the secondary antibody. Hematoxylin was used as a counterstain. Membranous staining on the human breast carcinoma, performed on a Leica Biosystems BOND™ RX instrument.
The section was incubated with ab224803 for 10 mins at room temperature.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human lung squamous carcinoma tissue stained for beta Catenin withab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human breast ductal carcinoma tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Flow Cytometry (Intracellular) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Flow Cytometry (Intracellular) staining using rabbit Anti-beta Catenin antibody


Flow cytometry analysis of HeLa (human cervix adenocarcinoma epithelial cell) labeling beta Catenin with purified ab224803 at 1/200 dilution (0.495 µg/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as a secondary antibody. Isotype control - Rabbit monoclonal IgG (ab172730) (black). Unlableled control - Unlabelled cells (blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human liver tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human renal cell carcinoma tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-beta Catenin antibody


Immunocytochemistry/ Immunofluorescence analysis of HeLa (human cervix adenocarcinoma epithelial cell) cells labeling beta Catenin with purified ab224803 at 1/10 (9.9 µg/ml). Cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human colon carcinoma tissue sections labeling beta Catenin with ab224803 at 1/100 dilution (1.20 µg/ml). Heat mediated antigen retrieval with sodium Citrate buffer (pH 6.0, epitope retrieval solution 1) for 10mins. Goat Anti-Rabbit & Mouse IgG (HRP) was used as the secondary antibody. Hematoxylin was used as a counterstain. Mainly membranous with nuclear staining on the human colon carcinoma, performed on a Leica Biosystems BOND™ RX instrument.
The section was incubated with ab224803 for 10 mins at room temperature.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Formalin-fixed, paraffin-embedded human stomach tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody

Formalin-fixed, paraffin-embedded human breast tissue stained for beta Catenin with ab224803 at 1/400 dilution in immunohistochemical analysis.



Immunoprecipitation - Anti-beta Catenin antibody [SP328] (ab224803)

beta Catenin was immunoprecipitated from HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate with ab224803 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab224803 at 1/1000 dilution.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

All lanes:
Immunoprecipitation - Anti-beta Catenin antibody [SP328] (ab224803) at 1/1000 dilution

Lane 1:
HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate (Input) at 10 µg

Lane 2:
HeLa (human cervical adenocarcinoma epithelial cell) whole cell lysate (-)


Lane 3:
Rabbit monoclonal IgG (ab172730) instead of ab224803 in HeLa whole cell lysate (-)

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Observed band size: 85 kDa


Exposure time: 3s



Immunocytochemistry/ Immunofluorescence - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-beta Catenin antibody


Immunocytochemistry/ Immunofluorescence analysis of C6 (rat glial tumor glial cell) cells labeling beta Catenin with purified ab224803 at 1/10 (9.9 µg/ml). Cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1/200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Flow Cytometry (Intracellular) staining using rabbit Anti-beta Catenin antibody


Flow cytometry analysis of NIH/3T3 (mouse embryonic fibroblast) labeling beta Catenin with purified ab224803 at 1/200 dilution (0.495 µg/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as a secondary antibody. Isotype control - Rabbit monoclonal IgG (ab172730) (black). Unlableled control -Unlabelled cells (blue).



Flow Cytometry (Intracellular) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Flow Cytometry (Intracellular) staining using rabbit Anti-beta Catenin antibody


Flow cytometry analysis of C6 (rat glial tumor glial cell) labeling beta Catenin with purified ab224803 at 1/200 dilution (0.495 µg/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as a secondary antibody. Isotype control - Rabbit monoclonal IgG (ab172730) (black). Unlableled control - Unlabelled cells (blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Mouse pancreas tissue sections labeling beta Catenin with ab224803 at 1/100 dilution (1.20 µg/ml). Heat mediated antigen retrieval with sodium Citrate buffer (pH 6.0, epitope retrieval solution 1) for 10mins. Goat Anti-Rabbit & Mouse IgG (HRP) was used as the secondary antibody. Hematoxylin was used as a counterstain. Mainly membranous staining on the mouse pancreas, performed on a Leica Biosystems BOND™ RX instrument.
The section was incubated with ab224803 for 10 mins at room temperature.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-beta Catenin antibody [SP328] (ab224803)

Beta Catenin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-beta Catenin antibody

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Rat pancreas tissue sections labeling beta Catenin with ab224803 at 1/100 dilution (1.20 µg/ml). Heat mediated antigen retrieval with sodium Citrate buffer (pH 6.0, epitope retrieval solution 1) for 10mins. Goat Anti-Rabbit & Mouse IgG (HRP) was used as the secondary antibody. Hematoxylin was used as a counterstain. Mainly membranous staining on the rat pancreas, performed on a Leica Biosystems BOND™ RX instrument.
The section was incubated with ab224803 for 10 mins at room temperature.



Immunoprecipitation - Anti-beta Catenin antibody [SP328] (ab224803)

beta Catenin was immunoprecipitated from Rat brain tissue lysate with ab224803 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab224803 at 1/1000 dilution.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

All lanes:
Immunoprecipitation - Anti-beta Catenin antibody [SP328] (ab224803) at 1/1000 dilution

Lane 1:
Rat brain tissue lysate (Input) at 10 µg

Lane 2:
Rat brain tissue lysate (+)


Lane 3:
Rabbit monoclonal IgG (ab172730) instead of ab224834 in rat brain tissue lysate (-)

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Observed band size: 85 kDa

Exposure time: 3s



Immunoprecipitation - Anti-beta Catenin antibody [SP328] (ab224803)

beta Catenin was immunoprecipitated from Mouse brain tissue lysate with ab224803 at 1/30 dilution (2µg in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab224803 at 1/1000 dilution.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

All lanes:
Immunoprecipitation - Anti-beta Catenin antibody [SP328] (ab224803) at 1/1000 dilution

Lane 1:

Mouse brain tissue lysate (Input) at 10 µg

Lane 2:
Mouse brain tissue lysate (+)


Lane 3:
Rabbit monoclonal IgG (ab172730) instead of ab224803 in mouse brain tissue lysate (-)

Secondary

All lanes:
Immunoprecipitation - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/5000 dilution

Observed band size: 85 kDa

Exposure time: 3s



Western blot - Anti-beta Catenin antibody [SP328] (ab224803)

False colour image of Western blot: Anti-beta Catenin antibody [SP328] staining at 1/400 dilution, shown in green; Mouse anti-Alpha Tubulin [DM1A] (ab7291) loading control staining at 1/20000 dilution, shown in red. In Western blot, ab224803 was shown to bind specifically to beta Catenin. A band was observed at 85/90 kDa in wild-type MCF7 cell lysates with no signal observed at this size in CTNNB1 knockout cell line ab286762. To generate this image, wild-type and CTNNB1 knockout MCF7 cell lysates were analysed. First, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3 % milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution.
All lanes:
Western blot - Anti-beta Catenin antibody [SP328] (ab224803) at 1/400 dilution

Lane 1:
Wild-type MCF7 cell lysate at 20 µg

Lane 2:
CTNNB1 knockout MCF7 cell lysate at 20 µg

Lane 3:
HeLa cell lysate at 20 µg


Secondary

All lanes:
Goat anti-Rabbit IgG H&L 800CW and Goat anti-Mouse IgG H&L 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 85 kDa

Observed band size: 85/90 kDa



Western blot - Anti-beta Catenin antibody [SP328] (ab224803)

All lanes:
Western blot - Anti-beta Catenin antibody [SP328] (ab224803) at 1/100 dilution

All lanes:
HEK-293 (human epithelial cell line from embryonic kidney) cell lysate

Predicted band size: 85 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.