abcam

ab227019

Anti-CD19 antibody

Lab Essentials

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7 Preservative: 0.025% Proclin 300 Constituents: PBS, 20% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Polyclonal
Immunogen	Recombinant Fragment Protein within Human CD19. The exact immunogen used to generate this antibody is proprietary information. Database link P15391
Purification technique	Affinity purification Immunogen
Concentration	0.21 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Mouse
Dilution info	1/50 - 1/200
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/500 - 1/3000
Notes	-

Species	Human
Dilution info	1/500 - 1/3000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100 - 1/1000
Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100 - 1/1000

Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information CD19

Function	Functions as a coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes (PubMed:29523808). Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens (PubMed:1373518, PubMed:16672701, PubMed:2463100). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed:12387743, PubMed:16672701, PubMed:9317126, PubMed:9382888). Is not required for early steps during B cell differentiation in the blood marrow (PubMed:9317126). Required for normal differentiation of B-1 cells (By similarity). Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed:1373518, PubMed:2463100). Required for normal levels of serum immunoglobulins, and for production of high-affinity antibodies in response to antigen challenge (PubMed:12387743, PubMed:16672701, PubMed:9317126).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD19 is a transmembrane protein also known as B4 or B-lymphocyte antigen CD19 with an approximate molecular weight of 95 kDa. It is expressed on the surface of B cells throughout their development from pro-B cells to mature B cells. CD19 functions mechanically as a coreceptor enhancing the sensitivity of B cell receptor (BCR) signaling. This protein plays a significant role in promoting the activation and proliferation of B cells by lowering the threshold for antigen receptor engagement.
Biological function summary	CD19 acts as an essential player in B cell signaling. CD19 integrates signals from the BCR with other coreceptors acting as a signaling hub. It participates in forming a signaling complex that includes proteins like CD21 and CD81 which further amplifies BCR-mediated signaling. This complex plays a critical role in B cell activation differentiation and survival ensuring that immune responses are efficiently mounted against pathogens.
Pathways	The CD19 protein fits into key immune response pathways such as the BCR signaling pathway and the PI3K-Akt signaling pathway. CD19 collaborates with other proteins like CD21 CD81 and PI3K within these pathways. Its interaction in the PI3K-Akt pathway for instance supports critical processes including B cell activation and homeostasis contributing to the adaptive immune response.
Associated diseases and disorders	CD19 is connected to B cell malignancies and autoimmune diseases. B cell acute lymphoblastic leukemia (ALL) often shows aberrant CD19 expression making it a valuable target for therapeutic interventions like anti-CD19 monoclonal antibodies. Additionally overexpression or mutations of CD19 may contribute to autoimmune diseases by disrupting normal B cell regulation. CD19's connection with proteins like CD22 and CD20 in these pathological contexts highlights its relevance in developing targeted therapies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


5 product images



Immunocytochemistry/ Immunofluorescence - Anti-CD19 antibody (ab227019)

CD19 Immunocytochemistry/ Immunofluorescence staining of Raji cells using rabbit Anti-CD19 antibody


Immunofluorescence analysis of Raji cells fixed in 4% paraformaldehyde at RT for 15 min labelling CD19 protein at cell membrane using ab227019 at a 1/500 dilution (Green). Hoechst 33342 was used for nuclear staining (Blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD19 antibody (ab227019)

CD19 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human Tonsil using rabbit Anti-CD19 antibody


Immunohistochemical analysis of human tonsil tissue stained for CD19 protein at cell membrane and cytoplasm with ab227019 at a 1/250 dilution.
Antigen retrieval: Citrate buffer, pH 6.0, 15 min.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD19 antibody (ab227019)

CD19 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human Tonsil using rabbit Anti-CD19 antibody


Immunohistochemical analysis of human tonsil tissue stained for CD19 protein at cell membrane with ab227019 at a 1/500 dilution.
Antigen retrieval: Citrate buffer, pH 6.0, 15 min.



Flow Cytometry - Anti-CD19 antibody (ab227019)

CD19 Flow Cytometry staining using rabbit Anti-CD19 antibody

Flow cytometric analysis of 4% paraformaldehyde-fixed mouse splenocytes labeling CD19 with ab227019 at 1/50 dilution (red) compared with an unlabeled sample as a control (black). Acquisition of 20,000 events were collected using a Dylight 488-conjugated secondary antibody for FACS analysis.



Western blot - Anti-CD19 antibody (ab227019)

7.5% SDS-PAGE gel.
All lanes:
Western blot - Anti-CD19 antibody (ab227019) at 1/1000 dilution

All lanes:
Mouse leukocyte extract at 50 µg

Predicted band size: 61 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.