abcam

ab227642

Anti-EGFR antibody [SP84] - C-terminal

Anti-EGFR antibody [SP84] - C-terminal (ab227642) is a rabbit monoclonal antibody detecting EGFR in **Western Blot, Flow Cytometry (Intra), IHC-P, ICC/IF**. Suitable for **Human**.

- Biophysical QC for unrivalled batch-batch consistency

Lab Essentials | Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.6 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	SP84
Purification technique	Affinity purification Protein A/G
Concentration	0.006 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from TCS by protein A/G.

Reactivity data

IHC-P

Tested

Species	Human

Dilution info	1/100
Notes	Primary antibody incubation for 30 minutes at room temperature. Perform heat-mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

Predicted

Species	Mouse, Rat, Cow, Pig
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	1/25
Notes	Primary antibody incubation for 1 hour at room temperature.

Predicted

Species	Mouse, Rat, Cow, Pig
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/30 - 1/1000 for 0.2 µg/mL
Notes	-

Predicted

Species	Mouse, Rat, Cow, Pig
Dilution info	-

Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/200
Notes	Permeabilize with 90% Methanol (-20 °C for 30min) after fixation.

Predicted

Species	Mouse, Rat, Cow, Pig
Dilution info	-
Notes	-

Target data

See full target information EGFR

Function

Receptor tyrosine kinase binding ligands of the EGF family and activating several signaling cascades to convert extracellular cues into appropriate cellular responses (PubMed:10805725, PubMed:27153536, PubMed:2790960, PubMed:35538033). Known ligands include EGF, TGFA/TGF-alpha, AREG, epigen/EPGN, BTC/betacellulin, epiregulin/EREG and HBEGF/heparin-binding EGF (PubMed:12297049, PubMed:15611079, PubMed:17909029, PubMed:20837704, PubMed:27153536, PubMed:2790960, PubMed:7679104, PubMed:8144591, PubMed:9419975). Ligand binding triggers receptor homo- and/or heterodimerization and autophosphorylation on key cytoplasmic residues. The phosphorylated receptor recruits adapter proteins like GRB2 which in turn activates complex downstream signaling cascades. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules (PubMed:27153536). May also activate the NF-kappa-B signaling cascade (PubMed:11116146). Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling (PubMed:11602604). Also phosphorylates MUC1 and increases its interaction with SRC and CTNNB1/beta-catenin (PubMed:11483589). Positively regulates cell migration via interaction with CCDC88A/GIV which retains EGFR at the cell membrane following ligand stimulation, promoting EGFR signaling which triggers cell migration (PubMed:20462955). Plays a role in enhancing learning and memory performance (By similarity). Plays a role in mammalian pain signaling (long-lasting hypersensitivity) (By similarity). Isoform 2 may act as an antagonist of EGF action. (Microbial infection) Acts as a receptor for hepatitis C virus (HCV) in hepatocytes and facilitates its cell entry. Mediates HCV entry by promoting the formation of the

CD81-CLDN1 receptor complexes that are essential for HCV entry and by enhancing membrane fusion of cells expressing HCV envelope glycoproteins.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-EGFR antibody [SP84] - C-terminal (ab227642) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Flow Cytometry (Flow Cyt), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human samples.

What is the molecular weight of EGFR?
Anti-EGFR [SP84] - C-terminal (ab227642) specifically detects a band for EGFR (UniProt: P07327) at a molecular weight of 134kDa.

Collaboration
Anti-EGFR [SP84] - C-terminal (ab227642) is a clone from the portfolio of Spring Bioscience (Roche) SP clones which have been optimised for immunohistochemistry (IHC).

Other related products
We have a range of other formats of antibody clone [SP84] also available for your convenience: ab227642, Carrier free - ab238803, ab279512, Alexa Fluor® 488 - ab320599, Alexa Fluor® 647 - ab320600, Alexa Fluor® 555 - ab320601, Alexa Fluor® 594 - ab320602, PE - ab320603, APC - ab320604, Alexa Fluor® 750 - ab321433

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	EGFR or Epidermal Growth Factor Receptor is a transmembrane glycoprotein that acts as a receptor for members of the epidermal growth factor family. Known

	alternatively as ErbB1 or HER1 this receptor has an approximate molecular weight of 170 kDa. EGFR is expressed in various cell types notably on epithelial cells and can influence multiple cellular processes through its kinase activity. It participates in the regulation of cell growth multiplication and survival by activating its kinase domain upon ligand binding.
Biological function summary	The EGFR protein plays an important role in cellular communication and signaling processes. EGFR pairs with other receptor family members to form active dimers or even higher-order complexes which in turn initiate intracellular signaling cascades. Through these complexes EGFR influences many processes including cell differentiation and repair. This function of EGFR makes it an integral part of mammalian biology affecting how cells respond to their environment by mediating changes in gene expression.
Pathways	EGFR is a central player in the MAPK and PI3K/Akt signaling pathways. Alongside other protein partners like KRAS and PI3 kinase it contributes to transmitting signals from the cell surface to the nucleus affecting gene transcription and cell behavior. These pathways are important for normal cell growth and division and aberrations in these pathways can lead to excessive or insufficient cell proliferation.
Associated diseases and disorders	EGFR is pertinent to cancer biology including non-small cell lung cancer and glioblastoma where mutations or overexpression of the receptor frequently occur. It connects to proteins such as PTEN and BRAF which influence tumor progression and response to targeted therapies. EGFR's involvement in these disorders highlights its significance as a therapeutic target since it can be manipulated to alter disease progression.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.


Full details and terms and conditions can be found here:
Terms & Conditions.

16 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)


Formalin-fixed, paraffin-embedded human lung squamous cell carcinoma tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-EGFR antibody


Formalin-fixed, paraffin-embedded human colon adenocarcinoma tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-EGFR antibody


ab227642 staining EGFR in HCT116 (positive) and HCT116-EGFR KO (negative) cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated for 2 hours at room temperature with ab227642 at 0.2 µg/ml (shown in green) and ab7291 Anti-alpha Tubulin antibody [DM1A] - Loading Control (shown in pseudocolour magenta). Cells were then incubated with ab150081, Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution and ab150120, Goat Anti-Mouse IgG H&L (Alexa Fluor® 594) preadsorbed at 1/1000 dilution for 1 hour at room temperature. Nuclear DNA was labeled with DAPI (shown in blue. Image was acquired with a high-content analyser (Operetta CLS, Revvity) and a maximum intensity projection of confocal sections is shown.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-EGFR antibody


Formalin-fixed, paraffin-embedded human piver tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-EGFR antibody


Formalin-fixed, paraffin-embedded human breast ductal carcinoma tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-EGFR antibody


Formalin-fixed, paraffin-embedded human placenta tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-EGFR antibody


Formalin-fixed, paraffin-embedded human renal cell carcinoma tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-EGFR antibody


Formalin-fixed, paraffin-embedded human liver hepatocellular carcinoma tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-EGFR antibody


Formalin-fixed, paraffin-embedded human stomach adenocarcinoma tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-EGFR antibody


Formalin-fixed, paraffin-embedded human bladder transitional cell carcinoma tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-EGFR antibody


Formalin-fixed, paraffin-embedded human lung squamous cell carcinoma tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-EGFR antibody


Immunocytochemistry/ Immunofluorescence analysis of A431(human epidermoid carcinoma epithelial cell) cells labeling EGFR with purified ab227642 at 1:50 (1.92 µg/ml). Cells were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. Cells were counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1:1000 (2 µg/ml) dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.



Flow Cytometry (Intracellular) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Flow Cytometry (Intracellular) staining using rabbit Anti-EGFR antibody


Flow Cytometry analysis of A431(Human epidermoid carcinoma epithelial cell) cells labeling EGFR with purified ab227642 at 1:200 dilution (0.48 µg/ml) Red. Cells were fixed with 4% paraformaldehyde . A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1:2000 dilution. Isotype control - Rabbit monoclonal IgG (ab172730) / Black. Unlabeled control - Unlabelled cells / Blue.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-EGFR antibody

Formalin-fixed, paraffin-embedded human lung adenocarcinoma tissue stained for EGFR using ab227642 at 1/100 dilution in immunohistochemical analysis.



Copyright © 2026 Abcam Limited

Western blot - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

EGFR Western blot staining using rabbit Anti-EGFR antibody

Western blot: Anti-EGFR antibody [SP84] - C-terminal ab227642 staining at 1/200 dilution, shown in green; Mouse anti alpha Tubulin ab7291 loading control staining at 1/20000 dilution, shown in magenta. A band was observed at 160 kDa in Wild-type A549 cell lysates with no signal observed at this size in EGFR knockout A549 cell line. To generate this image, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 3% Milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit 800CW and Goat anti-Mouse 680RD at 1/20,000 dilution.
All lanes:
Western blot - Anti-EGFR antibody [SP84] - C-terminal (ab227642) at 1/200 dilution

Lane 1:
Wild-type A549 cell lysates at 20 µg

Lane 2:
EGFR knockout A549 cell lysates at 20 µg

Lane 3:
Wild-type HeLa ab255928 cell lysates at 20 µg

Lane 4:
Western blot - Human EGFR knockout HeLa cell line (ab255385) at 20 µg


Secondary

All lanes:
Goat anti-Rabbit 800CW & Goat anti-Mouse 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 134 kDa

Observed band size: 160 kDa



Western blot - Anti-EGFR antibody [SP84] - C-terminal (ab227642)

All lanes:
Western blot - Anti-EGFR antibody [SP84] - C-terminal (ab227642) at 1/25 dilution

All lanes:
A431 (human epidermoid carcinoma cell line) EGF treated cell lysate

Predicted band size: 134 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.