abcam

ab227665

Anti-CD16a antibody [SP189] - C-terminal

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.6 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	SP189
Purification technique	Affinity purification Protein A/G
Concentration	0.073 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from TCS by protein A/G.

Reactivity data

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/100
Notes	-

WB

Tested

Species	Human
Dilution info	1/100
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	Antigen retrieval: Boil tissue section in EDTA buffer, pH 8.0 for 10 minutes followed by cooling at room temperature for 20 minutes.

Target data

See full target information FCGR3A

Function

Receptor for the invariable Fc fragment of immunoglobulin gamma (IgG). Optimally activated upon binding of clustered antigen-IgG complexes displayed on cell surfaces, triggers lysis of antibody-coated cells, a process known as antibody-dependent cellular cytotoxicity (ADCC). Does not bind free monomeric IgG, thus avoiding inappropriate effector cell activation in the absence of antigenic trigger (PubMed:11711607, PubMed:21768335, PubMed:22023369, PubMed:24412922, PubMed:25786175, PubMed:25816339, PubMed:28652325, PubMed:8609432, PubMed:9242542). Mediates IgG effector functions on natural killer (NK) cells. Binds antigen-IgG complexes generated upon infection and triggers NK cell-dependent cytokine production and degranulation to limit viral load and propagation. Involved in the generation of memory-like adaptive NK cells capable to produce high amounts of IFNG and to efficiently eliminate virus-infected cells via ADCC (PubMed:24412922, PubMed:25786175). Regulates NK cell survival and proliferation, in particular by preventing NK cell progenitor apoptosis (PubMed:29967280, PubMed:9916693). Following the engagement of antigen-IgG complexes, triggers phosphorylation of immunoreceptor tyrosine-based activation motif (ITAM)-containing adapters with subsequent activation of phosphatidylinositol 3-kinase signaling and sustained elevation of intracellular calcium that ultimately drive NK cell activation. The ITAM-dependent signaling coupled to receptor phosphorylation by PKC mediates robust intracellular calcium flux that leads to production of pro-inflammatory cytokines, whereas in the absence of receptor phosphorylation it mainly activates phosphatidylinositol 3-kinase signaling leading to cell degranulation (PubMed:1825220, PubMed:23024279, PubMed:2532305). Costimulates NK cells and trigger lysis of

target cells independently of IgG binding (PubMed:10318937, PubMed:23006327). Mediates the antitumor activities of therapeutic antibodies. Upon ligation on monocytes triggers TNFA-dependent ADCC of IgG-coated tumor cells (PubMed:27670158). Mediates enhanced opsonisation and ADCC in response to afucosylated IgGs (PubMed:34485821, PubMed:28566370). (Microbial infection) Involved in Dengue virus pathogenesis via antibody-dependent enhancement (ADE) mechanism. Secondary infection with Dengue virus triggers elevated levels of afucosylated non-neutralizing IgG1s with reactivity to viral envelope/E protein. Viral antigen-IgG1 complexes bind with high affinity to FCGR3A, facilitating virus entry in myeloid cells and subsequent viral replication.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The CD16a protein also known as Fc gamma receptor IIIa (FcγRIIIa) plays an important role in the immune system. This receptor is about 50-70 kDa in mass and is expressed on the surface of natural killer (NK) cells macrophages and some T cells. CD16a facilitates antibody-dependent cellular cytotoxicity (ADCC) by binding to the Fc region of immunoglobulin G (IgG). This interaction allows effector immune cells to recognize and destroy antibody-coated target cells.
Biological function summary	CD16a is important for immune responses and occurs as part of a receptor complex. It signals downstream effects through its association with the CD3 zeta chain and the Fc epsilon receptor gamma chain which possess immunoreceptor tyrosine-based activation motifs (ITAMs). Upon binding to IgG CD16a triggers a cascade of events leading to the release of cytokines and cytotoxic granules. This action helps in eliminating infected or cancerous cells demonstrating its role in both innate and adaptive immunity.
Pathways	CD16a contributes significantly to the immunological synapse formation and NK cell activation pathway. This receptor cooperates with other proteins such as CD3

	and CD19 to modulate the immune response. It also participates in signaling pathways like the PI3K-Akt pathway which regulates cell survival and proliferation. CD16a's interaction with these pathways highlights its importance in immune cell communication and function.
Associated diseases and disorders	CD16a is linked to autoimmune diseases and certain cancers. In autoimmune disorders like systemic lupus erythematosus (SLE) abnormal CD16a function can contribute to the pathogenesis by altering immune cell targeting. Additionally in cancers such as lymphoma CD16a's interaction with IgG-coated cancer cells can influence the efficacy of monoclonal antibody therapies. The receptor's association with proteins like CD3 and CD19 in these disease contexts highlights its potential as a therapeutic target.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.


Full details and terms and conditions can be found here:
Terms & Conditions.

15 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)


Formalin-fixed, paraffin-embedded human B-cell lymphoma tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody


Formalin-fixed, paraffin-embedded human colon adenocarcinoma tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody


Formalin-fixed, paraffin-embedded human cervical squamous cell carcinoma tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human spleen tissue sections labeling CD16a with ab227665 at 1/100 dilution (1.47 µg/ml). Heat mediated antigen retrieval was performed Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 10 mins. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody


Formalin-fixed, paraffin-embedded human stomach adenocarcinoma tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody


Formalin-fixed, paraffin-embedded human liver tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Flow Cytometry (Intracellular) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Flow Cytometry (Intracellular) staining using rabbit Anti-CD16a antibody


Flow cytometric analysis of Jurkat (human T cell leukemia cell line from peripheral blood) cell line labeling CD16a with ab227665 at 1/100 (green) compared with a rabbit isotype control (blue).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody


Formalin-fixed, paraffin-embedded human lung tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody


Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human liver tissue sections labeling CD16a with ab227665 at 1/100 dilution (1.47 µg/ml). Heat mediated antigen retrieval was performed Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution 2) for 10 mins. Rabbit specific IHC polymer detection kit HRP/DAB (ab209101) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody


Formalin-fixed, paraffin-embedded human HK lymphoma tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody


Formalin-fixed paraffin-embedded human liver hepatocellular carcinoma tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody


Formalin-fixed, paraffin-embedded human placenta tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)


CD16a Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16a antibody

Formalin-fixed, paraffin-embedded human spleen tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

Formalin-fixed, paraffin-embedded human thymus tissue stained for CD16a with ab227665 at 1/100 dilution in immunohistochemical analysis.



Western blot - Anti-CD16a antibody [SP189] - C-terminal (ab227665)

All lanes:
Western blot - Anti-CD16a antibody [SP189] - C-terminal (ab227665) at 1/100 dilution

All lanes:
K562 (human chronic myelogenous leukemia cell line from bone marrow) cell lysate

Developed using the ECL technique.

Predicted band size: 29 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.