abcam

ab228455

Anti-CD24 antibody [EPR19925] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR19925
Purification technique	Affinity purification Protein A
Concentration	1.046 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	-

Notes	-

ICC/IF

Tested

Species	Human
Dilution info	-
Notes	-

Target data

See full target information CD24

Function	May have a pivotal role in cell differentiation of different cell types. Signaling could be triggered by the binding of a lectin-like ligand to the CD24 carbohydrates, and transduced by the release of second messengers derived from the GPI-anchor. Modulates B-cell activation responses. Promotes AG-dependent proliferation of B-cells, and prevents their terminal differentiation into antibody-forming cells (PubMed:11313396). In association with SIGLEC10 may be involved in the selective suppression of the immune response to danger-associated molecular patterns (DAMPs) such as HMGB1, HSP70 and HSP90. Plays a role in the control of autoimmunity (By similarity).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab228455 is the carrier-free version of ab202073.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD24 also known as heat-stable antigen or CD24A is a glycosylphosphatidylinositol (GPI)-anchored membrane protein with a molecular mass of approximately 30-50 kDa. It is expressed on the surface of many cell types including B cells T cells neutrophils and epithelial cells. CD24 plays a role in cell adhesion and signaling processes impacting immune response and cancer progression. Its expression is often used as a marker in identification and classification procedures such as flow cytometry.
Biological function summary	CD24 functions like a molecular "switch" modulating signals sent between cells. It acts as a ligand for multiple receptors and participates in cellular communication especially influencing the immune system. Part of its biological role involves interaction within cellular complexes impacting immune modulation and inflammatory responses. CD24's location at the interface of cellular microenvironments makes it critical for cellular response adaptation to varied external stimuli.
Pathways	CD24 participates in both immune signaling and cancer-related pathways. It plays a significant role in the suppression of the immune response often interacting with proteins like Siglec-10 in these pathways. CD24's involvement in these pathways contributes to cell survival and proliferation particularly within cancer development. The protein’s interaction within specified signaling pathways allows it to influence downstream targets that regulate gene expression and cellular functions.
Associated diseases and disorders	CD24 has significant connections to cancer and inflammatory diseases. It is highly expressed in various cancers including breast and prostate cancer where it

correlates with poor prognosis due to its role in apoptosis evasion and metastasis. CD24 interacts with proteins like HMGB1 and is implicated in inflammatory conditions where it modulates immune cell activity. Extensive research into CD24's involvement in these disorders may lead to targeted therapies including the development of anti-CD24 antibodies for therapeutic interventions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

4 product images




Flow Cytometry - Anti-CD24 antibody [EPR19925] - BSA and Azide free (ab228455)

CD24 Flow Cytometry staining using rabbit Anti-CD24 antibody

Flow cytometric analysis of MDA-MB-231 (human breast adenocarcinoma cell line, Left) and MCF7 breast adenocarcinoma cell line, Right) cell lines labeling CD24 with ab202073 at 1/500 (red) compared with a Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) (black) and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) at 1/2000 dilution was used as the secondary antibody.
Gated on total viable cells. Negative control: MDA-MB-231 (PMID: 18433506 and 22934288).

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab202073).



Immunocytochemistry/ Immunofluorescence - Anti-CD24 antibody [EPR19925] - BSA and Azide free (ab228455)


CD24 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-CD24 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized MCF7 (human breast adenocarcinoma cell line) and MDA-MB-231 (human breast adenocarcinoma cell line) cells labeling CD24 with ab202073 at 1/100 dilution, followed by Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution (green). Confocal image showing membranous staining on MCF7 cell line.
Negative control: MDA-MB-231 (PMID: 18433506).

The nuclear counter stain is DAPI (blue). Tubulin is detected with Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) (ab195889) (red) at 1/200 dilution.

Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab202073).




Flow Cytometry - Anti-CD24 antibody [EPR19925] - BSA and Azide free (ab228455)

CD24 Flow Cytometry staining of Human peripheral blood mononuclear cells using rabbit Anti-CD24 antibody

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab202073).
Flow cytometry staining of human peripheral blood mononuclear cells (PBMCs) with ab202073 (right) or Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (left). PBMCs were incubated for 30 min on ice in 1x PBS containing 10 µg/ml human IgG and 10 % normal goat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody ab202073 or Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (1x 10^6 in 100 µl at 5.0 µg/ml (1/400)) for 30min on ice. The cells were simultaneously stained with CD3.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min on ice

Acquisition of >30000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter. Events were gated on live cells.



Flow Cytometry - Anti-CD24 antibody [EPR19925] - BSA and Azide free (ab228455)

CD24 Flow Cytometry staining of Human peripheral blood mononuclear cells using rabbit Anti-CD24 antibody

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab202073).

Flow cytometry staining of human peripheral blood mononuclear cells (PBMCs) with ab202073 (right) or Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (left). PBMCs were incubated for 30 min on ice in 1x PBS containing 10 µg/ml human IgG and 10 % normal goat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody ab202073 or Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (1x 10^6 in 100 µl at 5.0 µg/ml (1/400)) for 30min on ice. The cells were simultaneously stained with CD19.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min on ice

Acquisition of >30000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter. Events were gated on live cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.