abcam

ab234111

# Anti-CPT1A antibody [EPR21843-71-2F]

Anti-CPT1A antibody [EPR21843-71-2F] (ab234111) is a rabbit monoclonal antibody detecting CPT1A in **Western Blot, Flow Cytometry (Intra), IHC-P**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 30 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

## Key facts

| | |
|---|---|
| **Isotype** | IgG |
| **Host species** | Rabbit |
| **Storage buffer** | pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA |
| **Form** | Liquid |
| **Clonality** | Monoclonal |
| **Immunogen** | The exact immunogen used to generate this antibody is proprietary information. |
| **Clone number** | EPR21843-71-2F |
| **Purification technique** | Affinity purification Protein A |
| **Specificity** | We detected weak cross-reactivity with CPT1B with the recombinant protein only. Our WB images were generated by testing unboiled samples only. |
| **Concentration** | 0.48 mg/mL The concentration of this product may be batch-dependent<br>Batch concentration finder → |

## Reactivity data

### WB

### Tested

| | |
|---|---|
| **Species** | Mouse |
| **Dilution info** | 1/1000 |
| **Notes** | - |

| | |
|---|---|
| **Species** | Rat |
| **Dilution info** | 1/1000 |
| **Notes** | - |

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/1000 |
| **Notes** | - |

## Flow Cyt (Intra)

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/50 |
| **Notes** | - |

### Expected

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## IHC-P

### Tested

| | |
|---|---|
| **Species** | Mouse |
| **Dilution info** | 1/1000 |
| **Notes** | - |

| | |
|---|---|
| **Species** | Rat |

| | |
|---|---|
| **Dilution info** | 1/1000 |
| **Notes** | - |

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 1/1000 |
| **Notes** | - |

## Target data

See full target information CPT1A

| | |
|---|---|
| **Function** | Catalyzes the transfer of the acyl group of long-chain fatty acid-CoA conjugates onto carnitine, an essential step for the mitochondrial uptake of long-chain fatty acids and their subsequent beta-oxidation in the mitochondrion (PubMed:11350182, PubMed:14517221, PubMed:16651524, PubMed:9691089). Also possesses a lysine succinyltransferase activity that can regulate enzymatic activity of substrate proteins such as ENO1 and metabolism independent of its classical carnitine O-palmitoyltransferase activity (PubMed:29425493). Plays an important role in hepatic triglyceride metabolism (By similarity). Also plays a role in inducible regulatory T-cell (iTreg) differentiation once activated by butyryl-CoA that antagonizes malonyl-CoA-mediated CPT1A repression (By similarity). Sustains the IFN-I response by recruiting ZDHCC4 to palmitoylate MAVS at the mitochondria leading to MAVS stabilization and activation (PubMed:38016475). Promotes ROS-induced oxidative stress in liver injury via modulation of NFE2L2 and NLRP3-mediated signaling pathways (By similarity). |

## Storage

| | |
|---|---|
| **Shipped at conditions** | Conditional Ambient |
| **Appropriate short-term storage duration** | 1-2 weeks |
| **Appropriate short-term storage conditions** | +4°C |
| **Appropriate long-term storage conditions** | -20°C |
| **Aliquoting information** | Upon delivery aliquot |
| **Storage information** | Avoid freeze / thaw cycle |

## Notes

**What is this antibody validated in?**
Anti-CPT1A antibody [EPR21843-71-2F] (ab234111) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunohistochemistry (IHC-P) in Human, Mouse, Rat samples.

**What is the molecular weight of CPT1A?**
Anti-CPT1A [EPR21843-71-2F] (ab234111) specifically detects a band for CPT1A (UniProt: P50416) at a molecular weight of 88kDa.

**Trusted by the scientific community**
Anti-CPT1A [EPR21843-71-2F] (ab234111) was first used in a scientific publication in 2018 and has been cited over 30 times in peer-reviewed journals.

**Specificity confirmed**
The specificity of Anti-CPT1A antibody [EPR21843-71-2F] (ab234111) has been confirmed by Western blot testing in CPT1A Knockout HAP1 cells.

**Other related products**
We have a range of other formats of antibody clone [EPR21843-71-2F] also available for your convenience: ab234111, Carrier free - ab235841

**Patented technology**
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

**What are the advantages of a recombinant monoclonal antibody?**
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

**Trial sizes available!**
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

## Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

| | |
|---|---|
| **Activity summary** | CPT1A also known as carnitine palmitoyltransferase 1A is a vital enzyme involved in the transport of long-chain fatty acids across the mitochondrial membrane. This protein facilitates the conversion of fatty acids into acyl-carnitine a process necessary for beta-oxidation within the mitochondria. The molecular weight of CPT1A is approximately 88 kDa. CPT1A predominantly expresses in liver cells where energy metabolism from fatty acids is important. Alternate names for this target include CPT 1 and CTP1A reflecting its central role in metabolic processes related to lipid utilization. |

| | |
|---|---|
| **Biological function summary** | CPT1A plays a vital role in energy production by facilitating mitochondrial fatty acid oxidation. It does not function as a solitary enzyme but often associates with other enzymes forming a multiprotein complex that includes acyl-CoA synthetase. This complex enables efficient fatty acid transport and oxidation converting stored fats into usable energy. As CPT1A primarily operates in the liver its activity significantly impacts overall lipid and energy homeostasis demonstrating its critical regulatory role. |
| **Pathways** | CPT1A is central to the fatty acid beta-oxidation pathway an important process for breaking down fatty acids to produce energy. This pathway also involves proteins such as CPT1B a related isoform present in muscle tissues. CPT1A's function is important in the liver's capacity to regulate energy balance and respond to metabolic demands. Additionally CPT1A interacts within the signaling pathway of AMPK (AMP-activated protein kinase) which further integrates it into broader metabolic regulation networks linking energy status to cellular function. |
| **Associated diseases and disorders** | CPT1A is notably associated with metabolic syndromes and fatty liver disease. Mutations or deficiencies in CPT1A can lead to disorders like hepatic carnitine palmitoyltransferase 1A deficiency characterized by hypoketotic hypoglycemia and hepatomegaly. Furthermore its dysregulation can impact proteins such as ACC1 (acetyl-CoA carboxylase) in the context of metabolic syndrome. Research continues to explore the implications of CPT1A in non-alcoholic steatohepatitis highlighting its relevance in liver energy metabolism disorders and their pathophysiology. |

## Product promise

**Tested**

We have tested this species and application combination and it works. It is covered by our product promise.

**Expected**

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

**Predicted**

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

**Not recommended**

We do not recommend this combination. It is not covered by our product promise.


---

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

## 12 product images





### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

CPT1A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CPT1A antibody

Immunohistochemical analysis of paraffin-embedded human kidney tissue labeling CPT1A with ab234111 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Granular cytoplasmic staining in human kidney (PMID: 18192268; PMID: 28956034; PMID: 15363638; PMID: 8679700) is observed. Counter stained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.





### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

CPT1A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CPT1A antibody

Immunohistochemical analysis of paraffin-embedded human heart tissue labeling CPT1A with ab234111 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Cytoplasmic staining on human heart is observed. Counter stained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




### Flow Cytometry (Intracellular) - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)


CPT1A Flow Cytometry (Intracellular) staining of HAP1 using rabbit Anti-CPT1A antibody

Flow cytometry overlay histogram showing left wild-type HAP1 positive cells and right negative CPT1A knockout HAP1 stained with ab234111 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilised with 0.1% PBS-Triton X-100 for 15 min. The cells were then incubated in 1x PBS containing 10% normal goat serum to block non-specific protein-protein interaction followed by the antibody (ab234111) (1x $10^6$ in 100µl at 1 µg/ml (1/481)) for 30min at 22°C.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min at 22°C.

Isotype control antibody was Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (black line) used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.


Acquisition of >5000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter.




## Flow Cytometry (Intracellular) - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

CPT1A Flow Cytometry (Intracellular) staining using rabbit Anti-CPT1A antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol-permeabilized SK-OV-3 (human ovarian cancer cell line) cell line labeling CPT1A with ab234111 at 1/50 dilution (red) compared with a Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) (black) and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) at 1/2000 dilution was used as the secondary antibody.





## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

CPT1A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CPT1A antibody

Immunohistochemical analysis of paraffin-embedded human ovarian carcinoma tissue labeling CPT1A with ab234111 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Granular cytoplasmic staining in human ovarian carcinoma (PMID: 26716645) is observed. Counter stained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.





## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

CPT1A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CPT1A antibody

Immunohistochemical analysis of paraffin-embedded mouse kidney tissue labeling CPT1A with ab234111 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Cytoplasmic staining in mouse kidney (PMID: 18192268; PMID: 28956034; PMID: 15363638; PMID: 8679700) is observed. Counter stained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.





## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

CPT1A Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CPT1A antibody

Immunohistochemical analysis of paraffin-embedded rat kidney tissue labeling CPT1A with ab234111 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Granular cytoplasmic staining in rat kidney (PMID: 18192268; PMID: 28956034; PMID: 15363638; PMID: 8679700) is observed. Counter stained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

---




## Western blot - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

CPT1A Western blot staining using rabbit Anti-CPT1A antibody

**Lanes 1- 2:** Merged signal (red and green). Green - ab234111 observed at 88 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (ab8245) observed at 37 kDa.
ab234111 was shown to react with CPT1A in wild-type HEK-293T cells in western blot. Loss of signal was observed when knockout cell line ab266319 (knockout cell lysate ab256880) was used. Wild-type HEK-293T and CPT1A knockout HEK-293T cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab234111 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111) at 1/1000 dilution

Lane 1:
Wild-type HEK-293T cell lysate at 20 µg


Lane 2:
Western blot - Human CPT1A knockout HEK-293T cell lysate (ab256880) at 20 µg

Performed under reducing conditions.

Predicted band size: 88 kDa

Observed band size: 88 kDa

---




## Western blot - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

Blocking/Dilution buffer: 5% NFDM/TBST.
ab234111 was shown to specifically react with CPT1A in wild-type HAP1 cells as signal was lost in CPT1A knockout cells. Wild-type and CPT1A knockout samples were subjected to SDS-PAGE. ab234111 and ab181602 (Rabbit anti-GAPDH loading control) were incubated 1 hour at room temperature at 1/1000 dilution and 1/200,000 dilution respectively. Blots were developed with Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) secondary antibody at 1/100,000

dilution for 1 hour at room temperature before imaging. The blot was developed on a BIO-RAD® ChemiDoc™ MP instrument using the ECL technique.
All lanes:
Western blot - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111) at 1/1000 dilution

Lane 1:
Wild-type HAP1 whole cell lysate at 20 µg

Lane 2:
CPT1A knockout HAP1 whole cell lysate at 20 µg

Lane 3:
HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 20 µg

Lane 4:
SK-OV-3 (human ovarian cancer cell line) whole cell lysate at 20 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 88 kDa

Observed band size: 88 kDa

Exposure time: 92s




## Western blot - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

Blocking and diluting buffer: 5% NFDM/TBST
Lanes 1 - 3:
Western blot - Anti-CPT1A antibody [EPR21843-71-2F] - BSA and Azide free (ab235841) at 1/1000 dilution

Lanes 1 - 3:
Western blot - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111) at 1/1000 dilution

Lane 1:
MCF7 (Human breast adenocarcinoma epithelial cell) cell lysate at 20 µg

Lane 2:
HeLa (Human cervix adenocarcinoma epithelial cell) cell lysate at 20 µg


Lane 3:
HepG2 (Human hepatocellular carcinoma epithelial cell) cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/20000 dilution

Observed band size: 88 kDa

Exposure time: 180s




## Western blot - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

Blocking and diluting buffer: 5% NFDM /TBST
Lanes 1 - 2:
Western blot - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111) at 1/1000 dilution

Lanes 1 - 2:
Western blot - Anti-CPT1A antibody [EPR21843-71-2F] - BSA and Azide free (ab235841) at 1/1000 dilution

Lane 1:
Human kidney tissue lysate at 20 µg


Lane 2:
Human liver tissue lysate at 1/20 dilution

Secondary

All lanes:
Western blot - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/1000 dilution

Observed band size: 88 kDa

Exposure time: 92s



## Western blot - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111)

CPT1A Western blot staining using rabbit Anti-CPT1A antibody

**Exposure time :** Lanes 1 and 2: 3 minutes; Lane 3: 37 seconds; Lane 4: 114 seconds; Lane 5: 15 seconds; Lanes 6 and 7: 92 seconds.
Blocking/Dilution buffer: 5% NFDM/TBST.

All lanes:
Western blot - Anti-CPT1A antibody [EPR21843-71-2F] (ab234111) at 1/1000 dilution

Lane 1:
C6 (rat glial tumor cell line) whole cell lysate at 20 µg

Lane 2:
MCF7 (human breast adenocarcinoma cell line) whole cell lysate at 20 µg

Lane 3:
Mouse kidney lysate at 20 µg

Lane 4:
Rat kidney lysate at 20 µg

Lane 5:
Human kidney lysate at 20 µg

Lane 6:
HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 20 µg

Lane 7:
SK-OV-3 (human ovarian cancer cell line) whole cell lysate at 20 µg

Secondary

Lanes 1, 2, 3, 4, 6 and 7:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Lane 5:
Western blot - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/1000 dilution

Predicted band size: 88 kDa

Observed band size: 88 kDa

---

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.