abcam

ab236868

Anti-SCD1 antibody [EPR21963]

Anti-SCD1 antibody [EPR21963] (ab236868) is a rabbit monoclonal antibody detecting SCD1 in **Western Blot, Flow Cytometry (Intra), IP, IHC-P, ICC/IF**. Suitable for **Human, Mouse, Rat**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 20 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogens	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR21963
Purification technique	Affinity purification Protein A
Specificity	This SCD1 antibody can bind to SCD5 very weakly in WB when SCD5 is overexpressed (<10% cross-reactivity).
Concentration	0.606 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	1/30
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/500
Notes	-

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Human
Dilution info	1/2000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information SCD

Function	Stearoyl-CoA desaturase that utilizes O(2) and electrons from reduced cytochrome b5 to introduce the first double bond into saturated fatty acyl-CoA substrates (PubMed:15907797, PubMed:18765284). Catalyzes the insertion of a cis double bond at the delta-9 position into fatty acyl-CoA substrates including palmitoyl-CoA and stearoyl-CoA (PubMed:15907797, PubMed:18765284). Gives rise to a mixture of 16:1 and 18:1 unsaturated fatty acids (PubMed:15610069). Plays an important role in lipid biosynthesis. Plays an important role in regulating the expression of genes that are involved in lipogenesis and in regulating mitochondrial fatty acid oxidation (By similarity). Plays an important role in body energy homeostasis (By similarity). Contributes to the biosynthesis of membrane phospholipids, cholesterol esters and triglycerides (By similarity).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-SCD1 antibody [EPR21963] (ab236868) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Immunoprecipitation (IP), Immunohistochemistry (IHC-P), Immunocytochemistry/immunofluorescence (ICC/IF) in Human, Mouse, Rat samples.

What is the molecular weight of SCD1?
Anti-SCD1 [EPR21963] (ab236868) specifically detects a band for SCD1 (UniProt: O00767) at a molecular weight of 42kDa.

Trusted by the scientific community
Anti-SCD1 [EPR21963] (ab236868) was first used in a scientific publication in 2018 and has been cited over 20 times in peer-reviewed journals.

Specificity confirmed

The specificity of Anti-SCD1 antibody [EPR21963] (ab236868) has been confirmed by Western blot testing in SCD Knockout HeLa cell line, ab265220.

Other related products
We have a range of other formats of antibody clone [EPR21963] also available for your convenience: ab236868, Carrier free - ab238171, Alexa Fluor® 647 - ab314925, Alexa Fluor® 488 - ab315182

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Stearoyl-CoA desaturase-1 (SCD1) also known as delta-9-desaturase is an enzyme integral to the synthesis of monounsaturated fats. This enzyme converts saturated fatty acyl-CoAs to monounsaturated fatty acyl-CoAs by introducing a double bond between the ninth and tenth carbon atoms a process called desaturation. SCD1 has a molecular weight of approximately 45 kDa and expresses in a variety of tissues including the liver adipose tissue and heart.
Biological function summary	SCD1 plays a significant role in fatty acid metabolism by regulating the balance of saturated and unsaturated fatty acids which impacts membrane fluidity and cellular signaling. SCD1 does not function as part of a multi-protein complex acting independently in the endoplasmic reticulum. Its expression and activity influence lipid biosynthesis and storage affecting energy homeostasis and lipid composition in cells.
Pathways	Fatty acid synthesis and oxidation pathways extensively involve SCD1. It participates mainly in the lipogenesis pathway interacting with proteins like acetyl-CoA carboxylase (ACC) and fatty acid synthase (FAS). SCD1 contributes to maintaining the ratio of oleate and palmitoleate key monounsaturated fatty acids important for cellular functions by integrating into the lipid metabolism network.
Associated diseases and disorders	SCD1 is implicated in conditions like obesity and metabolic syndrome. Altered SCD1 activity links to insulin resistance as seen in obesity by modulating lipid profiles that influence insulin sensitivity. SCD1 associates with proteins like

peroxisome proliferator-activated receptor gamma (PPARγ) which regulates adipogenesis emphasizing its role in metabolic disorders.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


12 product images



Immunocytochemistry/ Immunofluorescence - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SCD1 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized SK-MEL-28 (Human malignant melanoma) cells labeling SCD1 with ab236868 at 1/100 dilution, followed by ab150077 AlexaFluor® 488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in SK-MEL-28 cell line. Counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) at 1/200 dilution. Nuclear counterstain is DAPI.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is ab150077 AlexaFluor® 488 Goat anti-Rabbit secondary at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human adipose tissue within cardiac muscle tissue using rabbit Anti-SCD1 antibody

Immunohistochemical analysis of paraffin-embedded human adipose tissue within cardiac muscle tissue labeling SCD1 with ab236868 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining in human adipose cells in cardiac muscle (PMID: 15907797) is observed. Counter stained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.


Heat-mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).



Immunocytochemistry/ Immunofluorescence - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SCD1 antibody

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HepG2 (Human hepatocellular carcinoma epithelial cell) cells labeling SCD1 with ab236868 at 1/100 dilution, followed by ab150077 AlexaFluor®488 Goat anti-Rabbit secondary antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in HepG2 cell line. Counterstained with ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) at 1/200 dilution. Nuclear counterstain is DAPI.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is ab150077 AlexaFluor®488 Goat anti-Rabbit secondary at 1/1000 dilution.




Immunocytochemistry/ Immunofluorescence - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-SCD1 antibody

ab236868 staining SCD1 in wild-type HeLa cells, with negative expression in SCD1 knockout HeLa cells. The cells were fixed with 4% formaldehyde (10 min), permeabilised with 0.1% Triton x-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab236868 at 0.04 µg/ml and ab7291, Mouse monoclonal [DM1A] to alpha Tubulin at 0.5 µg/ml. Cells were then incubated with ab150081, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 488), pre-adsorbed at 1/1000 dilution (shown in green) and ab150119, Goat polyclonal Secondary Antibody to Mouse IgG - H&L (Alexa Fluor® 647), pre-adsorbed at 1/1000 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).
Image was acquired with a high-content analyser (Operetta CLS, Perkin Elmer) and a maximum intensity projection of confocal sections is shown.


This product also work with 100% methanol (5 min) fixation under the same testing conditions.



Flow Cytometry (Intracellular) - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 Flow Cytometry (Intracellular) staining using rabbit Anti-SCD1 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed, 90% methanol permeabilized HepG2 (Human hepatocellular carcinoma epithelial cell) cell line labeling SCD1 with ab236868 at 1/500 dilution (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 was used as the secondary antibody.



Immunoprecipitation - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 was immunoprecipitated from 10 µg of HepG2 (Human hepatocellular carcinoma epithelial cell) whole cell lysate with ab236868 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab236868 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used for detection at 1/5000 dilution.
Lane 1: HepG2 whole cell lysate 10 µg (input).
Lane 2: ab236868 IP in HepG2 whole cell lysate (+).
Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab236868 in HepG2 whole cell lysate (-).


Blocking and dilution buffer: 5% NFDM/TBST.

Exposure time: 30 seconds.

The expression profile observed is consistent with what has been described in the literature (PMID: 20876744; PMID: 9843580; PMID: 17449569). The full-length protein migrates at 37 kDa; the 28 kDa fragment may represent an SCD1 cleavage product.

All lanes:
Immunoprecipitation - Anti-SCD1 antibody [EPR21963] (ab236868)

Predicted band size: 42 kDa



Immunoprecipitation - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 was immunoprecipitated from 0.35 mg of HEK-293 (Human embryonic kidney epithelial cell) whole cell lysate with ab236868 at 1/30 dilution. Western blot was performed from the immunoprecipitate using ab236868 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used for detection at 1/5000 dilution.
Lane 1: HEK-293 whole cell lysate 10 µg (input).
Lane 2: ab236868 IP in HEK-293 whole cell lysate (+).
Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab236868 in HEK-293 whole cell lysate (-).


Blocking and dilution buffer: 5% NFDM/TBST.

Exposure time: 30 seconds.

The expression profile observed is consistent with what has been described in the literature (PMID: 20876744; PMID: 9843580; PMID: 17449569). The full-length protein migrates at 37 kDa; the 28 kDa fragment may represent an SCD1 cleavage product.

All lanes:
Immunoprecipitation - Anti-SCD1 antibody [EPR21963] (ab236868)

Predicted band size: 42 kDa




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SCD1 antibody

Immunohistochemical analysis of paraffin-embedded mouse adipose tissue of stomach tissue labeling SCD1 with ab236868 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining in adipose cells in mouse stomach (PMID: 11500518) is observed. Counter stained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Heat-mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-SCD1 antibody

Immunohistochemical analysis of paraffin-embedded rat adipose tissue of pancreas tissue labeling SCD1 with ab236868 at 1/2000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) ready to use. Positive staining in adipose cells in rat pancreas (PMID: 11500518) is observed. Counter stained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) ready to use.

Heat-mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).



Western blot - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 Western blot staining using rabbit Anti-SCD1 antibody

Blocking/dilution buffer: 5% NFDM/TBST.
The expression profile observed is consistent with what has been described in the literature (PMID: 20876744; PMID: 9843580; PMID: 17449569). The full-length protein migrates at 37 kDa; the 28 kDa fragment may represent an SCD1 cleavage product.

We recommend that customers **do not boil** samples to prevent protein aggregation.

All lanes:
Western blot - Anti-SCD1 antibody [EPR21963] (ab236868) at 1/1000 dilution

Lane 1:
HepG2 (human hepatocellular carcinoma epithelial cell), whole cell lysate at 20 µg

Lane 2:
HEK-293 (human embryonic kidney epithelial cell), whole cell lysate at 20 µg


Lane 3:
SK-MEL-28 (human malignant melanoma), whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/50000 dilution

Predicted band size: 42 kDa

Exposure time: 4s



Western blot - Anti-SCD1 antibody [EPR21963] (ab236868)


SCD1 Western blot staining using rabbit Anti-SCD1 antibody

ab236868 was shown to react with SCD1 in wild-type HeLa cells in Western blot with loss of signal observed in SCD1 knockout cell line ab265220. Wild-type HeLa and SCD1 knockout cell lysates were subjected to SDS-PAGE. Membranes were blocked in 5% milk in TBST for 1 hr before incubation with ab236868 overnight at 4 °C at a 1/1000 dilution. Blots were incubated with secondary antibodies at 0.2 µg/mL before imaging.

These data were provided by YCharOS Inc., an open science company with the mission of characterizing commercially available antibody reagents for all human proteins. Abcam and YCharOS are working together to help address the reproducibility crisis by enabling the life science community to better evaluate commercially available antibodies.
All lanes:
Western blot - Anti-SCD1 antibody [EPR21963] (ab236868) at 1/1000 dilution

Lane 1:
Wild-type HeLa lysate at 35 µg

Lane 2:
SCD1 knock-out HeLa lysate at 35 µg



Western blot - Anti-SCD1 antibody [EPR21963] (ab236868)

SCD1 Western blot staining using rabbit Anti-SCD1 antibody

anes 1-3: Merged signal (red and green). Green - ab236868. Red - loading control ab8245 observed at 50 kDa.
ab236868 Anti-SCD1 antibody [EPR21963] was shown to specifically react with SCD1 in wild-type HeLa cells. Loss of signal was observed when knockout cell line ab265220 (knockout cell lysate ab257658) was used. Wild-type and SCD1 knockout samples were subjected to SDS-PAGE. ab236868 and Anti-tubulin antibody [DM1A] - Loading Control (ab7291) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1 in 10000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-SCD1 antibody [EPR21963] (ab236868) at 1/1000 dilution

Lane 1:
Wild-type HeLa cell lysate at 20 µg

Lane 2:
SCD knockout HeLa cell lysate at 20 µg

Lane 2:

Western blot - Human SCD (SCD1) knockout HeLa cell line (ab265220)

Lane 3:
HepG2 cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) at 1/10000 dilution

Predicted band size: 42 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.