abcam

ab237267

Alexa Fluor® 488 Anti-Heme Oxygenase 1 antibody [EPR18161-128]

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Conjugation	Alexa Fluor® 488
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	pH: 7.4 Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR18161-128
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	1/100
Notes	-

Species	Mouse
Dilution info	1/100
Notes	-

Species	Rat
Dilution info	1/100
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/2500
Notes	This product gave a positive signal in HeLa fixed with 100% methanol (5 min).

Expected

Species	Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Flow Cyt (Intra)

Expected

Species	Human, Mouse, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information HMOX1

Function	Heme oxygenase 1. Catalyzes the oxidative cleavage of heme at the alpha-methene bridge carbon, released as carbon monoxide (CO), to generate biliverdin IXalpha, while releasing the central heme iron chelate as ferrous iron (PubMed:11121422, PubMed:19556236, PubMed:7703255). Affords protection against programmed cell death and this cytoprotective effect relies on its ability to catabolize free heme and prevent it from sensitizing cells to undergo apoptosis (PubMed:20055707). Heme oxygenase 1. (Microbial infection) During SARS-COV-2 infection, promotes SARS-CoV-2 ORF3A-mediated autophagy but is unlikely to be required for ORF3A-mediated induction of reticulophagy. Heme oxygenase 1 soluble form. Catalyzes the oxidative cleavage of heme at the alpha-methene bridge carbon, released as carbon monoxide (CO), to generate biliverdin IXalpha, while releasing the central heme iron chelate as ferrous iron.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Stable for 12 months at -20°C, Store in the dark

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies

Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Heme Oxygenase 1 also known as HO-1 or HMOX1 is an enzyme that plays an important mechanistic role in heme catabolism. It catalyzes the degradation of heme into biliverdin carbon monoxide and free iron. This process involves the cleavage of the heme ring. HO-1 has a molecular weight of approximately 32 kDa. It is widely expressed in numerous tissues but is especially abundant in the liver and spleen. Its expression is induced by heme and other stress stimuli such as heavy metals cytokines and reactive oxygen species.
Biological function summary	Heme Oxygenase 1 serves important protective functions in the body. It is not part of a larger complex but its products such as carbon monoxide and biliverdin have their own biological activities. Carbon monoxide produced by HO-1 has antiflammatory properties and can modulate apoptotic pathways. Biliverdin is reduced to bilirubin which acts as an antioxidant. The enzyme therefore directly influences cellular stress responses and maintains cellular homeostasis through these processes.
Pathways	Heme Oxygenase 1 is integrally involved in oxidative stress response and heme metabolism. It participates in the cellular response to oxidative damage by reducing oxidative stress and promoting cytoprotection. Through its heme degradation activity it is connected with the synthesis of biologically active molecules like bilirubin and carbon monoxide. Heme Oxygenase 1 activity is related to other proteins in oxidative stress pathways such as Nuclear Factor Erythroid 2-Related Factor 2 (Nrf2) which regulates its expression and globins which are sources of heme for HO-1 activity.
Associated diseases and disorders	Heme Oxygenase 1 has been linked to conditions like cardiovascular diseases and neurodegenerative disorders. Its expression can attenuate the severity of atherosclerosis where oxidative stress is an important factor. In neurodegenerative diseases HO-1’s antioxidant properties may provide neuroprotection by mitigating oxidative damage. The protein's interactions with inflammatory cytokines such as Interleukin-6 and tumor necrosis factor-alpha influence its activity in these disease contexts.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-Heme Oxygenase 1 antibody [EPR18161-128] (ab237267)

Heme Oxygenase 1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Heme Oxygenase 1 antibody

ab237267 staining Heme Oxygenase 1 in HeLa cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab237267 at 1/2500 dilution (shown in green) and ab195889, Mouse monoclonal to alpha Tubulin (Alexa Fluor® 594), at 1/250 dilution (shown in red). Nuclear DNA was labeled with DAPI (shown in blue). mage was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.