abcam

ab239376

Anti-DPP4 antibody [EPR22233-135] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR22233-135
Purification technique	Affinity purification Protein A

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	-
Notes	-

Target data

See full target information DPP4

Function	Cell surface glycoprotein receptor involved in the costimulatory signal essential for T-cell receptor (TCR)-mediated T-cell activation (PubMed:10900005, PubMed:10951221, PubMed:11772392, PubMed:17287217). Acts as a positive regulator of T-cell coactivation, by binding at least ADA, CAV1, IGF2R, and PTPRC (PubMed:10900005, PubMed:10951221, PubMed:11772392, PubMed:14691230). Its binding to CAV1 and CARD11 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner (PubMed:17287217). Its interaction with ADA also regulates lymphocyte-epithelial cell adhesion (PubMed:11772392). In association with FAP is involved in the pericellular proteolysis of the extracellular matrix (ECM), the migration and invasion of endothelial cells into the ECM (PubMed:10593948, PubMed:16651416). May be involved in the promotion of lymphatic endothelial cells adhesion, migration and tube formation (PubMed:18708048). When overexpressed, enhanced cell proliferation, a process inhibited by GPC3 (PubMed:17549790). Also acts as a serine exopeptidase with a dipeptidyl peptidase activity that regulates various physiological processes by cleaving peptides in the circulation, including many chemokines, mitogenic growth factors, neuropeptides and peptide hormones such as brain natriuretic peptide 32 (PubMed:10570924, PubMed:16254193). Removes N-terminal dipeptides sequentially from polypeptides having unsubstituted N-termini provided that the penultimate residue is proline (PubMed:10593948). (Microbial infection) Acts as a receptor for human coronavirus MERS-CoV-2.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab239376 is the carrier-free version of ab229018.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity

- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Dipeptidyl peptidase 4 also known as DPP4 or CD26 functions as a serine exopeptidase. This enzyme is involved in cleaving dipeptides from the N-terminus of polypeptides. DPP4 with a mass of approximately 110 kDa is expressed in various tissues such as the liver intestines kidney and lymphocytes. It exists in both a membrane-bound form and a soluble form in blood plasma.
Biological function summary	DPP4 plays a significant role in glucose metabolism and immune function. The enzyme interacts with members of the immune system and regulates proteins such as chemokines growth factors and neuropeptides. DPP4 acts independently and is not known to be part of a larger complex. It modulates biological processes by inactivating peptide hormones and signaling molecules which impacts glucose homeostasis and immune responses.
Pathways	DPP4 impacts the incretin signaling pathway and the immune response pathway. Incretin hormones like glucagon-like peptide-1 (GLP-1) and glucose-dependent insulinotropic polypeptide (GIP) are substrates of DPP4 and play important roles in glucose metabolism. DPP4 also interacts with proteins such as adenosine deaminase (ADA) which impacts T-cell function and immune regulation. The regulation of these pathways by DPP4 influences metabolic and immune health.
Associated diseases and disorders	DPP4 involvement is relevant in type 2 diabetes and immunological disorders. In type 2 diabetes DPP4 inhibitors modulate the incretin hormones improving glycemic control. Proteins like GLP-1 are affected by DPP4 activity connecting it to diabetes management. In autoimmune diseases altered DPP4 expression can impact immune cell functions influencing conditions such as rheumatoid arthritis. Through its effects on ADA and associated pathways DPP4 can influence both metabolic and immune-related disorders.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

1 product image



Flow Cytometry - Anti-DPP4 antibody [EPR22233-135] - BSA and Azide free (ab239376)

DPP4 Flow Cytometry staining using rabbit Anti-DPP4 antibody

Flow cytometric analysis of human primary peripheral blood mononuclear cells (PBMC) labeling CD26 with ab229018 at 1/600 (right) compared with a isotype control Rabbit monoclonal IgG (ab172730) (left). Goat anti rabbit IgG Fc (Dylight® 488, ab98462), at 1/2000 dilution was used as the secondary antibody.
Gated on viable cells.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol and sodium azide (ab229018).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.