abcam

ab240198

# Anti-Histone H2B (acetyl K20) antibody [EPR859] - BSA and Azide free

Recombinant | RabMAb | Advanced Validation

## Key facts

| | |
|---|---|
| **Isotype** | IgG |
| **Host species** | Rabbit |
| **Storage buffer** | pH: 7.2 - 7.4 Constituents: PBS |
| **Form** | Liquid |
| **Clonality** | Monoclonal |
| **Immunogen** | The exact immunogen used to generate this antibody is proprietary information. |
| **Carrier free** | Yes |
| **Clone number** | EPR859 |
| **Purification technique** | Affinity purification Protein A |
| **Concentration** | 0.996 mg/mL The concentration of this product may be batch-dependent<br>Batch concentration finder → |

## Reactivity data

## ChIP

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | - |

| | |
|---|---|
| **Notes** | - |

### Expected

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## WB

### Tested

| | |
|---|---|
| **Species** | Mouse, Human |
| **Dilution info** | - |
| **Notes** | - |

### Expected

| | |
|---|---|
| **Species** | Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## PepArr

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | - |
| **Notes** | - |

### Expected

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## ICC/IF

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | - |
| **Notes** | - |

### Expected

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## Flow Cyt (Intra)

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | - |
| **Notes** | ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody. |

### Expected

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## IHC-P

### Tested

| | |
|---|---|
| **Species** | Human, Mouse, Rat |
| **Dilution info** | - |
| **Notes** | - |

## ChIP-seq

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | - |
| **Notes** | - |

### Expected

| | |
|---|---|
| **Species** | Mouse, Rat |
| **Dilution info** | Use at an assay dependent concentration. |
| **Notes** | - |

## Target data

See full target information H2BC21 acK20

**Function** Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Has broad antibacterial activity. May contribute to the formation of the functional antimicrobial barrier of the colonic epithelium, and to the bactericidal activity of amniotic fluid.

## Storage

| | |
|---|---|
| **Shipped at conditions** | Blue Ice |
| **Appropriate short-term storage conditions** | +4°C |
| **Appropriate long-term storage conditions** | +4°C |
| **Storage information** | Do Not Freeze |

## Notes

ab240198 is the carrier-free version of ab177430.

**Patented technology**

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For

details on our patents, please refer to RabMAb® patents.

**What are the advantages of a recombinant monoclonal antibody?**
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

**Conjugation ready**
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

**Compatibility**
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

## Product promise

**Tested**

We have tested this species and application combination and it works. It is covered by our product promise.

**Expected**

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

**Predicted**

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

**Not recommended**


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


## 7 product images




### Flow Cytometry (Intracellular) - Anti-Histone H2B (acetyl K20) antibody [EPR859] - BSA and Azide free (ab240198)

Histone H2B (acetyl K20) Flow Cytometry (Intracellular) staining using rabbit Anti-Histone H2B (acetyl K20) antibody

Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed HeLa (Human epithelial cells from cervix adenocarcinoma) cells treated with 500ng/ml Trichostatin A for 4 hours, labeling Histone H2B (acetyl K20) with ab177430 at 1/300 dilution (red) compared with a rabbit monoclonal IgG isotype control (black) and a unlabelled control (cells without incubation with primary antibody and secondary antibody; blue). Goat anti rabbit IgG (FITC) at 1/150 dilution was used as the secondary antibody.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab177430).





### Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H2B (acetyl K20) antibody [EPR859] - BSA and Azide free (ab240198)

Histone H2B (acetyl K20) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Histone H2B (acetyl K20) antibody

Immunohistochemical analysis of paraffin-embedded Human colon tissue labeling Histone H2B (acetyl K20) with ab177430 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution. Nucleus staining on Human colon is observed. Counter stained with Hematoxylin.
Negative control: Using PBS instead of primary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab177430).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.





### Immunocytochemistry/ Immunofluorescence - Anti-Histone H2B (acetyl K20) antibody [EPR859] - BSA and Azide free (ab240198)

Histone H2B (acetyl K20) Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Histone H2B (acetyl K20) antibody


Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cells from cervix adenocarcinoma) cells, treated with Trichostatin A (500 ng/ml) for 4 hours or untreated, labeling Histone H2B (acetyl K20) with ab177430 at 1/2000 dilution, followed by Goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/500 dilution (green). Confocal image showing nuclear staining on HeLa cell line is observed. Acetylation level increased after treatment with Trichostatin A (500 ng/ml) for 4 hours. The nuclear counter stain is DAPI (blue). Tubulin is detected with ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution and ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution (red).
The negative controls are as follows;

1. ab177430 at 1/2000 dilution followed by ab150120 (AlexaFluor®594 Goat anti-Mouse secondary) at 1/500 dilution.
2. ab7291 (anti-Tubulin mouse mAb) at 1/1000 dilution followed by ab150077 (Alexa Fluor®488 Goat Anti-Rabbit IgG H&L) at 1/500 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab177430).




## ChIP - Anti-Histone H2B (acetyl K20) antibody [EPR859] - BSA and Azide free (ab240198)

Chromatin was prepared from HeLa (Human epithelial cells from cervix adenocarcinoma) cells according to the Abcam X-ChIP protocol. Cells were fixed with formaldehyde for 10 minutes. The ChIP was performed with 25µg of chromatin, 2µg of ab177430 (blue), and 20µl of Anti rabbit IgG sepharose beads. 2µg of rabbit normal IgG was added to the beads control (yellow). The immunoprecipitated DNA was quantified by real time PCR (Sybr green approach). Primers and probes are located in the first kb of the transcribed region.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab177430).





## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H2B (acetyl K20) antibody [EPR859] - BSA and Azide free (ab240198)

Histone H2B (acetyl K20) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Histone H2B (acetyl K20) antibody

Immunohistochemical analysis of paraffin-embedded Mouse kidney tissue labeling Histone H2B (acetyl K20) with ab177430 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution. Nucleus staining on mouse kidney is observed. Counter stained with Hematoxylin.
Negative control: Using PBS instead of primary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab177430).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.





## Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H2B (acetyl K20) antibody [EPR859] - BSA and Azide free (ab240198)

Histone H2B (acetyl K20) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Histone H2B (acetyl K20) antibody

Immunohistochemical analysis of paraffin-embedded Rat colon tissue labeling Histone H2B (acetyl K20) with ab177430 at 1/1000 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/500 dilution. Nucleus staining on Rat colon is observed. Counter stained with Hematoxylin.
Negative control: Using PBS instead of primary antibody.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab177430).

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




## Peptide Array - Anti-Histone H2B (acetyl K20) antibody [EPR859] - BSA and Azide free (ab240198)

ab177430 was tested in Peptide array against 501 different modified and unmodified histone peptides; each peptide is printed on the array at six concentrations (each in triplicate).
Circle area represents affinity between the antibody and a peptide: all antigen-containing peptides are displayed as red circles, all other peptides as blue circles. The affinity is calculated as area under curve when antibody binding values are plotted against the corresponding peptide concentration. Each circle area is normalized to the peptide with the strongest affinity.
The complete dataset, including full list of all peptides and information on the position of each peptide in the diagram, is available under the Product Protocol section.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab177430).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.