abcam

ab240558

Anti-CD11c antibody [EPR21826] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR21826
Purification technique	Affinity purification Protein A
Concentration	1.031 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Mouse
Dilution info	-

Notes	During IHC-P detection, we recommend using the manual two-step method and selecting polymer-based secondary antibodies to enhance signal intensity.

Flow Cyt

Tested

Species	Mouse
Dilution info	-
Notes	-

Target data

See full target information Itgax

Function	Integrin alpha-X/beta-2 is a receptor for fibrinogen. It recognizes the sequence G-P-R in fibrinogen. It mediates cell-cell interaction during inflammatory responses. It is especially important in monocyte adhesion and chemotaxis. Functions as a receptor of the erythrocyte-specific adhesion molecule ICAM4 and mediates erythrophagocytosis.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab240558 is the carrier-free version of ab219799.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD11c also known as ITGAX is a subunit of integrin proteins that plays a pivotal role in immune cell function. This protein has a molecular mass of about 150 kDa and is primarily expressed on the surface of dendritic cells monocytes and macrophages. It serves as a marker for these cells commonly utilized in applications like CD11c flow cytometry and CD11c IHC. Researchers often study this target using CD11c antibodies to explore immune system dynamics and cell surface characteristics.
Biological function summary	CD11c functions as an integrin alpha X subunit forming a heterodimer with the beta 2 subunit known as ITGB2. This complex is important for cell adhesion and migration processes. It mediates the binding of leukocytes to endothelial cells facilitating their migration from the bloodstream to sites of inflammation or infection. CD11c's presence on immune cells makes it essential for presenting antigens and initiating immune responses therefore having critical implications for immunity.
Pathways	CD11c plays a significant role in the immune system particularly in the leukocyte adhesion and migration pathways. It partners with proteins like ITGB2 to mediate these functions. The signaling pathways involving CD11c and related molecules regulate leukocyte trafficking and activation ensuring proper localization and function at sites of damage or infection. These pathways are essential for maintaining immune homeostasis and prompt responses to pathogens.
Associated diseases and disorders	CD11c has significant connections to autoimmune diseases and chronic inflammatory conditions. Aberrant CD11c expression or activity can contribute to the pathogenesis of conditions like rheumatoid arthritis where leukocyte migration and persistent inflammation play a role. Studies also link CD11c with proteins involved in these disorders highlighting its importance in disease progression and potential as a therapeutic target. The understanding of CD11c's

role in these conditions can guide the development of treatments aimed at modulating immune responses.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

2 product images



Flow Cytometry - Anti-CD11c antibody [EPR21826] - BSA and Azide free (ab240558)


CD11c Flow Cytometry staining using rabbit Anti-CD11c antibody

Flow cytometric analysis of mouse primary splenocytes labeling CD11c with ab219799 at a 1/50 dilution.
Cells were stained with BV421 conjugated CD11b and rabbit IgG (Left panel) or CD11c (Right panel).

Goat anti rabbit IgG (Alexa Fluor® 488, ab150077), at 1/2000 dilution was used as the secondary antibody.

Gated on viable cells.

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab219799).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD11c antibody [EPR21826] - BSA and Azide free (ab240558)

CD11c Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD11c antibody

Immunohistochemical analysis of paraffin-embedded mouse spleen tissue labeling CD11c with ab219799 at 1/100 dilution, followed by ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on dendritic cells of mouse spleen (PMID: 20194631, PMID: 15795237) is observed. Counterstained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol and sodium azide (ab219799).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.