abcam

ab241941

# violetFluor™ 450 Anti-CD3 antibody [UCHT1]

Anti-CD3 antibody [UCHT1] conjugated to violetFluor™ 450 - suitable for flow cytometry, and reacts with human samples. Clone [UCHT1] is a mouse monoclonal antibody and one of the most commonly used clones for flow cytometry.

Design your panel with our interactive flow cytometry panel builder. Click here.

## Key facts

| | |
|---|---|
| **Isotype** | IgG1 |
| **Host species** | Mouse |
| **Conjugation** | violetFluor™ 450 |
| **Excitation/Emission** | Ex: 405nm, Em: 450nm |
| **Storage buffer** | pH: 7.2 Preservative: 0.09% Sodium azide Constituents: 0.87% Sodium chloride, 0.12% Sodium dihydrogen phosphate, 0.1% Gelatin |
| **Form** | Liquid |
| **Clonality** | Monoclonal |
| **Clone number** | UCHT1 |
| **Purification technique** | Affinity purification |
| **Specificity** | ab241941 is specific for human CD3e, also known as CD3 epsilon. |
| **Light chain type** | kappa |
| **Concentration** | 0.1 mg/mL The concentration of this product may be batch-dependent<br>Batch concentration finder → |
| **Purification notes** | Purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. |

## Reactivity data

## Flow Cyt

### Tested

| | |
|---|---|
| **Species** | Human |
| **Dilution info** | 5 µL for $10^5$-$10^8$ Cells |
| **Notes** | Use 5 µl (0.5 µg)/test. |

## Target data

See full target information CD3D

**Function**

Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD247/CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain. Upon TCR engagement, these motifs become phosphorylated by Src family protein tyrosine kinases LCK and FYN, resulting in the activation of downstream signaling pathways (PubMed:2470098). In addition of this role of signal transduction in T-cell activation, CD3D plays an essential role in thymocyte differentiation. Indeed, participates in correct intracellular TCR-CD3 complex assembly and surface expression. In absence of a functional TCR-CD3 complex, thymocytes are unable to differentiate properly. Interacts with CD4 and CD8 and thus serves to establish a functional link between the TCR and coreceptors CD4 and CD8, which is needed for activation and positive selection of CD4 or CD8 T-cells (PubMed:12215456).

## Storage

| | |
|---|---|
| **Shipped at conditions** | Blue Ice |
| **Appropriate short-term storage conditions** | +4°C |
| **Appropriate long-term storage conditions** | +4°C |
| **Storage information** | Store in the dark |

# Product promise

**Tested**

We have tested this species and application combination and it works. It is covered by our product promise.

**Expected**

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

**Predicted**

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

**Not recommended**

We do not recommend this combination. It is not covered by our product promise.

---


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

## 1 product image




### Flow Cytometry - violetFluor™ 450 Anti-CD3 antibody [UCHT1] (ab241941)

CD3 Flow Cytometry staining using mouse Anti-CD3 antibody

Flow cytometric analysis of human peripheral blood lymphocytes labeling CD3 with 5 µl (0.5 µg) ab241941 (solid line) or 0.5 µg violetFluor™ 450 Mouse IgG1 isotype control (dashed line).

---

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.