abcam

ab243091

Anti-IL-18 antibody [EPR19954-188]

Anti-IL-18 antibody [EPR19954-188] (ab243091) is a rabbit monoclonal antibody detecting IL-18 in **Western Blot, Flow Cytometry (Intra), IHC-P**. Suitable for **Human**.

- KO validated for confirmed specificity
- Biophysical QC for unrivalled batch-batch consistency
- Over 20 publications

Recombinant | RabMAb | KO Validated | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR19954-188
Purification technique	Affinity purification Protein A
Concentration	0.486 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

WB

Tested

Species	Human
Dilution info	1/1000
Notes	-

ICC/IF

Not recommended

Species	Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	1/50
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/1000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Target data

See full target information IL18

Function Pro-inflammatory cytokine primarily involved in epithelial barrier repair, polarized T-helper 1 (Th1) cell and natural killer (NK) cell immune responses (PubMed:10653850). Upon binding to IL18R1 and IL18RAP, forms a signaling ternary complex which activates NF-kappa-B, triggering synthesis of inflammatory mediators (PubMed:14528293, PubMed:25500532, PubMed:37993714). Synergizes with IL12/interleukin-12 to induce IFNG synthesis from T-helper 1 (Th1) cells and natural killer (NK) cells (PubMed:10653850). Involved in transduction of

inflammation downstream of pyroptosis: its mature form is specifically released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33883744).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What is this antibody validated in?
Anti-IL-18 antibody [EPR19954-188] (ab243091) is a rabbit recombinant monoclonal antibody and is validated for use in Western Blot (WB), Flow Cytometry (Intra), Flow Cytometry (Flow Cyt), Immunohistochemistry (IHC-P) in Human samples.

What is the molecular weight of IL-18?
Anti-IL-18 [EPR19954-188] (ab243091) specifically detects a band for IL-18 (UniProt: Q14116) at a molecular weight of 22kDa.

Trusted by the scientific community
Anti-IL-18 [EPR19954-188] (ab243091) was first used in a scientific publication in 2018 and has been cited over 20 times in peer-reviewed journals.

Specificity confirmed
The specificity of Anti-IL-18 antibody [EPR19954-188] (ab243091) has been confirmed by Western blot testing in IL18 Knockout HeLa cell line, ab265274.

Other related products
We have a range of other formats of antibody clone [EPR19954-188] also available for your convenience: ab207324, Carrier free - ab222926, Carrier free - ab242590

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Trial sizes available!
Test your antibody or perform pre-screening before committing to a larger quantity. Sold in 20µl. Discover our selection of trial-size antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

6 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IL-18 antibody [EPR19954-188] (ab243091)

IL-18 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IL-18 antibody

Immunohistochemical analysis of paraffin-embedded human liver tissue labeling IL-18 with ab243091 at 1/250 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Cytoplasmic staining on Kupffer cells in human liver (PMID: 19084941). Counter stained with hematoxylin. Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IL-18 antibody [EPR19954-188] (ab243091)

IL-18 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-IL-18 antibody

Immunohistochemical analysis of paraffin-embedded human tonsil tissue labeling IL-18 with ab243091 at 1/250 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Positive staining on subsets of macrophages in tonsil (PMID: 28842466). Counter stained with hematoxylin. Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.


Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).



Flow Cytometry (Intracellular) - Anti-IL-18 antibody [EPR19954-188] (ab243091)

IL-18 Flow Cytometry (Intracellular) staining using rabbit Anti-IL-18 antibody


Intracellular flow cytometric analysis of4% paraformaldehyde-fixed, 90% methanol-permeabilized Jurkat (human T cell leukemia cell line from peripheral blood) (left panel) and PC-3 (human prostate adenocarcinoma cell line) (right panel) cells labeling IL-18 with ab243091 at 1/50 dilution (red) compared with a Rabbit IgG, monoclonal [EPR25A] - Isotype Control (ab172730) (black) and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) (ab150077) at 1/2000 dilution was used as the secondary antibody.
Negative control: Jurkat (PMID 15086390).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-IL-18 antibody [EPR19954-188] (ab243091)

Immunohistochemical analysis of paraffin-embedded human kidney tissue labeling IL-18 with ab243091 at 1/250 dilution, followed by Goat Anti-Rabbit IgG H&L (HRP) Ready to use. Cytoplasmic and nuclear staining on distal convoluted tubules of human kidney (PMID: 17687255). Counter stained with hematoxylin.
Secondary antibody only control: Used PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG H&L (HRP) Ready to use.

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0).



Western blot - Anti-IL-18 antibody [EPR19954-188] (ab243091)

IL-18 Western blot staining using rabbit Anti-IL-18 antibody

Lanes 1-4: Merged signal (red and green). Green - ab243091 observed at 22 kDa. Red - loading control ab8245 observed at 37 kDa.
ab243091 Anti-IL-18 antibody [EPR19954-188] was shown to specifically react with IL-18 in wild-type HeLa cells. Loss of signal was observed when knockout cell line ab265274 (knockout cell lysate ab256952) was used. Wild-type and IL-18 knockout samples were subjected to SDS-PAGE. ab243091 and Anti-alpha Tubulin antibody [EP1332Y] - Microtubule Marker (ab52866) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

All lanes:
Western blot - Anti-IL-18 antibody [EPR19954-188] (ab243091) at 1/1000 dilution

Lane 1:
Wild-type HeLa cell lysate at 20 µg

Lane 2:
IL18 knockout HeLa cell lysate at 20 µg

Lane 2:
Western blot - Human IL18 knockout HeLa cell line (ab265274)

Lane 3:
A431 cell lysate at 20 µg

Lane 4:
Jurkat cell lysate at 20 µg


Secondary

All lanes:
Western blot - Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (ab216773) at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 22 kDa

Observed band size: 22 kDa



Western blot - Anti-IL-18 antibody [EPR19954-188] (ab243091)

Exposure time : Lanes 1 & 2: 3 minutes. Lane 3: 26 seconds.
Blocking and dilution buffer: 5% NFDM/TBST.

All lanes:
Western blot - Anti-IL-18 antibody [EPR19954-188] (ab243091) at 1/1000 dilution

Lane 1:
PC-3 (human prostate adenocarcinoma cell line) whole cell lysate at 20 µg

Lane 2:
HeLa (human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 20 µg

Lane 3:
HaCaT (Human keratinocyte cell line) whole cell lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 22 kDa

Observed band size: 14 kDa, 22 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.