abcam

ab243650

Anti-IL-2 antibody [EPR16615-341]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR16615-341
Purification technique	Affinity purification Protein A
Concentration	0.455 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

IP

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Mouse, Human, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

WB

Not recommended

Species	Human, Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information Il2

Function	Cytokine produced by activated CD4-positive helper T-cells and to a lesser extend activated CD8-positive T-cells and natural killer (NK) cells that plays pivotal roles in the immune response and tolerance (PubMed:14614860, PubMed:9814585). Binds to a receptor complex composed of either the high-affinity trimeric IL-2R (IL2RA/CD25, IL2RB/CD122 and IL2RG/CD132) or the low-affinity dimeric IL-2R (IL2RB and IL2RG). Interaction with the receptor leads to oligomerization and conformation changes in the IL-2R subunits resulting in downstream signaling starting with phosphorylation of JAK1 and JAK3. In turn, JAK1 and JAK3 phosphorylate the receptor to form a docking site leading to the phosphorylation of several substrates including STAT5 (PubMed:14614860, PubMed:27018889). This process leads to activation of several pathways including STAT, phosphoinositide-3-kinase/PI3K and mitogen-activated protein kinase/MAPK pathways. Functions as a T-cell growth factor and can increase NK-cell cytolytic activity as well. Promotes strong proliferation of activated B-cells and subsequently immunoglobulin production. Plays a pivotal role in regulating the adaptive immune system by controlling the survival and proliferation of regulatory T-cells, which are required for the maintenance of immune tolerance (PubMed:14614860). Moreover, participates in the differentiation and homeostasis of effector T-cell subsets, including Th1, Th2, Th17 as well as memory CD8-positive T-cells (PubMed:9814585).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


4 product images



Flow Cytometry (Intracellular) - Anti-IL-2 antibody [EPR16615-341] (ab243650)

IL-2 Flow Cytometry (Intracellular) staining using rabbit Anti-IL-2 antibody


Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized EL4.IL-2 (mouse lymphoma T lymphocyte) treated with cell stimulation cocktail (80nM PMA + 1.34uM Ionomycin + 10.6uM Brefeldin A + 2uM Monensin) for 6 hours (Red)/ Untreated control (Green) cells labelling IL-2 with ab243650 at 1/50 dilution (1ug) (Red) compared with a Rabbit monoclonal IgG (ab172730) (Black) isotype control and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.



Flow Cytometry (Intracellular) - Anti-IL-2 antibody [EPR16615-341] (ab243650)

IL-2 Flow Cytometry (Intracellular) staining using rabbit Anti-IL-2 antibody


Intracellular flow cytometric analysis of 2% paraformaldehyde fixed 0.1% Tween-20 permeabilized Mouse splenocytes treated with cell stimulation cocktail (80nM PMA+1.34uM Ionomycin+10.6uM Brefeldin A+2uM Monensin) for 6 hours (Right)/ Untreated control (Left). cells labelling IL-2 with ab243650 at 1/500 dilution (1ug)/ Left and Right (Red) compared with a isotype control. A Goat anti rabbit IgG (Alexa Fluor® 647, ab150079) at 1/2000 dilution was used as the secondary antibody. Cells were surface stained with anti-CD3 conjugated to FITC. Then fixed with 2% PFA for 10min followed by intracellularly stained with ab243650. Gated on lymphocytes population. Data was kindly provided by an anonymous collaborator.



Immunocytochemistry/ Immunofluorescence - Anti-IL-2 antibody [EPR16615-341] (ab243650)

IL-2 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-IL-2 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized EL4.IL-2 cells labelling IL-2 with ab243650 at 1/50 (9.1 ug/ml) dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green) Confocal image showing cytoplasmic staining in EL4.IL-2 cell line treated with Cell Stimulation Cocktail (80nM PMA + 1.34uM Ionomycin + 10.6uM Brefeldin A + 2uM Monensin) for 6 h. ab195889 Anti-alpha Tubulin mouse monoclonal antibody - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.



Flow Cytometry (Intracellular) - Anti-IL-2 antibody [EPR16615-341] (ab243650)

IL-2 Flow Cytometry (Intracellular) staining of Mouse splenocytes using rabbit Anti-IL-2 antibody

Flow cytometry staining of C57 BL/6 mouse splenocytes (top) or C57 BL/6 mouse splenocytes treated with 80nM PMA/TPA, 1.34uM Ionomycin and 1µg/mL Brefeldin A for 18 Hours (bottom) with ab243650 (right) or Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (left). Cells were fixed and permeabilised with BD Cytofix/Cytoperm™ for 20 min. Cells were incubated for

30min on ice in 1x PBS containing 10µg/ml anti CD16/CD32 and 10% normal goat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody ab243650 or Recombinant Rabbit IgG, monoclonal [EPR25A] - Isotype Control (1x 10^6in 100 µl at 1.0 µg/ml (1/500)) for 30min at 22°C. The cells were simultaneously stained with CD4.

The secondary antibody Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) preadsorbed was incubated at 1/4000 for 30min at 22°C.

Acquisition of >30000 events were collected using a 50 mW Blue laser (488nm) and 525/40 bandpass filter. Events were gated on live cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.