abcam

ab243925

Anti-CD16 antibody [SP175] - BSA and Azide free

Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925) is a rabbit recombinant monoclonal antibody provided in a PBS only buffer for easy conjugation detecting CD16 in **Flow Cytometry (Intra), IHC-P**. Suitable for **Human**.

- BSA, sodium azide, and glycerol-free for easy conjugation
- Biophysical QC for unrivalled batch-batch consistency

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	SP175
Purification technique	Affinity purification Protein A/G
Concentration	1.04 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	Purified from TCS by protein A/G.

Reactivity data

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	-
Notes	Antigen retrieval: Boil tissue section in EDTA buffer, pH 8.0 for 10 min followed by cooling at room temperature for 20 min.

Target data

See full target information FCGR3A

Function

Receptor for the invariable Fc fragment of immunoglobulin gamma (IgG). Optimally activated upon binding of clustered antigen-IgG complexes displayed on cell surfaces, triggers lysis of antibody-coated cells, a process known as antibody-dependent cellular cytotoxicity (ADCC). Does not bind free monomeric IgG, thus avoiding inappropriate effector cell activation in the absence of antigenic trigger (PubMed:11711607, PubMed:21768335, PubMed:22023369, PubMed:24412922, PubMed:25786175, PubMed:25816339, PubMed:28652325, PubMed:8609432, PubMed:9242542). Mediates IgG effector functions on natural killer (NK) cells. Binds antigen-IgG complexes generated upon infection and triggers NK cell-dependent cytokine production and degranulation to limit viral load and propagation. Involved in the generation of memory-like adaptive NK cells capable to produce high amounts of IFNG and to efficiently eliminate virus-infected cells via ADCC (PubMed:24412922, PubMed:25786175). Regulates NK cell survival and proliferation, in particular by preventing NK cell progenitor apoptosis (PubMed:29967280, PubMed:9916693). Following the engagement of antigen-IgG complexes, triggers phosphorylation of immunoreceptor tyrosine-based activation motif (ITAM)-containing adapters with subsequent activation of phosphatidylinositol 3-kinase signaling and sustained elevation of intracellular calcium that ultimately drive NK cell activation. The ITAM-dependent signaling coupled to receptor phosphorylation by PKC mediates robust intracellular calcium flux that leads to production of pro-inflammatory cytokines, whereas in the absence of receptor phosphorylation it mainly activates phosphatidylinositol 3-kinase signaling leading to cell degranulation (PubMed:1825220, PubMed:23024279, PubMed:2532305). Costimulates NK cells and trigger lysis of target cells independently of IgG binding (PubMed:10318937, PubMed:23006327).

Mediates the antitumor activities of therapeutic antibodies. Upon ligation on monocytes triggers TNFA-dependent ADCC of IgG-coated tumor cells (PubMed:27670158). Mediates enhanced opsonisation and ADCC in response to afucosylated IgGs (PubMed:34485821, PubMed:28566370). (Microbial infection) Involved in Dengue virus pathogenesis via antibody-dependent enhancement (ADE) mechanism. Secondary infection with Dengue virus triggers elevated levels of afucosylated non-neutralizing IgG1s with reactivity to viral envelope/E protein. Viral antigen-IgG1 complexes bind with high affinity to FCGR3A, facilitating virus entry in myeloid cells and subsequent viral replication.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

What is this antibody validated in?
Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925) is a rabbit recombinant monoclonal antibody and is validated for use in Flow Cytometry (Intra), Flow Cytometry (Flow Cyt), Immunohistochemistry (IHC-P) in Human samples.

Collaboration
Anti-CD16 [SP175] - BSA and Azide free (ab243925) is a clone from the portfolio of Spring Bioscience (Roche) SP clones which have been optimised for immunohistochemistry (IHC).

Other related products
We have a range of other formats of antibody clone [SP175] also available for your convenience: ab183354, Carrier free - ab243925, ab279462, PE - ab305743, APC - ab305744, HRP - ab305745, ab308605, Carrier free - ab308606, ab308607, Carrier free - ab308608, Alkaline Phosphatase - ab308988, Alexa Fluor® 488 - ab309932, Alexa Fluor® 647 - ab310299, Alexa Fluor® 594 - ab310761, Alexa Fluor® 555 - ab312288, Alexa Fluor® 568 - ab312783, Alexa Fluor® 750 - ab321276

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD16 also known as FCGR3A or CD16A protein is a low-affinity receptor for the Fc region of IgG. It weighs approximately 50-70 kDa and exists on the surface of several immune cells such as natural killer (NK) cells some populations of T cells and macrophages. The expression of CD16 mediates various cellular responses including antibody-dependent cellular cytotoxicity (ADCC) where it acts by engaging with immune complexes. Researchers often use anti-CD16 antibodies in flow cytometry to analyze the presence and quantity of CD16-expressing cells.
Biological function summary	This receptor plays a significant role in immune system regulation by contributing to the activation of immune effector cells. CD16 is not part of a larger complex but it operates in concert with CD3 proteins to trigger intracellular signaling pathways upon engagement with its ligand. Its function is important for the antibody-dependent clearance of infected or malignant cells from the body representing an essential linkage between innate and adaptive immunity.
Pathways	Signaling involving CD16 is integrated into the immune response mechanisms that include the Fc gamma receptor-mediated phagocytosis pathway. This pathway involves interactions with IgG antibodies and complements the activity of other Fc gamma receptors such as CD32 and CD64. Through these interactions CD16 enhances phagocytic activity and stimulates the secretion of cytokines and chemokines important for orchestrating an effective immune response.
Associated diseases and disorders	CD16 plays an important role in conditions like autoimmune diseases and certain cancers. Its expression and function are important in the pathophysiology of systemic lupus erythematosus where an imbalance in immune complex clearance leads to tissue damage. In oncology CD16 aids in the immune system's recognition and elimination of tumor cells working alongside proteins such as CD3 in modulating the body's anti-tumor response. Understanding CD16's involvement in these diseases helps in developing therapeutic strategies that harness the immune system's power such as monoclonal antibody therapies targeting CD16.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


12 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody


Immunohistochemical analysis of paraffin embedded human liver hepatocellular carcinoma tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody


Immunohistochemical analysis of paraffin embedded human uterus tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody


Immunohistochemical analysis of paraffin embedded human HK lymphoma tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody


Immunohistochemical analysis of paraffin embedded human placenta tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody


Immunohistochemical analysis of paraffin embedded human lung tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody


Immunohistochemical analysis of paraffin embedded human spleen tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Flow Cytometry - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Flow Cytometry staining using rabbit Anti-CD16 antibody


Flow cytometric analysis of Jurkat (Human T cell leukemia cell line from peripheral blood) cells labeling CD16 with ab183354 at a 1/100 dilution (green) compared to a negative control Rabbit IgG (blue).
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody


Immunohistochemical analysis of paraffin embedded human pancreatic adenocarcinoma tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody


Immunohistochemical analysis of paraffin embedded human B cell lymphoma tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody


Immunohistochemical analysis of paraffin embedded human liver tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody

Immunohistochemical analysis of paraffin embedded human lung adenocarcinoma tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD16 antibody [SP175] - BSA and Azide free (ab243925)

CD16 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD16 antibody

Immunohistochemical analysis of paraffin embedded human thymus tissue labeling CD16 with ab183354 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183354).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.