abcam

ab244557

Anti-IL-2 Receptor alpha antibody [OX39] - BSA and Azide free

Key facts

Isotype	IgG1
Host species	Mouse
Storage buffer	Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	OX39
Purification technique	Affinity purification
Light chain type	kappa
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Rat
Dilution info	0.2 µg/mL

Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

IHC-Fr

Tested

Species	Human
Dilution info	5 µg/mL
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Target data

See full target information IL2RA

Function	Receptor for interleukin-2. The receptor is involved in the regulation of immune tolerance by controlling regulatory T cells (TREGs) activity. TREGs suppress the activation and expansion of autoreactive T-cells.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	+4°C
Aliquoting information	Upon delivery aliquot
Storage information	Do Not Freeze

Notes

ab244557 is the carrier-free version of ab6411.

Want a custom formulation?

This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com

Conjugation ready

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

2 product images



Immunohistochemistry (Frozen sections) - Anti-IL-2 Receptor alpha antibody [OX39] - BSA and Azide free (ab244557)


IL-2 Receptor alpha Immunohistochemistry (Frozen sections) staining using mouse Anti-IL-2 Receptor alpha antibody

This data was developed using the same antibody clone in a different buffer formulation (ab6411). IHC image of IL2 receptor alpha staining in a section of frozen normal rat spleen.

The section was fixed using 10% formaldehyde in 1XPBS for 10 minutes. No antigen retrieval step was performed prior to staining. Non-specific protein-protein interactions were then blocked in TBS containing 0.025% (v/v) Triton X-100, 0.3M glycine and 1% (w/v) BSA for 1h at room temperature. The section was then incubated overnight at +4°C in TBS containing 0.025% (v/v) Triton X-100 and 1% (w/v) BSA with ab6411 at 5µg/ml and ab16669 (Rabbit monoclonal [SP7] to CD3) at 1/150. The section was then incubated with ab150117 (Goat Anti-Mouse IgG H&L (Alexa Fluor® 488), 1/1000)) (shown in green) and ab150080 Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 594), 1/1000) (shown in red) for 1 hour at room temperature. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM. The secondary-only control insert image is taken from an identical assay without primary antibody. The section was then mounted using Fluoromount®.

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

For IHC staining systems (automated and non-automated), customers should optimize variable parameters such as antibody concentrations and incubation times.



Flow Cytometry - Anti-IL-2 Receptor alpha antibody [OX39] - BSA and Azide free (ab244557)

IL-2 Receptor alpha Flow Cytometry staining using mouse Anti-IL-2 Receptor alpha antibody

This data was developed using the same antibody clone in a different buffer formulation (ab6411). Lewis rat splenocytes (top) or Lewis rat splenocytes treated with 5 µg/ml ConA for 3 days (bottom) were stained with ab6411 (right) or mouse IgG1 kappa (ab170190) isotype (left).

Splenocytes were incubated for 30 min on ice in 1x PBS containing 10 % rat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody (ab6411) or mouse IgG1 kappa (ab170190) isotype (1x 10^6 in 100 µl at 0.2 µg/ml) for 30 min on ice.

The secondary antibody Goat anti-mouse IgG H&L (Alexa Fluor ® 488, pre-adsorbed) (ab150117) was used at 1/2000 dilution for 30 min on ice.

The cells were simultaneously stained with CD4.

Acquisition of >30,000 events were collected using a 50 mW Blue laser (488nm) and 530/30 bandpass filter. Events were gated on CD3 positive T cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.