abcam

ab244581

Anti-CD44 antibody [OX50] - BSA and Azide free

Key facts

Isotype	IgG1
Host species	Mouse
Storage buffer	Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	OX50
Purification technique	Affinity purification
Light chain type	kappa
Concentration	1 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Rat
Dilution info	0.2 µg/mL
Notes	-

Target data

See full target information CD44

Function	Cell-surface receptor that plays a role in cell-cell interactions, cell adhesion and migration, helping them to sense and respond to changes in the tissue microenvironment (PubMed:16541107, PubMed:19703720, PubMed:22726066). Participates thereby in a wide variety of cellular functions including the activation, recirculation and homing of T-lymphocytes, hematopoiesis, inflammation and response to bacterial infection (PubMed:7528188). Engages, through its ectodomain, extracellular matrix components such as hyaluronan/HA, collagen, growth factors, cytokines or proteases and serves as a platform for signal transduction by assembling, via its cytoplasmic domain, protein complexes containing receptor kinases and membrane proteases (PubMed:18757307, PubMed:23589287). Such effectors include PKN2, the RhoGTPases RAC1 and RHOA, Rho-kinases and phospholipase C that coordinate signaling pathways promoting calcium mobilization and actin-mediated cytoskeleton reorganization essential for cell migration and adhesion (PubMed:15123640). Upon interaction with LGALS9 ligand, activates downstream signaling components including LCK, ERK and MAPK to promotes NK cell activation (PubMed:37006235). (Microbial infection) Promotes foot-and-mouth disease virus internalization via macropinocytosis.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Aliquoting information	Upon delivery aliquot
Storage information	Do Not Freeze

Notes

Want a custom formulation?
This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

1 product image



Flow Cytometry - Anti-CD44 antibody [OX50] - BSA and Azide free (ab244581)

CD44 Flow Cytometry staining using mouse Anti-CD44 antibody

This data was developed using the same antibody clone in a different buffer formulation (ab238465)
Lewis rat splenocytes stained with ab238465 (right) or mouse IgG1κ (ab170190) isotype (left). Lewis rat splenocytes were incubated for 30 min on ice in 1x PBS / 10 % rat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody (ab238465) or mouse IgG1κ (ab170190) ($1x^{106}$ in 100 µl at 0.2 µg/ml) for 30 min on ice.

The secondary antibody Goat anti-mouse IgG H&L (Alexa Fluor ® 488, pre-adsorbed) (ab150117) was used at 1/2000 dilution for 30 min on ice. The cells were simultaneously stained with CD3 antibody.

Acquisition of >30,000 events were collected using a 50 mW Blue laser (488nm) and 530/30 bandpass filter. Events were gated on viable lymphocytes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.