abcam

ab246415

APC Anti-SIGN-R1 antibody [22D1]

Anti-SIGN-R1 antibody [22D1] conjugated to APC - suitable for flow cytometry, and reacts with mouse samples. Clone [22D1] is a Armenian Hamster monoclonal antibody.

Design your panel with our interactive flow cytometry panel builder. Click here.

Key facts

Isotype	IgG
Host species	Armenian hamster
Conjugation	APC
Excitation/Emission	Ex: 650nm, Em: 660nm
Storage buffer	pH: 7.2 Preservative: 0.09% Sodium azide Constituents: 0.87% Sodium chloride, 0.16% Sodium phosphate, 0.1% Gelatin
Form	Liquid
Clonality	Monoclonal
Clone number	22D1
Purification technique	Affinity purification
Concentration	0.2 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →
Purification notes	ab246415 was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation.

Reactivity data

Flow Cyt

Tested

Species	Mouse
Dilution info	-
Notes	0.25 µg

Target data

See full target information CD209

Function

Pathogen-recognition receptor expressed on the surface of immature dendritic cells (DCs) and involved in initiation of primary immune response. Thought to mediate the endocytosis of pathogens which are subsequently degraded in lysosomal compartments. The receptor returns to the cell membrane surface and the pathogen-derived antigens are presented to resting T-cells via MHC class II proteins to initiate the adaptive immune response. On dendritic cells (DCs) it is a high affinity receptor for ICAM2 and ICAM3 by binding to mannose-like carbohydrates (PubMed:10721994, PubMed:10721995, PubMed:11017109, PubMed:12574325). May act as a DC rolling receptor that mediates transendothelial migration of DC presursors from blood to tissues by binding endothelial ICAM2 (PubMed:11017109). Forms a first contact between DC and resting T cell, througth ICAM3 binding, facilitating the downstream DC-T cell clustering process and DC-induced proliferation of resting T Cells (PubMed:10721994, PubMed:10721995). (Microbial infection) Acts as an attachment receptor for HIV-1 and HIV-2. (Microbial infection) Acts as an attachment receptor for Ebolavirus. (Microbial infection) Acts as an attachment receptor for Cytomegalovirus. (Microbial infection) Acts as an attachment receptor for HCV. (Microbial infection) Acts as an attachment receptor for Dengue virus. (Microbial infection) Acts as an attachment receptor for Measles virus. (Microbial infection) Acts as an attachment receptor for Herpes simplex virus 1. (Microbial infection) Acts as an attachment receptor for Influenzavirus A. (Microbial infection) Acts as an attachment receptor for SARS-CoV. (Microbial infection) Acts as an attachment receptor for Japanese encephalitis virus. (Microbial infection) Acts as an attachment receptor for Lassa virus (PubMed:23966408). Acts as an attachment receptor for Marburg virusn. (Microbial infection) Acts as an attachment receptor for Respiratory syncytial virus. (Microbial infection) Acts as an attachment receptor for Rift valley fever virus and uukuniemi virus. (Microbial infection) Acts as an attachment receptor for West-nile virus. (Microbial infection) Probably recognizes in a calcium-dependent manner high mannose N-linked oligosaccharides in a variety of bacterial pathogen antigens, including Leishmania pifanoi LPG, Lewis-x antigen in Helicobacter pylori LPS, mannose in Klebsiella pneumonae LPS, di-mannose and tri-mannose in Mycobacterium tuberculosis ManLAM and Lewis-x antigen in Schistosoma mansoni SEA (PubMed:16379498). Recognition of M.tuberculosis by dendritic cells occurs partially via this molecule (PubMed:16092920, PubMed:21203928).

Storage

Shipped at conditions	Blue Ice

Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Store in the dark

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Flow Cytometry - APC Anti-SIGN-R1 antibody [22D1] (ab246415)

SIGN-R1 Flow Cytometry staining using armenian hamster Anti-SIGN-R1 antibody

C57Bl/6 splenocytes were stained with PE Anti-Mouse CD11b and 0.25 µg ab246415 (Right panel) or 0.25 µg APC Armenian Hamster IgG isotype control (Left panel).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.