abcam

ab246508

Anti-BST2/Tetherin antibody [EPR23597-202]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23597-202
Purification technique	Affinity purification Protein A
Concentration	0.625 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Mouse
Dilution info	1/2000
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

Flow Cyt

Tested

Species	Mouse
Dilution info	1/500
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

WB

Not recommended

Species	Mouse, Rat, Human
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Mouse
Dilution info	1/100
Notes	-

Not recommended

Species	Human, Rat
Dilution info	-
Notes	-

Target data

See full target information Bst2

Function IFN-induced antiviral host restriction factor which efficiently blocks the release of diverse mammalian enveloped viruses by directly tethering nascent virions to the membranes of infected cells. Acts as a direct physical tether, holding virions to the cell membrane and linking virions to each other. The tethered virions can be internalized by endocytosis and subsequently degraded or they can remain on the cell surface. In either case, their spread as cell-free virions is restricted. Its target viruses belong to diverse families, including retroviridae: human immunodeficiency virus type 1 (HIV-1), mouse mammary tumor virus (MMTV) and murine leukemia virus (MLV), filoviridae: ebola virus (EBOV), arenaviridae: lassa virus (LASV), and rhabdoviridae: vesicular stomatitis virus (VSV). Can inhibit cell surface proteolytic activity of MMP14 causing decreased activation of MMP15 which results in inhibition of cell growth and migration. Can stimulate signaling by LILRA4/ILT7 and consequently provide negative feedback to the production of IFN by plasmacytoid dendritic cells in response to viral infection. Plays a role in the organization of the subapical actin cytoskeleton in polarized epithelial cells.

Storage

Shipped at conditions Conditional Ambient

Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

7 product images




Flow Cytometry - Anti-BST2/Tetherin antibody [EPR23597-202] (ab246508)

BST2/Tetherin Flow Cytometry staining using rabbit Anti-BST2/Tetherin antibody

Flow cytometric analysis of Mouse splenocyte cells labelling BST2/Tetherin with ab246508 at 1/500 dilution (Right) compared with a Rabbit monoclonal IgG (ab172730) isotype control (Left).A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.
Cells were stained with rabbit IgG (Left) or ab246508 (Right). Then stained with anti-CD45R conjugated to Alexa Fluor® 647.

Gated on viable cells.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-BST2/Tetherin antibody [EPR23597-202] (ab246508)

BST2/Tetherin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-BST2/Tetherin antibody

Immunohistochemical analysis of paraffin-embedded Mouse spleen tissue labeling BST2/Tetherin with ab246508 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on mouse spleen (PMID: 19903902).The section was incubated with ab246508 for 30 mins at room temperature.The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins.




Immunocytochemistry/ Immunofluorescence - Anti-BST2/Tetherin antibody [EPR23597-202] (ab246508)

BST2/Tetherin Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-BST2/Tetherin antibody

Immunofluorescent analysis of 100% methanol-fixed EL.4 cells labelling BST2/Tetherin with ab246508 at 1/100 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing cytoplasmic staining in EL.4 cell line. 100% methanol fixation is recommended. ab195889 Anti-alpha Tubulin antibody (Alexa Fluor® 594)

was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-BST2/Tetherin antibody [EPR23597-202] (ab246508)

BST2/Tetherin Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-BST2/Tetherin antibody

Immunohistochemical analysis of paraffin-embedded Mouse skeletal muscle tissue labeling BST2/Tetherin with ab246508 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). **Negative control:** No staining on mouse skeletal muscle (PMID: 19903902). The section was incubated with ab246508 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument. Counterstained with Hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).


Heat mediated antigen retrieval with Citrate buffer (pH 6.0, epitope retrieval solution 1) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-BST2/Tetherin antibody [EPR23597-202] (ab246508)

BST2/Tetherin Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-BST2/Tetherin antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized Mouse splenocyte cells labelling BST2/Tetherin with ab246508 at 1/100 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing cytoplamic staining in mouse splenocyte ab195889 Anti-alpha Tubulin antibody (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 dilution (Red). The Nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 dilution.



Immunoprecipitation - Anti-BST2/Tetherin antibody [EPR23597-202] (ab246508)

BST2/Tetherin was immunoprecipitated from 0.35 mg Mouse spleen tissue lysate 10ug with ab246508 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab246508 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/5000 dilution. Lane 1: Mouse spleen tissue lysate 10ug
Lane 2: ab246508 IP in Mouse spleentissue lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab246508 in mouse spleen tissue lysate


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 15 seconds

BST2 is type II transmembrane glycoprotein with a molecular mass of 28-40 KD, which is consistent to the literature(PMID: 22520941; PMID: 19737401).
All lanes:
Immunoprecipitation - Anti-BST2/Tetherin antibody [EPR23597-202] (ab246508)

Predicted band size: 20 kDa

Observed band size: 28-40 kDa



Immunoprecipitation - Anti-BST2/Tetherin antibody [EPR23597-202] (ab246508)

BST2/Tetherin was immunoprecipitated from 0.35 mg EL4 (mouse lymphoma T lymphocyte) whole cell lysate with ab246508 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab246508 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP) (ab131366) was used at 1/5000 dilution.
Lane 1: EL4 (mouse lymphoma T lymphocyte), whole cell lysate 10 ug

Lane 2: ab246508 IP in EL4 whole cell lysate

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab246508 in EL4 whole cell lysate.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 15 seconds

BST2 is type II transmembrane glycoprotein with a molecular mass of 28-40 KD, which is consistent to the literature(PMID: 22520941; PMID: 19737401).

All lanes:
Immunoprecipitation - Anti-BST2/Tetherin antibody [EPR23597-202] (ab246508)

Predicted band size: 20 kDa

Observed band size: 28-40 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.