abcam

ab246709

Anti-CD42b antibody [SP219] - Low endotoxin, Azide free

Recombinant RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	SP219
Purification technique	Affinity purification Protein A/G
Purification notes	Purified from TCS by protein A/G. Endotoxin level is less than 1 EU/ml as determined by the TAL test.

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	-

Notes	-

Expected

Species	Mouse
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Dog
Dilution info	-
Notes	-

IHC-Fr

Tested

Species	Mouse
Dilution info	-
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Predicted

Species	Dog
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	-
Notes	ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Species	Human
Dilution info	-
Notes	ab199376 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

Predicted

Species	Dog
Dilution info	-
Notes	-

Target data

See full target information GP1BA

Function	GP-Ib, a surface membrane protein of platelets, participates in the formation of platelet plugs by binding to the A1 domain of vWF, which is already bound to the subendothelium.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab246709 is the carrier-free version of ab183345.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

What does low endotoxin mean?
Our low endotoxin, azide-free formats have low endotoxin level (1 EU/mg, determined by the TAL assay) and are free from azide, to achieve consistent experimental results in functional assays.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


7 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD42b antibody [SP219] - Low endotoxin, Azide free (ab246709)

CD42b Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining of Human spleen using rabbit Anti-CD42b antibody


This data was developed using ab183345, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of formalin fixed paraffin embedded human spleen labelling CD42b with ab183345 at a concentration of 1 µg/ml. The immunostaining was performed on a Ventana DISCOVERY ULTRA (Roche Tissue Diagnostics) instrument with a OptiView DAB IHC Detection Kit. The heat-mediated antigen retrieval was conducted with at 100 °C, pH 8.5 for 32 mins. ab183345 Anti-CD42b antibody [SP219] was incubated for 16 mins, at 37 °C. Sections were counterstained with Hematoxylin II. Image inset shows absence of staining in secondary antibody only control. Customers are encouraged to optimise antigen retrieval conditions, antibody concentration, incubation times and temperature for best results in their own IHC assay workflow (automated and manual).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD42b antibody [SP219] - Low endotoxin, Azide free (ab246709)

CD42b Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD42b antibody


Immunohistochemical analysis of paraffin embedded human spleen tissue labeling CD42b with ab183345 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183345).



Flow Cytometry (Intracellular) - Anti-CD42b antibody [SP219] - Low endotoxin, Azide free (ab246709)

CD42b Flow Cytometry (Intracellular) staining using rabbit Anti-CD42b antibody


Intracellular Flow Cytometry analysis of HEL (human bone marrow erythroleukemia) labeling CD42b with purified ab183345 at 1/20 dilution (9.7µg/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilized with 90% methanol. Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as a secondary antibody. Isotype control - Rabbit monoclonal IgG (ab172730) (Black). Unlabeled control - Unlabeled cells (blue).
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183345).



Flow Cytometry (Intracellular) - Anti-CD42b antibody [SP219] - Low endotoxin, Azide free (ab246709)

CD42b Flow Cytometry (Intracellular) staining using rabbit Anti-CD42b antibody


Intracellular flow cytometric analysis of rabbit anti-CD42b (SP219) antibody using ab183345 (diluted 1/400) in Jurkat cells (green) compared to negative control of rabbit IgG (blue).
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183345).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CD42b antibody [SP219] - Low endotoxin, Azide free (ab246709)

CD42b Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CD42b antibody


Immunohistochemical analysis of paraffin embedded human bone marrow tissue labeling CD42b with ab183345 at 1/100 dilution.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183345).



Immunohistochemistry (Frozen sections) - Anti-CD42b antibody [SP219] - Low endotoxin, Azide free (ab246709)

CD42b Immunohistochemistry (Frozen sections) staining using rabbit Anti-CD42b antibody

Immunohistochemistry (Frozen) analysis of mouse spleen tissue section labeling CD42b with purified ab183345 at 1/100 (1.9 µg/ml). Sections were fixed in 4% paraformaldehyde and permeabilized with 0.2% Triton X-100. Antigen retrieval was heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20). Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) was used as the secondary antibody at 1/1000 (2 µg/ml) dilution. DAPI was used as nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183345).



Flow Cytometry (Intracellular) - Anti-CD42b antibody [SP219] - Low endotoxin, Azide free (ab246709)

CD42b Flow Cytometry (Intracellular) staining using rabbit Anti-CD42b antibody

Intracellular Flow Cytometry analysis of mouse splenocytes cell labeling CD42b with purified ab183345 at 1/20 dilution (9.7µg/ml) (red). Cells were fixed with 4% paraformaldehyde and permeabilised with 90% methanol. Goat anti rabbit IgG (Alexa Fluor® 488, ab150081) at 1/2000 dilution was used as a secondary antibody. Isotype control -Rabbit monoclonal IgG (ab172730) (Black). Unlableled control -Unlabeled cells (blue).
This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA and sodium azide (ab183345).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.