abcam

ab247661

Anti-Bcl10 antibody [EPR3174] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR3174
Purification technique	Affinity purification
Concentration	1.019 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Not recommended

Species	Human
Dilution info	-
Notes	-

IP

Not recommended

Species	Human
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Target data

See full target information BCL10

Function	Plays a key role in both adaptive and innate immune signaling by bridging CARD domain-containing proteins to immune activation (PubMed:10187770, PubMed:10364242, PubMed:10400625, PubMed:24074955, PubMed:25365219).

Acts by channeling adaptive and innate immune signaling downstream of CARD domain-containing proteins CARD9, CARD11 and CARD14 to activate NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed:24074955). Recruited by activated CARD domain-containing proteins: homooligomerized CARD domain-containing proteins form a nucleating helical template that recruits BCL10 via CARD-CARD interaction, thereby promoting polymerization of BCL10, subsequent recruitment of MALT1 and formation of a CBM complex (PubMed:24074955). This leads to activation of NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed:18287044, PubMed:24074955, PubMed:27777308). Activated by CARD9 downstream of C-type lectin receptors; CARD9-mediated signals are essential for antifungal immunity (PubMed:26488816). Activated by CARD11 downstream of T-cell receptor (TCR) and B-cell receptor (BCR) (PubMed:18264101, PubMed:18287044, PubMed:24074955, PubMed:27777308). Promotes apoptosis, pro-caspase-9 maturation and activation of NF-kappa-B via NIK and IKK (PubMed:10187815).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab247661 is the carrier-free version of ab108412.

Species reactivity

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	Bcl10 also known as B-cell lymphoma 10 is a protein involved in several cellular processes. It has a molecular mass of approximately 26 kDa and shows expression in various lymphoid tissues. Bcl10 plays a significant role in signaling pathways by transmitting signals from the cell surface to the nucleus impacting cell growth and immune responses. Researchers often study Bcl10 using immunohistochemistry (IHC) techniques referred to as Bcl10 IHC to analyze its expression and distribution within tissues.
Biological function summary	Bcl10 activates downstream signaling pathways that are critical for immune cell functioning and survival. It forms part of the CARD-BCL10-MALT1 (CBM) complex which is important for the activation of the NF-kB pathway. This complex formation is essential for lymphocyte activation and proliferation. Bcl10 interactions within this complex influence cellular responses to various stimuli helping organisms to mount appropriate immune responses.
Pathways	Bcl10 serves as an integral component in the NF-kB and TCR signaling pathways. It functions alongside proteins such as MALT1 and CARD9 in these pathways. The proper functioning of Bcl10 in these pathways ensures accurate transcriptional regulation necessary for immune responses. Misregulation in these pathways can lead to immune disorders and aberrant cell behavior linking Bcl10 to further biological implications.
Associated diseases and disorders	Bcl10 shows associations with mucosa-associated lymphoid tissue (MALT) lymphoma and other inflammatory diseases. Aberrant Bcl10 expression or mutations can lead to unchecked activation of NF-kB contributing to the progression of these conditions. In MALT lymphoma Bcl10 often cooperates with MALT1 both contributing to pathological cell survival and proliferation highlighting the importance of Bcl10 in understanding these diseases.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


2 product images



Flow Cytometry (Intracellular) - Anti-Bcl10 antibody [EPR3174] - BSA and Azide free (ab247661)

Bcl10 Flow Cytometry (Intracellular) staining using rabbit Anti-Bcl10 antibody

This data was developed using ab108412, the same antibody clone in a different buffer formulation.
Intracellular flow cytometric analysis of Bcl10 in permeabilized Raji cells, using ab108412 at a 1/500 dilution (red) or a Rabbit IgG (negative) (green).



Western blot - Anti-Bcl10 antibody [EPR3174] - BSA and Azide free (ab247661)

Bcl10 Western blot staining of Daudi cell lysate using rabbit Anti-Bcl10 antibody

This data was developed using ab108412, the same antibody clone in a different buffer formulation.
All lanes:
Western blot - Anti-Bcl10 antibody [EPR3174] - BSA and Azide free (ab247661) at 1/1000 dilution

All lanes:
Daudi cell lysate at 10 µg

Predicted band size: 26 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.