abcam

ab248335

Anti-CtBP1 antibody [EPR6800] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR6800
Purification technique	Affinity purification Protein A
Dissociation constant	1.5 x 10^{-11} M
Concentration	1.061 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	-

Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information CTBP1

Function	Corepressor targeting diverse transcription regulators such as GLIS2 or BCL6. Has dehydrogenase activity. Involved in controlling the equilibrium between tubular and stacked structures in the Golgi complex. Functions in brown adipose tissue (BAT) differentiation.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab248335 is the carrier-free version of ab129181.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


5 product images



Flow Cytometry (Intracellular) - Anti-CtBP1 antibody [EPR6800] - BSA and Azide free (ab248335)

CtBP1 Flow Cytometry (Intracellular) staining using rabbit Anti-CtBP1 antibody

This data was developed using ab129181, the same antibody clone in a different buffer formulation.
Intracellular Flow Cytometry analysis of Raji (Human Burkitt's lymphoma B lymphocyte) cells labeling CtBP1 (red) with ab129181 at a 1/2000 dilution. Cells were fixed with 4% paraformaldehyde and permeabilized with 90% methanol. A goat anti-rabbit IgG (Alexa Fluor® 488) (ab150077) was used as the secondary antibody at a 1/2000 dilution. Black - Rabbit monoclonal IgG (Black) (ab172730). Blue (unlabeled control) - Cell without incubation with primary antibody and secondary antibody (Blue).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CtBP1 antibody [EPR6800] - BSA and Azide free (ab248335)

CtBP1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CtBP1 antibody

This data was developed using ab129181, the same antibody clone in a different buffer formulation.
ab129181, at 1/100 dilution, staining CtBP1 in formalin-fixed, paraffin-embedded human colon tissue by immunohistochemistry.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-CtBP1 antibody [EPR6800] - BSA and Azide free (ab248335)

CtBP1 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-CtBP1 antibody

This data was developed using ab129181, the same antibody clone in a different buffer formulation.
Immunocytochemistry/Immunofluorescence analysis of HeLa cells labelling CtBP1 with ab129181 at 1/500. Cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% Triton X-100. ab150077, an Alexa Fluor® 488-conjugated goat anti-rabbit IgG (1/1000) was used as the secondary antibody.
Control: PBS only.
Nuclear counter stain: DAPI.



Western blot - Anti-CtBP1 antibody [EPR6800] - BSA and Azide free (ab248335)

CtBP1 Western blot staining using rabbit Anti-CtBP1 antibody

This data was developed using ab129181, the same antibody clone in a different buffer formulation.
All lanes:
Western blot - Anti-CtBP1 antibody [EPR6800] (ab129181) at 1/1000 dilution

Lane 1:
SH-SY5Y cell lysate at 10 µg

Lane 2:
Raji cell lysate at 10 µg

Lane 3:
A549 cell lysate at 10 µg


Lane 4:
HeLa cell lysate at 10 µg

Secondary

All lanes:
HRP-conjugated goat anti-rabbit IgG

Predicted band size: 47 kDa

Observed band size: 48 kDa



OI-RD Scanning - Anti-CtBP1 antibody [EPR6800] - BSA and Azide free (ab248335)

We have systematically measured KD (the equilibrium dissociation constant between the antibody and its antigen), of more than 840 recombinant antibodies to assess not only their individual KD values but also to see the average affinity of antibody. Based on the comparison with published literature values for mouse monoclonal antibodies, Recombinant antibodies appear to be on average 1-2 order of magnitude higher affinity.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.