abcam

ab248716

Anti-MCM7/PRL antibody [EPR1973Y] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR1973Y
Purification technique	Affinity purification Protein A
Concentration	1.009 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Not recommended

Species	Human
Dilution info	-

Notes	-

WB

Tested

Species	Human
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human
Dilution info	-
Notes	-

ICC/IF

Not recommended

Species	Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Target data

See full target information MCM7

Function	Acts as a component of the MCM2-7 complex (MCM complex) which is the replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. Core component of CDC45-MCM-GINS (CMG) helicase, the molecular machine that unwinds template DNA during replication, and around which the replisome is built (PubMed:25661590, PubMed:32453425, PubMed:34694004, PubMed:34700328, PubMed:35585232, PubMed:9305914). The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity (PubMed:32453425). Required for S-phase checkpoint activation upon UV-induced damage.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab248716 is the carrier-free version of ab134194.

Species reactivity

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes,

oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The MCM7 (Minichromosome Maintenance Complex Component 7) also known as PRL serves an important mechanical role in DNA replication. It is part of the MCM2-7 complex a helicase essential for the initiation and elongation phases of DNA replication. MCM7 has a molecular mass of approximately 81 kDa and is found expressed in the nucleus of proliferating cells. It is essential for unwinding the DNA helix preparing it for replication.
Biological function summary	MCM7 is involved in maintaining genomic stability and ensuring accurate DNA replication. As part of the larger MCM2-7 hexameric complex it collaborates with other members to form the core component of the pre-replicative complex (pre-RC). By loading onto replication origins MCM7 helps license the DNA for duplication. This function contributes to cell cycle progression particularly the S phase where DNA synthesis occurs.
Pathways	MCM7 plays an important role in DNA replication-related signaling pathways. It functions within the cell cycle and DNA replication pathways interacting closely with other essential proteins like CDC45 and GINS complex. These partnerships facilitate the transition from the G1 to the S phase triggering the replication machinery's activation and ensuring cell division processes continue smoothly.
Associated diseases and disorders	MCM7 has been associated with various cancers including breast and prostate cancer. Aberrant expression or dysregulation of MCM7 contributes to uncontrolled cell proliferation linking it to tumorigenesis. Also MCM7 interacts with proteins such as cyclin D1 in certain oncogenic pathways emphasizing its role in the progression of cancerous conditions. Understanding these interactions offers insights into potential therapeutic targets for cancer treatment.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---


2 product images



Flow Cytometry (Intracellular) - Anti-MCM7/PRL antibody [EPR1973Y] - BSA and Azide free (ab248716)

MCM7/PRL Flow Cytometry (Intracellular) staining using rabbit Anti-MCM7/PRL antibody

This data was developed using ab134194, the same antibody clone in a different buffer formulation. Intracellular Flow Cytometry analysis of HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling MCM7/PRL with purified ab134194 at 1/600 dilution (1µg/mL) (Red). Cells were fixed with 4% Paraformaldehyde and permeabilised with 90% Methanol. A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) secondary antibody was used at 1/2000. Isotype control - Rabbit monoclonal IgG (Black). Unlabeled control - Cells without incubation with primary antibody and secondary antibody (Blue).



Western blot - Anti-MCM7/PRL antibody [EPR1973Y] - BSA and Azide free (ab248716)

MCM7/PRL Western blot staining using rabbit Anti-MCM7/PRL antibody

This data was developed using ab134194, the same antibody clone in a different buffer formulation.
All lanes:
Western blot - Anti-MCM7/PRL antibody [EPR1973Y] (ab134194) at 1/2000 dilution

Lane 1:
HeLa cell lysate at 10 µg

Lane 2:
Jurkat cell lysate at 10 µg

Secondary

All lanes:
HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 81 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.