abcam

ab248928

Anti-KHSRP antibody [EPR9865] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR9865
Purification technique	Affinity purification Protein A
Concentration	1.015 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	-

Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information KHSRP

Function	Binds to the dendritic targeting element and may play a role in mRNA trafficking (By similarity). Part of a ternary complex that binds to the downstream control sequence (DCS) of the pre-mRNA. Mediates exon inclusion in transcripts that are subject to tissue-specific alternative splicing. May interact with single-stranded DNA from the far-upstream element (FUSE). May activate gene expression. Also involved in degradation of inherently unstable mRNAs that contain AU-rich elements (AREs) in their 3'-UTR, possibly by recruiting degradation machinery to ARE-containing mRNAs.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab248928 is the carrier-free version of ab140648.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

10 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-KHSRP antibody [EPR9865] - BSA and Azide free (ab248928)

KHSRP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-KHSRP antibody

This data was developed using ab140648, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of paraffin embedded Human Cervical carcinoma tissue using ab140648 showing +ve staining.


Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-KHSRP antibody [EPR9865] - BSA and Azide free (ab248928)

KHSRP Flow Cytometry (Intracellular) staining using rabbit Anti-KHSRP antibody

This data was developed using ab140648, the same antibody clone in a different buffer formulation.
Overlay histogram showing Hela cells stained with ab140648 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab140648, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was Alexa Fluor® 488 goat anti-rabbit IgG (H&L) (ab150077) at 1/2000 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1 µg/1x10^6 cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in HeLa cells fixed with 4% paraformaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-KHSRP antibody [EPR9865] - BSA and Azide free (ab248928)

KHSRP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-KHSRP antibody

This data was developed using ab140648, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of paraffin embedded Normal Human Uterus tissue using ab140648 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-KHSRP antibody [EPR9865] - BSA and Azide free (ab248928)

KHSRP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-KHSRP antibody

This data was developed using ab140648, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of paraffin embedded Human Endometrial carcinoma tissue using ab140648 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-KHSRP antibody [EPR9865] - BSA and Azide free (ab248928)

KHSRP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-KHSRP antibody

This data was developed using ab140648, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of paraffin embedded Human Ovarian carcinoma tissue using ab140648 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-KHSRP antibody [EPR9865] - BSA and Azide free (ab248928)

KHSRP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-KHSRP antibody

This data was developed using ab140648, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of paraffin embedded Human brain tissue labelling KHSRP with ab140648 at a 1/100 dilution.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-KHSRP antibody [EPR9865] - BSA and Azide free (ab248928)

KHSRP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-KHSRP antibody

This data was developed using ab140648, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of paraffin embedded Human colon tissue labelling KHSRP with ab140648 at a 1/100 dilution.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-KHSRP antibody [EPR9865] - BSA and Azide free (ab248928)

KHSRP Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-KHSRP antibody

This data was developed using ab140648, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of paraffin embedded Normal Human Colon tissue using ab140648 showing +ve staining.

Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Western blot - Anti-KHSRP antibody [EPR9865] - BSA and Azide free (ab248928)

KHSRP Western blot staining using rabbit Anti-KHSRP antibody

This data was developed using ab140648, the same antibody clone in a different buffer formulation.
All lanes:
Western blot - Anti-KHSRP antibody [EPR9865] (ab140648) at 1/1000 dilution

Lane 1:
293T cell lysate at 10 µg

Lane 2:
HeLa cell lysate at 10 µg


Lane 3:
Jurkat cell lysate at 10 µg

Lane 4:
A673 cell lysate at 10 µg

Secondary

All lanes:
HRP conjugated goat anti-rabbit at 1/2000 dilution

Predicted band size: 73 kDa



Western blot - Anti-KHSRP antibody [EPR9865] - BSA and Azide free (ab248928)

KHSRP Western blot staining using rabbit Anti-KHSRP antibody

This data was developed using the same antibody clone in a different buffer formulation containing PBS, BSA, glycerol, and sodium azide (ab140648).

Western blot: Anti-KHSRP antibody [EPR9865] ab140648 staining at 1/1000 dilution, shown in green; Mouse anti GAPDH ab8245 loading control staining at 1/20000 dilution, shown in magenta. A band was observed at 80-90 kDa in Wild-type HeLa cell line with no signal observed at this size in KHSRP knockout HeLa cell line. To generate this image, samples were run on an SDS-PAGE gel then transferred onto a nitrocellulose membrane. Membranes were blocked in 5% Milk in TBS-0.1 % Tween® 20 (TBS-T) before incubation with primary antibodies overnight at 4 °C. Blots were washed four times in TBS-T, incubated with secondary antibodies for 1 h at room temperature, washed again four times then imaged. Secondary antibodies used were Goat anti-Rabbit 800CW and Goat anti-Mouse 680RD at 1/20,000 dilution.
All lanes:
Western blot - Anti-KHSRP antibody [EPR9865] (ab140648) at 1/1000 dilution

Lane 1:
Wild-type HeLa cell lysate at 20 µg

Lane 2:
Western blot - Human KHSRP knockout HeLa cell line (ab274903) at 20 µg

Lane 3:
HepG2 cell lysate at 20 µg

Lane 4:
LNCaP cell lysate at 20 µg

Secondary

All lanes:
Goat anti-Rabbit 800CW & Goat anti-Mouse 680RD at 1/20000 dilution

Performed under reducing conditions.

Predicted band size: 73 kDa

Observed band size: 80-90 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.