abcam

ab250189

Anti-RNA Helicase A antibody [EPR13520(B)] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR13520(B)
Purification technique	Affinity purification
Concentration	1.03 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Human
Dilution info	-

Notes	-

Predicted

Species	Orangutan
Dilution info	-
Notes	-

IP

Not recommended

Species	Human, Orangutan
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Orangutan
Dilution info	-
Notes	-

IHC-P

Not recommended

Species	Human, Orangutan
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Orangutan
Dilution info	-
Notes	-

Target data

See full target information DHX9

Function

Multifunctional ATP-dependent nucleic acid helicase that unwinds DNA and RNA in a 3' to 5' direction and that plays important roles in many processes, such as DNA replication, transcriptional activation, post-transcriptional RNA regulation, mRNA translation and RNA-mediated gene silencing (PubMed:11416126, PubMed:12711669, PubMed:15355351, PubMed:16680162, PubMed:17531811, PubMed:20669935, PubMed:21561811, PubMed:24049074, PubMed:24990949, PubMed:25062910, PubMed:28221134, PubMed:9111062, PubMed:37467750). Requires a 3'-single-stranded tail as entry site for acid nuclei unwinding activities as well as the binding and hydrolyzing of any of the four ribo- or deoxyribo-nucleotide triphosphates (NTPs) (PubMed:1537828). Unwinds numerous nucleic acid substrates such as double-stranded (ds) DNA and RNA, DNA:RNA hybrids, DNA and RNA forks composed of either partially complementary DNA duplexes or DNA:RNA hybrids, respectively, and also DNA and RNA displacement loops (D- and R-loops), triplex-helical DNA (H-DNA) structure and DNA and RNA-based G-quadruplexes (PubMed:20669935, PubMed:21561811, PubMed:24049074). Binds dsDNA, single-stranded DNA (ssDNA), dsRNA, ssRNA and poly(A)-containing RNA (PubMed:10198287, PubMed:9111062). Also binds to circular dsDNA or dsRNA of either linear and/or circular forms and stimulates the relaxation of supercoiled DNAs catalyzed by topoisomerase TOP2A (PubMed:12711669). Plays a role in DNA replication at origins of replication and cell cycle progression (PubMed:24990949). Plays a role as a transcriptional coactivator acting as a bridging factor between polymerase II holoenzyme and transcription factors or cofactors, such as BRCA1, CREBBP, RELA and SMN1 (PubMed:11038348, PubMed:11149922, PubMed:11416126, PubMed:15355351, PubMed:28221134, PubMed:9323138, PubMed:9662397). Binds to the CDKN2A promoter (PubMed:11038348). Plays several roles in post-transcriptional regulation of gene expression (PubMed:28221134, PubMed:28355180). In cooperation with NUP98, promotes pre-mRNA alternative splicing activities of a subset of genes (PubMed:11402034,

PubMed:16680162, PubMed:28221134, PubMed:28355180). As component of a large PER complex, is involved in the negative regulation of 3' transcriptional termination of circadian target genes such as PER1 and NR1D1 and the control of the circadian rhythms (By similarity). Also acts as a nuclear resolvase that is able to bind and neutralize harmful massive secondary double-stranded RNA structures formed by inverted-repeat Alu retrotransposon elements that are inserted and transcribed as parts of genes during the process of gene transposition (PubMed:28355180). Involved in the positive regulation of nuclear export of constitutive transport element (CTE)-containing unspliced mRNA (PubMed:10924507, PubMed:11402034, PubMed:9162007). Component of the coding region determinant (CRD)-mediated complex that promotes cytoplasmic MYC mRNA stability (PubMed:19029303). Plays a role in mRNA translation (PubMed:28355180). Positively regulates translation of selected mRNAs through its binding to post-transcriptional control element (PCE) in the 5'-untranslated region (UTR) (PubMed:16680162). Involved with LARP6 in the translation stimulation of type I collagen mRNAs for CO1A1 and CO1A2 through binding of a specific stem-loop structure in their 5'-UTRs (PubMed:22190748). Stimulates LIN28A-dependent mRNA translation probably by facilitating ribonucleoprotein remodeling during the process of translation (PubMed:21247876). Plays also a role as a small interfering (siRNA)-loading factor involved in the RNA-induced silencing complex (RISC) loading complex (RLC) assembly, and hence functions in the RISC-mediated gene silencing process (PubMed:17531811). Binds preferentially to short double-stranded RNA, such as those produced during rotavirus intestinal infection (PubMed:28636595). This interaction may mediate NLRP9 inflammasome activation and trigger inflammatory response, including IL18 release and pyroptosis (PubMed:28636595). Finally, mediates the attachment of heterogeneous nuclear ribonucleoproteins (hnRNPs) to actin filaments in the nucleus (PubMed:11687588). (Microbial infection) Plays a role in HIV-1 replication and virion infectivity (PubMed:11096080, PubMed:19229320, PubMed:25149208, PubMed:27107641). Enhances HIV-1 transcription by facilitating the binding of RNA polymerase II holoenzyme to the proviral DNA (PubMed:11096080, PubMed:25149208). Binds (via DRBM domain 2) to the HIV-1 TAR RNA and stimulates HIV-1 transcription of transactivation response element (TAR)-containing mRNAs (PubMed:11096080, PubMed:9892698). Involved also in HIV-1 mRNA splicing and transport (PubMed:25149208). Positively regulates HIV-1 gag mRNA translation, through its binding to post-transcriptional control element (PCE) in the 5'-untranslated region (UTR) (PubMed:16680162). Binds (via DRBM domains) to a HIV-1 double-stranded RNA region of the primer binding site (PBS)-segment of the 5'-UTR, and hence stimulates DHX9 incorporation into virions and virion infectivity (PubMed:27107641). Also plays a role as a cytosolic viral MyD88-dependent DNA and RNA sensors in plasmacytoid dendritic cells (pDCs), and hence induce antiviral innate immune responses (PubMed:20696886, PubMed:21957149). Binds (via the OB-fold region) to viral single-stranded DNA unmethylated C-phosphate-G (CpG) oligonucleotide (PubMed:20696886).

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C

Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab250189 is the carrier-free version of ab180189.

Species reactivity

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	RNA Helicase A also known as DHX9 is a protein essential for various cellular processes. This helicase protein belongs to the DEXH-box family and facilitates the unwinding of RNA structures which is critical for RNA metabolism. The molecular weight of RNA Helicase A is approximately 141 kDa. It is expressed in a variety of human tissues including the nucleus cytoplasm and sometimes nucleoli. The protein’s helicase activity is central to its mechanical role in cellular functioning.

Biological function summary	RNA Helicase A is involved in numerous processes such as transcription RNA processing and the assembly of ribonucleoprotein complexes. The protein plays an important role in facilitating RNA polymerase II transcription and also has functions in RNA splicing through its association with spliceosomal components. RNA Helicase A does not function in isolation; instead it forms part of multi-protein complexes that are necessary for its diverse cellular roles. It interacts with various proteins to achieve its functions impacting both RNA and DNA substrates.
Pathways	RNA Helicase A influences the regulation of gene expression and is a significant player in the innate immune response. It participates in the Toll-like receptor (TLR) signaling pathway where it helps detect viral RNA and triggers immune responses. It also interacts with proteins like TLR3 and TRIF which are involved in the signaling cascade essential for producing type I interferons and pro-inflammatory cytokines.
Associated diseases and disorders	RNA Helicase A has associations with cancer and viral infections. Abnormal regulation of RNA Helicase A has been linked with tumorigenesis where its altered expression contributes to oncogenic processes. It also plays a part in the host response to viral infections where the protein aids in recognizing viral nucleic acids. RNA Helicase A interacts with viral proteins thereby influencing viral replication and host antiviral responses. These connections highlight the protein's relevance in both cancer biology and infectious disease pathogenesis.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


3 product images



Flow Cytometry (Intracellular) - Anti-RNA Helicase A antibody [EPR13520(B)] - BSA and Azide free (ab250189)

RNA Helicase A Flow Cytometry (Intracellular) staining using rabbit Anti-RNA Helicase A antibody


This data was developed using ab180189, the same antibody clone in a different buffer formulation.
Intracellular flow cytometrical analysis of permeabilized Jurkat cells labeling RNA Helicase A with ab180189 at 1/10 (red) or a rabbit IgG negative (green).



Immunocytochemistry/ Immunofluorescence - Anti-RNA Helicase A antibody [EPR13520(B)] - BSA and Azide free (ab250189)

RNA Helicase A Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-RNA Helicase A antibody

This data was developed using ab180189, the same antibody clone in a different buffer formulation.
Immunofluorescent analysis of 293T cells labeling RNA Helicase A with ab180189 at 1/100.



Western blot - Anti-RNA Helicase A antibody [EPR13520(B)] - BSA and Azide free (ab250189)

RNA Helicase A Western blot staining using rabbit Anti-RNA Helicase A antibody

This data was developed using ab180189, the same antibody clone in a different buffer formulation.
All lanes:
Western blot - Anti-RNA Helicase A antibody [EPR13520(B)] - N-terminal (ab180189) at 1/1000 dilution

Lane 1:
BxPC-3 cell lysate at 10 µg

Lane 2:
Jurkat cell lysate at 10 µg

Lane 3:
HeLa cell lysate at 10 µg

Lane 4:
293T (Human embryonic kidney epithelial cell) cell lysate at 10 µg

Secondary

All lanes:
HRP labeled goat anti-rabbit at 1/2000 dilution

Predicted band size: 141 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.