abcam

ab250318

Anti-UAP56 antibody [EPR13144-62] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR13144-62
Purification technique	Affinity purification
Concentration	1.063 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IHC-P

Tested

Species	Human
Dilution info	-

Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information DDX39B

Function

Involved in nuclear export of spliced and unspliced mRNA (PubMed:15833825, PubMed:15998806, PubMed:17190602). Component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA (PubMed:15833825, PubMed:15998806, PubMed:17190602). The TREX complex is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm via the TAP/NXF1 pathway (PubMed:15833825, PubMed:15998806, PubMed:17190602). The THOC1-THOC2-THOC3 core complex alone is sufficient to promote ATPase activity of DDX39B; in the complex THOC2 is the only component that directly interacts with DDX39B (PubMed:33191911). Associates with SARNP/CIP29, which facilitates RNA binding of DDX39B and likely plays a role in mRNA export (PubMed:37578863). May undergo several rounds of ATP hydrolysis during assembly of TREX to drive subsequent loading of components such as ALYREF/THOC4 and CHTOP onto mRNA. Also associates with pre-mRNA independent of ALYREF/THOC4. Involved in the nuclear export of intronless mRNA; the ATP-bound form is proposed to recruit export adapter ALYREF/THOC4 to intronless mRNA; its ATPase activity is cooperatively stimulated by RNA and ALYREF/THOC4 and ATP hydrolysis is thought to trigger the dissociation from RNA to allow the association of ALYREF/THOC4 and the NXF1-NXT1 heterodimer. Involved in transcription elongation and genome stability. Splice factor that is required for the first ATP-dependent step in spliceosome assembly and for the interaction of U2 snRNP with the branchpoint. Has both RNA-stimulated ATP binding/hydrolysis activity and ATP-dependent RNA unwinding activity. Even with the stimulation of RNA, the ATPase activity is weak. Can only hydrolyze ATP but not other NTPs. The RNA stimulation of ATPase activity does not have a strong preference for the sequence and length of the RNA. However, ssRNA stimulates the ATPase activity much more strongly than dsRNA. Can unwind 5' or 3' overhangs or blunt end RNA duplexes in vitro. The ATPase and

helicase activities are not influenced by U2AF2; the effect of ALYREF/THOC4 is reported conflictingly with [PubMed:23299939] reporting a stimulatory effect. (Microbial infection) The TREX complex is essential for the export of Kaposi's sarcoma-associated herpesvirus (KSHV) intronless mRNAs and infectious virus production.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab250318 is the carrier-free version of ab181061.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready

Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility

This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	UAP56 also known as DDX39B or SUB2 is an ATP-dependent RNA helicase with a molecular mass of approximately 46 kDa. It plays a significant role in RNA processing by unwinding RNA duplexes. UAP56 is found throughout the nucleus and functions in various cell types with high expression levels in actively dividing cells. It binds RNA substrates using ATP hydrolysis facilitating movement along RNA strands and this action is vital for splicing and mRNA export from the nucleus to the cytoplasm.
Biological function summary	This protein contributes to the assembly of the spliceosome complex which is essential for pre-mRNA splicing. By interacting with other splicing factors UAP56 ensures accurate removal of introns and the joining of exons in the mRNA. Its function is also necessary for the proper export of mRNA through the nuclear pore complex. Through these activities UAP56 supports gene expression by ensuring that only mature mRNA is available for translation.
Pathways	UAP56 operates in the mRNA processing and transport pathways. In the mRNA export pathway it collaborates with other proteins like Aly/REF and the TREX complex facilitating the translocation of mRNA from the nucleus to the cytoplasm. In the splicing pathway UAP56 interacts with small nuclear ribonucleoproteins (snRNPs) to ensure correct splicing. These interactions help maintain the efficiency and fidelity of gene expression.
Associated diseases and disorders	Defects in UAP56 or its interactions may contribute to cancer and neurodegenerative diseases. In cancer abnormal mRNA export and splicing can lead to the expression of oncogenes or the loss of tumor suppressors implicating UAP56 in tumor progression. In neurodegenerative disorders UAP56 dysfunction might result in improper mRNA processing leading to the accumulation of misfolded proteins. Its role is related to proteins such as TDP-43 which is involved in neurodegenerative pathologies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.


4 product images



Flow Cytometry (Intracellular) - Anti-UAP56 antibody [EPR13144-62] - BSA and Azide free (ab250318)

UAP56 Flow Cytometry (Intracellular) staining using rabbit Anti-UAP56 antibody

This data was developed using ab181061, the same antibody clone in a different buffer formulation.
Intracellular Flow Cytometry analysis of Jurkat cells labeling UAP56 using ab181061 at a 1/10 dilution (pink). Goat anti rabbit IgG (FITC) used as the secondary at a 1/150 dilution. Isotype control Rabbit monoclonal IgG (green). Cells were fixed in 2% paraformaldehyde.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-UAP56 antibody [EPR13144-62] - BSA and Azide free (ab250318)

UAP56 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-UAP56 antibody

This data was developed using ab181061, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of paraffin embedded Human infiltrating duct breast carcinoma tissue sections labeling UAP56 using ab181061 at a 1/250 dilution. A ready to use HRP Polymer for Rabbit IgG was used as the secondary. Hematoxylin counterstain.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunocytochemistry/ Immunofluorescence - Anti-UAP56 antibody [EPR13144-62] - BSA and Azide free (ab250318)

UAP56 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-UAP56 antibody

This data was developed using ab181061, the same antibody clone in a different buffer formulation.
Immunofluorescent analysis of acetone fixed HeLa cells labeling UAP56 using ab181061 at a 1/250 dilution. A Goat anti rabbit IgG (Alexa Fluor®555) was used as the secondary at a 1/200 dilution.



Western blot - Anti-UAP56 antibody [EPR13144-62] - BSA and Azide free (ab250318)

This data was developed using ab181061, the same antibody clone in a different buffer formulation.
Blocking and dilution buffer: 5% NFDM/TBST.

All lanes:
Western blot - Anti-UAP56 antibody [EPR13144-62] (ab181061) at 1/20000 dilution

Lane 1:
K562 cell lysate at 20 µg

Lane 2:
HeLa cell lysate at 20 µg

Lane 3:
Jurkat cell lysate at 20 µg

Lane 4:
293 cell lysate at 20 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 49 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.