abcam

ab250708

Anti-CAND1 antibody [EPR14241] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR14241
Purification technique	Affinity purification Protein A
Concentration	1.045 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Human
Dilution info	-

Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	-
Notes	-

Predicted

Species	Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information CAND1

Function	Key assembly factor of SCF (SKP1-CUL1-F-box protein) E3 ubiquitin ligase complexes that promotes the exchange of the substrate-recognition F-box subunit in SCF complexes, thereby playing a key role in the cellular repertoire of SCF complexes. Acts as a F-box protein exchange factor. The exchange activity of CAND1 is coupled with cycles of neddylation conjugation: in the deneddylated state, cullin-binding CAND1 binds CUL1-RBX1, increasing dissociation of the SCF complex and promoting exchange of the F-box protein. Probably plays a similar role in other cullin-RING E3 ubiquitin ligase complexes.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab250708 is the carrier-free version of ab183748.

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CAND1 also known as cullin-associated and neddylation-dissociated 1 is a scaffolding protein with a molecular mass of approximately 130 kDa. It is found in the cytoplasm and nucleus of eukaryotic cells. CAND1 plays a critical role in regulating the activity of cullin-RING ubiquitin ligases (CRLs) which are important for protein degradation processes. By interacting with neddylated cullin proteins CAND1 modulates the assembly and disassembly of these CRL complexes therefore influencing their activity.
Biological function summary	CAND1 mediates the dynamics of CRL complexes acting as a regulator of substrate specificity and turnover. CAND1 does not function alone; it is integral to a complex that includes cullin proteins and various CRL adaptors. This interaction ensures the right substrates are targeted for ubiquitination leading to proteasomal degradation. The ability of CAND1 to bind to cullin substrates is influenced by cullin neddylation and deneddylation cycles making it essential for the proper function of the ubiquitin-proteasome system.
Pathways	Studies have shown that CAND1 is important in the ubiquitin-proteasome pathway along with the neddylation pathway. Within these pathways it primarily interacts with proteins such as cullins 1 3 and 4 which are central to CRLs. By regulating these interactions CAND1 ensures that protein turnover is controlled which is important for processes like cell cycle progression and signal transduction.

Associated diseases and disorders	Research suggests a connection between CAND1 dysregulation and certain cancers as well as neurodegenerative disorders. In cancer aberrant regulation of CAND1 can lead to uncontrolled cell proliferation due to faulty protein degradation. In neurodegenerative diseases ineffective CAND1-mediated pathways may contribute to the accumulation of damaged proteins. Understanding CAND1's relationship with key proteins in these conditions such as p53 in cancer highlights its potential as a therapeutic target.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

4 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-CAND1 antibody [EPR14241] - BSA and Azide free (ab250708)

CAND1 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-CAND1 antibody

This data was developed using ab183748, the same antibody clone in a different buffer formulation.
Immunohistochemical analysis of Paraffin-embedded Human pancreas tissue labeling CAND1 with ab183748 at 1/100 dilution. HRP conjugated Rabbit IgG was used as a secondary antibody. Counter stained with Hematoxylin.

Perform heat mediated antigen retrieval with EDTA buffer pH 9 before commencing with IHC staining protocol.




Flow Cytometry (Intracellular) - Anti-CAND1 antibody [EPR14241] - BSA and Azide free (ab250708)

CAND1 Flow Cytometry (Intracellular) staining using rabbit Anti-CAND1 antibody

This data was developed using ab183748, the same antibody clone in a different buffer formulation.
Intracellular flow cytometric analysis of 293T cells (fixed in 2% paraformaldehyde) labeling CAND1 with ab183748 at 1/220 dilution (red).

Isotype control Rabbit monoclonal IgG (green). Secondary antibody Goat anti rabbit IgG (FITC) at a 1/150 dilution.




Immunoprecipitation - Anti-CAND1 antibody [EPR14241] - BSA and Azide free (ab250708)

This data was developed using ab183748, the same antibody clone in a different buffer formulation.
Western blot analysis of CAND1 in immunoprecipitation pellets from Jurkat lysate. ab183748 used at 1/50 dilution. Secondary antibody Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000.

All lanes:
Immunoprecipitation - Anti-CAND1 antibody [EPR14241] - N-terminal (ab183748)

Predicted band size: 136 kDa



Western blot - Anti-CAND1 antibody [EPR14241] - BSA and Azide free (ab250708)

This data was developed using ab183748, the same antibody clone in a different buffer formulation.
All lanes:
Western blot - Anti-CAND1 antibody [EPR14241] - N-terminal (ab183748) at 1/20000 dilution

Lane 1:
293T whole cell lysate at 20 µg

Lane 2:
Jurkat whole cell lysate at 20 µg

Lane 3:
Saos-2 whole cell lysate at 20 µg

Secondary

All lanes:

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 136 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.