abcam

ab250997

Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] - BSA and Azide free

Recombinant | RabMAb

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Constituents: PBS
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Carrier free	Yes
Clone number	EPR18407
Concentration	0.972 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Mouse, Human
Dilution info	-
Notes	-

Expected

Species	Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

WB

Tested

Species	Recombinant full length protein - Human, Mouse, Rat, Human
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	-
Notes	-

Expected

Species	Human, Rat
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Recombinant full length protein - Human
Dilution info	-
Notes	-

Target data

See full target information ENO1

Function	Enolase that catalyzes the conversion of 2-phosphoglycerate to phosphoenolpyruvate in glycolysis and the reverse reaction in gluconeogenesis (PubMed:1369209, PubMed:29775581). Also involved in various processes such as growth control, hypoxia tolerance and allergic responses (PubMed:10802057, PubMed:12666133, PubMed:2005901, PubMed:29775581). May also function in the intravascular and pericellular fibrinolytic system due to its ability to serve as a receptor and activator of plasminogen on the cell surface of several cell-types such as leukocytes and neurons (PubMed:12666133). Stimulates immunoglobulin production (PubMed:1369209). Isoform MBP-1. Binds to the myc promoter and acts as a transcriptional repressor. May be a tumor suppressor.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	+4°C
Storage information	Do Not Freeze

Notes

ab250997 is the carrier-free version of ab189891.

Patented technology

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Conjugation ready
Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.

Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with 1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.

Compatibility
This product is compatible with the Maxpar® Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar® is a trademark of Fluidigm Canada Inc.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	ENO1 ENO2 and ENO3 also known as alpha gamma and beta enolase are isoenzymes involved in the glycolytic pathway. Each enolase isoenzyme offers specific mechanical functions in this energy-producing process. ENO1 has an approximate mass of 47 kDa and is widely expressed in most tissues. ENO2 similar in size is primarily found in neural and neuroendocrine tissues. ENO3 also roughly 47 kDa predominantly expresses in muscle tissue. These proteins catalyze the reversible conversion of 2-phosphoglycerate to phosphoenolpyruvate playing an important role in glycolysis.
Biological function summary	The enolase family of isoenzymes acts as multifunctional catalysts and has roles beyond glycolysis. ENO1 serves as a plasminogen receptor on cell surfaces assisting in tissue remodeling and immune response. ENO2 often referred to as neuron-specific enolase (NSE) functions importantly in neural development and tumor cell metabolism. ENO3 or muscle-specific enolase contributes to muscle metabolism and energy turnover. While these proteins individually carry out essential tasks they also form a complex that facilitates efficient energy production and cellular regulation.
Pathways	The enolase enzymes integrate in the glycolytic and gluconeogenesis pathways central to energy metabolism. Glycolysis involves other key proteins like hexokinase and phosphofructokinase cooperating for glucose breakdown and energy release. In gluconeogenesis the enolases help generate glucose from non-carbohydrate sources important during fasting or intense exercise. Their activity along with related proteins impacts the efficiency and regulation of energy metabolism pathways.
Associated diseases and disorders	Enolase dysregulation associates with conditions like cancer and neuromuscular diseases. Elevated ENO2 levels often serve as a biomarker for neuroendocrine tumors and a target for cancer therapies. ENO3 mutations link to muscle metabolic disorders including glycogen storage diseases. ENO1 interaction with proteins such as plasminogen can influence cancer cell invasion and metastasis highlighting its relevance in disease progression and potential therapeutic interventions.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

7 product images



Immunocytochemistry/ Immunofluorescence - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] - BSA and Azide free (ab250997)

ENO1 + ENO2 + ENO3 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ENO1 + ENO2 + ENO3 antibody

This data was developed using ab189891, the same antibody clone in a different buffer formulation.

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (Human epithelial cell line from cervix adenocarcinoma) cells labeling ENO1 + ENO2 + ENO3 with ab189891 at 1/500 dilution, followed by Goat Anti-Rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution (green).

Confocal image showing mostly cytoplasmic staining on Hela cells.


The nuclear counterstain is DAPI (blue).

Tubulin is detected with Anti-alpha Tubulin - Loading Control (ab7291) at 1/1000 dilution and Goat Anti-Mouse IgG (AlexaFluor®594) preadsorbed (ab150120) at 1/1000 dilution (red).

The negative controls are as follows:-

-ve control 1: ab189891 at 1/500 dilution followed by ab150120 at 1/1000 dilution.


-ve control 2: ab7291 at 1/1000 dilution followed by ab150077 at 1/1000 dilution.



Flow Cytometry (Intracellular) - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] - BSA and Azide free (ab250997)

ENO1 + ENO2 + ENO3 Flow Cytometry (Intracellular) staining using rabbit Anti-ENO1 + ENO2 + ENO3 antibody

This data was developed using ab189891, the same antibody clone in a different buffer formulation.
Intracellular flow cytometric analysis of 4% paraformaldehyde-fixed NIH/3T3 (Mouse embryonic fibroblast cell line) labeling ENO1 + ENO2 + ENO3 with ab189891 at 1/70 dilution (red) compared with a Rabbit IgG,monoclonal - Isotype control (ab172730) (black) and an unlabelled control (cells without incubation with primary antibody and secondary antibody) (blue). Goat anti Rabbit IgG (Alexa Fluor® 488) at 1/500 dilution was used as the secondary antibody.



Immunocytochemistry/ Immunofluorescence - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] - BSA and Azide free (ab250997)

ENO1 + ENO2 + ENO3 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-ENO1 + ENO2 + ENO3 antibody

This data was developed using ab189891, the same antibody clone in a different buffer formulation.

Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 (Mouse embryonic fibroblast cell line) cells labeling ENO1 + ENO2 + ENO3 with ab189891 at 1/500 dilution, followed by Goat Anti-Rabbit IgG (Alexa Fluor® 488) (ab150077) secondary antibody at 1/1000 dilution (green).

Confocal image showing mostly cytoplasmic staining on NIH/3T3 cells.


The nuclear counterstain is DAPI (blue).

Tubulin is detected with Anti-alpha Tubulin antibody -Loading Control (ab7291) at 1/1000 dilution and Goat Anti-Mouse IgG (AlexaFluor®594) preadsorbed (ab150120) at 1/1000 dilution (red).

The negative controls are as follows:-

-ve control 1: ab189891 at 1/500 dilution followed by ab150120 at 1/1000 dilution.

-ve control 2: ab7291 at 1/1000 dilution followed by ab150077 at 1/1000 dilution.



Western blot - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] - BSA and Azide free (ab250997)

ENO1 + ENO2 + ENO3 Western blot staining using rabbit Anti-ENO1 + ENO2 + ENO3 antibody

This data was developed using ab189891, the same antibody clone in a different buffer formulation.
Blocking and dilution buffer: 5% NFDM/TBST.

Human ENO1 full length recombinant protein (ab89248) contains aa1-434. Human ENO2 full length recombinant protein contains aa1-434 with a His-Tag®. Human ENO3 full length recombinant protein (ab113127) contains aa1-434 with a His-Tag®. Human ENO2 full length recombinant protein was made in-house.

All lanes:
Western blot - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] (ab189891) at 1/1000 dilution

Lane 1:
Human ENO1 full length recombinant protein at 0.02 µg

Lane 2:
Human ENO2 full length recombinant protein at 0.02 µg

Lane 3:
Human ENO3 full length recombinant protein at 0.02 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 47 kDa

Observed band size: 47 kDa

Exposure time: 5s



Western blot - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] - BSA and Azide free (ab250997)

ENO1 + ENO2 + ENO3 Western blot staining using rabbit Anti-ENO1 + ENO2 + ENO3 antibody

This data was developed using ab189891, the same antibody clone in a different buffer formulation.
Blocking and dilution buffer: 5% NFDM/TBST.

Exposure times: Lanes 1-2: 5 seconds; Lane 3: 15 seconds.

All lanes:
Western blot - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] (ab189891) at 1/1000 dilution

Lane 1:
Human fetal heart lysate at 10 µg

Lane 2:
Human fetal kidney lysate at 10 µg


Lane 3:
Human fetal spleen lysate at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG Peroxidase Conjugate, specific to the non-reduced form of IgG at 1/10000 dilution

Predicted band size: 47 kDa

Observed band size: 47 kDa



Western blot - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] - BSA and Azide free (ab250997)

ENO1 + ENO2 + ENO3 Western blot staining using rabbit Anti-ENO1 + ENO2 + ENO3 antibody

This data was developed using ab189891, the same antibody clone in a different buffer formulation.
Blocking and dilution buffer: 5% NFDM/TBST.

All lanes:
Western blot - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] (ab189891) at 1/5000 dilution

Lane 1:
HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate at 20 µg

Lane 2:
Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate at 20 µg

Lane 3:
MCF7 (Human breast adenocarcinoma cell line) whole cell lysate at 20 µg

Lane 4:
A431 (Human epidermoid carcinoma cell line) whole cell lysate at 20 µg


Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 47 kDa

Observed band size: 47 kDa

Exposure time: 3s



Western blot - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] - BSA and Azide free (ab250997)

ENO1 + ENO2 + ENO3 Western blot staining using rabbit Anti-ENO1 + ENO2 + ENO3 antibody

This data was developed using ab189891, the same antibody clone in a different buffer formulation.
Blocking and dilution buffer: 5% NFDM/TBST.

Exposure time: Lanes 1-4: 5 seconds; Lanes 5-7: 1 second.

Lanes 1 - 4:
Western blot - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] (ab189891) at 1/1000 dilution

Lanes 5 - 7:
Western blot - Anti-ENO1 + ENO2 + ENO3 antibody [EPR18407] (ab189891) at 1/5000 dilution

Lane 1:
Mouse brain tissue lysate at 10 µg

Lane 2:
Mouse heart tissue lysate at 10 µg

Lane 3:
Rat brain tissue lysate at 10 µg

Lane 4:
Rat heart tissue lysate at 10 µg

Lane 5:
C6 (Rat glial tumor cell line) whole cell lysate at 10 µg

Lane 6:
RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) whole cell lysate at 10 µg

Lane 7:
NIH/3T3 (Mouse embryonic fibroblast cell line) whole cell lysate at 10 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/10000 dilution

Predicted band size: 47 kDa

Observed band size: 47 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.