abcam

ab254151

Anti-CD35 antibody [7G9]

Recombinant

Key facts

Isotype	IgG2a
Host species	Mouse
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	7G9
Purification technique	Affinity purification Protein A
Concentration	0.944 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Human
Dilution info	1/1000
Notes	-

ICC/IF

Tested

Species	Human
Dilution info	1/100
Notes	-

Target data

See full target information CR1

Function	Membrane immune adherence receptor that plays a critical role in the capture and clearance of complement-opsonized pathogens by erythrocytes and monocytes/macrophages (PubMed:2963069). Mediates the binding by these cells of particles and immune complexes that have activated complement to eliminate them from the circulation (PubMed:2963069). Also acts in the inhibition of spontaneous complement activation by impairing the formation and function of the alternative and classical pathway C3/C5 convertases, and by serving as a cofactor for the cleavage by factor I of C3b to iC3b, C3c and C3d,g, and of C4b to C4c and C4d (PubMed:2972794, PubMed:8175757). Also plays a role in immune regulation by contributing, upon ligand binding, to the generation of regulatory T cells from activated helper T cells (PubMed:25742728). (Microbial infection) Acts as a receptor for Epstein-Barr virus.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Want a custom formulation?
This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

2 product images




Immunocytochemistry/ Immunofluorescence - Anti-CD35 antibody [7G9] (ab254151)

CD35 Immunocytochemistry/ Immunofluorescence staining using mouse Anti-CD35 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized human PBMC cells labelling CD35 / CR1 with ab254151 at 1/100 dilution, followed by ab150113 Goat Anti-mouse IgG H&L (Alexa Fluor® 488) antibody at 1/1000 dilution (Green). Confocal image showing membranous and cytoplasmic staining in human PBMC. ab179513 Anti-beta Tubulin

rabbit monoclonal antibody was used to counterstain tubulin at 1/500 dilution, followed by ab150080 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 594) (Red). The nuclear counterstain was DAPI (Blue).
Negative control 1: ab254151 at a 1/100 dilution followed by ab150080 at a 1/500 diliution.

Negative control 2: ab179513 at a 1/500 dilution followed by ab150113 at a 1/1000 diliution.



Flow Cytometry - Anti-CD35 antibody [7G9] (ab254151)

CD35 Flow Cytometry staining using mouse Anti-CD35 antibody

Flow cytometric analysis of human peripheral blood mononuclear cell (PBMC) cells labelling CD35 / CR1 with ab254151 at 1/1000 dilution (0.1µg) (Right panel) compared with a mouse monoclonal IgG isotype control (Left panel). Goat anti mouse IgG (Alexa Fluor® 488, ab150113) at 1/2000 dilution was used as the secondary antibody.
Cells were stained with mouse IgG (Left) or ab254151 (Right), then stained with anti-CD19 conjugated to Alexa Fluor® 647.

Gated on viable cells.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.