abcam

ab254298

Anti-Eg5 antibody [EPR23276-52]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23276-52
Purification technique	Affinity purification Protein A
Concentration	0.662 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

IP

Tested

Species	Mouse
Dilution info	1/30
Notes	-

Species	Human

Dilution info	1/30
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

WB

Tested

Species	Mouse
Dilution info	1/1000
Notes	-

Species	Human
Dilution info	1/1000
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

ICC/IF

Tested

Species	Mouse
Dilution info	1/50
Notes	-

Species	Human
Dilution info	1/50

Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Mouse
Dilution info	1/600
Notes	-

Species	Human
Dilution info	1/600
Notes	-

Not recommended

Species	Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/4000
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Not recommended

Species	Mouse
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

Species	Rat
Dilution info	-
Notes	Perform heat-mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

IHC-Fr

Not recommended

Species	Rat, Mouse, Human
Dilution info	-
Notes	-

Target data

See full target information KIF11

Function	Motor protein required for establishing a bipolar spindle and thus contributing to chromosome congression during mitosis (PubMed:19001501, PubMed:37728657). Required in non-mitotic cells for transport of secretory proteins from the Golgi complex to the cell surface (PubMed:23857769).

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot

Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	The Eg5 protein also known scientifically as kinesin family member 11 (KIF11) plays an important role in cell mitosis. It functions mechanically as a plus-end directed motor involved in the separation of centrosomes during the early stage of cell division. Eg5 with an approximate mass of 116 kDa is responsible for the proper alignment and segregation of spindle fibers. The protein is expressed in rapidly dividing cells particularly within tissues with high turnover rates such as bone marrow and epithelium.
Biological function summary	The motor function of Eg5 is indispensable for the mitotic spindle organization. It provides the force necessary to push microtubules in opposite directions. Eg5 functions as part of a complex cellular machinery that includes proteins like dynein and other kinesins. By controlling spindle bipolarity it ensures accurate chromosome segregation during cell division preventing errors that could lead to aneuploidy.
Pathways	The action of Eg5 is fundamental in mitotic pathways including the spindle assembly checkpoint pathway. It interacts with proteins such as cyclin-dependent kinases (CDKs) which regulate its activity to ensure proper cell cycle progression. The protein's role is integral in facilitating the dynamics of the metaphase spindle checkpoint guaranteeing cell cycle fidelity.
Associated diseases and disorders	Eg5 is linked to certain types of cancers where there is abnormal cell division. The overexpression or dysregulation of Eg5 can contribute to tumorigenesis. Additionally Eg5 interacts with proteins involved in cancer such as p53 a tumor suppressor protein directly influencing cellular proliferation and survival. Researchers target Eg5 in the development of anticancer therapies to inhibit its activity aiming to obstruct malignant cell proliferation.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended

We do not recommend this combination. It is not covered by our product promise.


We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

---

12 product images




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Eg5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Eg5 antibody

Immunohistochemical analysis of paraffin-embedded human tonsil tissue labeling Eg5 with 254298 at 1/4000 (0.17µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Cytoplasmic staining in human tonsil is observed (PMID:25277178).
The section was incubated with ab254298 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Eg5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Eg5 antibody

Immunohistochemical analysis of paraffin-embedded human breast cancer tissue labeling Eg5 with ab254298 at 1/4000 (0.17µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Scattered cytoplasmic staining in human breast cancer cells is observed (PMID:29181100).
The section was incubated with ab254298 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).


Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunocytochemistry/ Immunofluorescence - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Eg5 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Eg5 antibody


Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa cells labeling Eg5 with ab254298 at 1/50 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 2 µg/ml dilution (Green). Confocal image showing cytoplasmic and spindle (arrow) staining in HeLa cells. ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5 µg/ml dilution (Red). The nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 2 µg/ml dilution.



Flow Cytometry (Intracellular) - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Eg5 Flow Cytometry (Intracellular) staining using rabbit Anti-Eg5 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized HeLa (Human cervix adenocarcinoma epithelial cell) cells labeling Eg5 with ab254298 at 1/600 dilution (0.1µg) (Red) compared with a Rabbit monoclonal IgG (ab172730) / Black isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Eg5 Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Eg5 antibody

Immunohistochemical analysis of paraffin-embedded human testis tissue labeling Eg5 with 254298 at 1/1/4000 (0.17µg/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Cytoplasmic staining in spermatocytes of human testis is observed.
The section was incubated with ab254298 for 30 mins at room temperature.
The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with hematoxylin.
Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Immunoprecipitation - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Eg5 was immunoprecipitated from 0.35 mg HeLa (human cervix adenocarcinoma epithelial cell), whole cell lysate 10 ug with ab254298 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab254298 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: HeLa whole cell lysate 10 ug.

Lane 2: ab254298 IP in HeLa whole cell lysate.

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab254298 in HeLa whole cell lysate.


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes.

Fresh lysates were used in this IP.

All lanes:
Immunoprecipitation - Anti-Eg5 antibody [EPR23276-52] (ab254298)


Predicted band size: 119 kDa



Flow Cytometry (Intracellular) - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Eg5 Flow Cytometry (Intracellular) staining using rabbit Anti-Eg5 antibody

Intracellular flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized NIH/3T3 (Mouse embryonic fibroblast) cells labeling Eg5 with ab254298 at 1/600 dilution (0.1µg) (Red) compared with a Rabbit monoclonal IgG (ab172730) / Black isotype control and an unlabeled control (cells without incubation with primary antibody and secondary antibody) (Blue). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunocytochemistry/ Immunofluorescence - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Eg5 Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Eg5 antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized NIH/3T3 cells labeling Eg5 with ab254298 at 1/50 dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 2 µg/ml dilution (Green). Confocal image showing cytoplasmic and spindle (arrow) staining in NIH/3T3 cells. ab195889 Anti-alpha Tubulin antibody

[DM1A] - Microtubule Marker (Alexa Fluor® 594) was used to counterstain tubulin at 1/200 2.5 µg/ml dilution (Red). The nuclear counterstain was DAPI (Blue).
Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 2 µg/ml dilution.



Immunoprecipitation - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Eg5 was immunoprecipitated from 0.35 mg NIH/3T3 (mouse embryonic fibroblast), whole cell lysate 10 ug with ab254298 at 1/30 dilution (2ug in 0.35mg lysates). Western blot was performed on the immunoprecipitate using ab254298 at 1/1000 dilution. VeriBlot for IP Detection Reagent (HRP)(ab131366) was used at 1/5000 dilution.
Lane 1: NIH/3T3 whole cell lysate 10 ug.

Lane 2: ab254298 IP in NIH/3T3 whole cell lysate.

Lane 3: Rabbit monoclonal IgG (ab172730) instead of ab254298 in NIH/3T3 whole cell lysate.


Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Exposure time: 3 minutes

Fresh lysates were used in this IP.

All lanes:
Immunoprecipitation - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Predicted band size: 119 kDa



Western blot - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
The expression profile observed is consistent with what has been described in the literature (PMID:23857769; 27492783).

Negative control: Human lung (PMID: 27279560).

The minor bands beneath the target band may correspond to degradation.

Exposure time: 48 seconds.

All lanes:
Western blot - Anti-Eg5 antibody [EPR23276-52] (ab254298) at 1/1000 dilution

Lane 1:
Human testis tissue lysate at 20 µg


Lane 2:
Human tonsil tissue lysate at 20 µg

Lane 3:
Human lung tissue lysate at 20 µg

Secondary

All lanes:
Western blot - VeriBlot for IP Detection Reagent (HRP) (ab131366) at 1/100000 dilution

Predicted band size: 119 kDa



Western blot - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Eg5 Western blot staining using rabbit Anti-Eg5 antibody

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
The expression profile observed is consistent with what has been described in the literature (PMID:23857769; 27492783).

Fresh lysates were tested in this WB.

The minor bands beneath the target band in lane 1 and lane 4 may correspond to degradation.

Exposure time: 15 seconds.

All lanes:
Western blot - Anti-Eg5 antibody [EPR23276-52] (ab254298) at 1/1000 dilution

Lane 1:
Jurkat (human T cell leukemia T lymphocyte), whole cell lysate at 20 µg

Lane 2:
NCI-H1975 (human adenocarcinoma lung epithelial cell) at 20 µg


Lane 3:
MCF7 (human breast adenocarcinoma epithelial cell) at 20 µg

Lane 4:
4T1 (mouse mammary gland carcinoma epithelial cell) at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 119 kDa



Western blot - Anti-Eg5 antibody [EPR23276-52] (ab254298)

Blocking and diluting buffer and concentration: 5% NFDM/TBST.
Negative control: Mouse brain (PMID: 17974955).

The minor bands beneath the target band may correspond to degradation.

Exposure time: 48 seconds.

All lanes:
Western blot - Anti-Eg5 antibody [EPR23276-52] (ab254298) at 1/1000 dilution

Lane 1:
Mouse testis tissue lysate at 20 µg

Lane 2:
Mouse brain tissue lysate at 20 µg

Secondary

All lanes:
Western blot - Goat Anti-Rabbit IgG H&L (HRP) (ab97051) at 1/100000 dilution

Predicted band size: 119 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.