abcam

ab254400

Anti-Histone H3.3 (mutated K36M) antibody [EPR23526-89]

Recombinant | RabMAb | 20ul selling size

Key facts

Isotype	IgG
Host species	Rabbit
Storage buffer	pH: 7.2 - 7.4 Preservative: 0.01% Sodium azide Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	EPR23526-89
Purification technique	Affinity purification Protein A
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

ICC/IF

Tested

Species	Transfected cell line
Dilution info	1/1000
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Synthetic peptide, Mouse, Rat
Dilution info	-
Notes	-

IP

Not recommended

Species	Human, Transfected cell line, Synthetic peptide, Mouse, Rat
Dilution info	-
Notes	-

ChIP

Not recommended

Species	Human, Transfected cell line, Synthetic peptide, Mouse, Rat
Dilution info	-
Notes	-

Dot

Tested

Species	Synthetic peptide
Dilution info	1/1000
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Transfected cell line, Mouse, Rat
Dilution info	-
Notes	-

WB

Tested

Species	Transfected cell line
Dilution info	1/1000
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Synthetic peptide, Mouse, Rat
Dilution info	-
Notes	-

IHC-P

Tested

Species	Human
Dilution info	1/1000

Notes	-

Not recommended

Species	Mouse, Rat, Transfected cell line, Synthetic peptide
Dilution info	-
Notes	-

I-ELISA

Tested

Species	Human
Dilution info	0.0065 g/mL
Notes	-

Not recommended

Species	Transfected cell line, Synthetic peptide, Mouse, Rat
Dilution info	-
Notes	-

Flow Cyt (Intra)

Tested

Species	Transfected cell line
Dilution info	1/500
Notes	-

Expected

Species	Human
Dilution info	Use at an assay dependent concentration.
Notes	-

Not recommended

Species	Synthetic peptide, Mouse, Rat
Dilution info	-
Notes	-

Target data

See full target information H3-3A K36M

Function	Variant histone H3 which replaces conventional H3 in a wide range of nucleosomes in active genes. Constitutes the predominant form of histone H3 in non-dividing cells and is incorporated into chromatin independently of DNA synthesis. Deposited at sites of nucleosomal displacement throughout transcribed genes, suggesting that it represents an epigenetic imprint of transcriptionally active chromatin. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Storage

Shipped at conditions	Conditional Ambient
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle

Notes

Patented technology
Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

What are the advantages of a recombinant monoclonal antibody?
This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free batch production

For more information, read more on recombinant antibodies.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

7 product images



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H3.3 (mutated K36M) antibody [EPR23526-89] (ab254400)


Histone H3.3 (mutated K36M) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Histone H3.3 (mutated K36M) antibody

Immunohistochemical analysis of paraffin-embedded Human gaint cell tumor of bone tissue labelling Histone H3 (mutated K36M) with ab254400 at 1/1000 (0.548 ug/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Counterstained with Hematoxylin.

Negative control: No staining in human gaint cell tumor of bone (PMID: 29757500).

Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).


Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0)



Flow Cytometry (Intracellular) - Anti-Histone H3.3 (mutated K36M) antibody [EPR23526-89] (ab254400)

Histone H3.3 (mutated K36M) Flow Cytometry (Intracellular) staining using rabbit Anti-Histone H3.3 (mutated K36M) antibody

Flow cytometric analysis of 4% paraformaldehyde fixed 90% methanol permeabilized 293T transfected with myc tagged Histone H3.3 construct (Left) or myc tagged Histone H3.3 H3K36M construct (Right) cells labelling Histone H3 (mutated K36M) with ab254400 at 1/500 dilution (0.1ug). A Goat anti rabbit IgG (Alexa Fluor® 488, ab150077) at 1/2000 dilution was used as the secondary antibody.




Immunocytochemistry/ Immunofluorescence - Anti-Histone H3.3 (mutated K36M) antibody [EPR23526-89] (ab254400)

Histone H3.3 (mutated K36M) Immunocytochemistry/ Immunofluorescence staining using rabbit Anti-Histone H3.3 (mutated K36M) antibody

Immunofluorescent analysis of 4% Paraformaldehyde-fixed, 0.1% TritonX-100 permeabilized 293T cells labelling Histone H3 (mutated K36M) with ab254400 at 1/1000 (0.548 ug/ml) dilution, followed by ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) antibody at 1/1000 (2 ug/ml) dilution (Green). Myc-Tag Mouse mAb (Alexa Fluor® 647) was used to counterstain tubulin at 1/200 (2.5 ug/ml) dilution (Red). The Nuclear counterstain was DAPI (Blue).
Confocal image showing nuclear staining in 293T cells transfected with myc-tagged H3K36M expression vector is observed.

Secondary antibody only control: Secondary antibody is ab150077 Goat Anti-Rabbit IgG H&L (Alexa Fluor® 488) at 1/1000 (2 ug/ml) dilution.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Histone H3.3 (mutated K36M) antibody [EPR23526-89] (ab254400)

Histone H3.3 (mutated K36M) Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) staining using rabbit Anti-Histone H3.3 (mutated K36M) antibody

Immunohistochemical analysis of paraffin-embedded Human chondroblastoma tissue labelling Histone H3 (mutated K36M) with ab254400 at 1/1000 (0.548 ug/ml) followed by a ready to use Goat Anti-Rabbit IgG H&L (HRP). Counterstained with Hematoxylin.
Nuclear staining in human chondroblastoma (PMID: 29757500).


Secondary antibody only control: Secondary antibody is a ready to use Goat Anti-Rabbit IgG H&L (HRP).

Heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0)



Indirect ELISA - Anti-Histone H3.3 (mutated K36M) antibody [EPR23526-89] (ab254400)

ELISA using ab254400 at varying antibody concentrations and antigen concentration at 100 ng/ml. An Alkaline Phosphatase-conjugated AffiniPure Goat Anti-Rabbit IgG (H+L) (1/2500) was used as the secondary antibody.



Western blot - Anti-Histone H3.3 (mutated K36M) antibody [EPR23526-89] (ab254400)

Blocking and diluting buffer and concentration: 5% NFDM/TBST
Exposure time: 92 seconds

All lanes:
Western blot - Anti-Histone H3.3 (mutated K36M) antibody [EPR23526-89] (ab254400) at 1/1000 dilution

Lane 1:
HEK-293 transfected with Histone H3.3 H3K36M expression vector containing a myc-His-tag®, whole cell lysate at 10 µg


Lane 2:
HEK-293 transfected with Histone H3.3 (WT) expression vector containing a myc-His-tag®, whole cell lysate at 10 µg

Secondary

All lanes:
Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/20000 dilution

Predicted band size: 15 kDa

Observed band size: 20 kDa



Dot Blot - Anti-Histone H3.3 (mutated K36M) antibody [EPR23526-89] (ab254400)

Dot blot analysis of Histone H3.3 (mutated K36M) using ab254400 at 1/1000 (0.548 ug/ml) followed by a Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ab97051) at 1/100,000 dilution.
Lane 1: Histone H3.3 (mutated K36M)peptide (aa28-40)

Lane 2: Histone H3.3 (mutated K36M) peptide (aa29-42)

Lane 3: Histone H3.3 (WT) peptide (aa28-42)

Exposure time: 3 minutes

Blocking and diluting buffer and concentration: 5% NFDM/TBST

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.