abcam

ab256263

Alexa Fluor® 488 Anti-CD45RA antibody [MRC OX-33]

Anti-CD45RA antibody [MRC OX-33] conjugated to Alexa Fluor® 488 - suitable for flow cytometry, and reacts with rat samples. Clone [MRC OX-33] is a mouse monoclonal antibody.

Design your panel with our interactive flow cytometry panel builder. Click here.

Key facts

Isotype	IgG1
Host species	Mouse
Conjugation	Alexa Fluor® 488
Excitation/Emission	Ex: 495nm, Em: 519nm
Storage buffer	Preservative: 0.02% Sodium azide Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA
Form	Liquid
Clonality	Monoclonal
Immunogen	The exact immunogen used to generate this antibody is proprietary information.
Clone number	MRC OX-33
Purification technique	Affinity purification
Specificity	This antibody is directed against a high molecular weight band of the leucocyte common antigen (LCA).
Light chain type	kappa
Concentration	0.5 mg/mL The concentration of this product may be batch-dependent Batch concentration finder →

Reactivity data

Flow Cyt

Tested

Species	Rat
Dilution info	1/2500
Notes	-

Target data

See full target information Ptprc

Function	Protein tyrosine-protein phosphatase required for T-cell activation through the antigen receptor. Acts as a positive regulator of T-cell coactivation upon binding to DPP4. The first PTPase domain has enzymatic activity, while the second one seems to affect the substrate specificity of the first one. Upon T-cell activation, recruits and dephosphorylates SKAP1 and FYN (By similarity). Dephosphorylates LYN, and thereby modulates LYN activity (By similarity). Interacts with CLEC10A at antigen presenting cell-T cell contact; CLEC10A on immature dendritic cells recognizes Tn antigen-carrying PTPRC/CD45 receptor on effector T cells and modulates T cell activation threshold to limit autoreactivity.

Storage

Shipped at conditions	Blue Ice
Appropriate short-term storage duration	1-2 weeks
Appropriate short-term storage conditions	+4°C
Appropriate long-term storage conditions	-20°C
Aliquoting information	Upon delivery aliquot
Storage information	Avoid freeze / thaw cycle, Stable for 12 months at -20°C, Store in the dark

Notes

Spleen cells from immunised BALB/c mice were fused with cells of the NSO/U myeloma cell line. This clone has been described reacting with paraffin embedded material following PLP fixation (see Whiteland et al., 1995). It only labels B cells among thoracic duct lymphocytes, with little labelling in bone marrow and none on thymocytes (Barclay et al.,

1987).Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Supplementary info

This supplementary information is collated from multiple sources and compiled automatically.

Activity summary	CD45RA also known as CD45RA antigen and T200 is an isoform of the CD45 protein family. It is a transmembrane protein tyrosine phosphatase with a molecular mass of approximately 220 kDa. CD45RA expression is abundant on naive T-cells and subsets of other immune cells like B cells. Researchers frequently use CD45RA antibodies including CD45RA-PE and CD45RA-FITC for flow cytometry to identify and study immune cell subsets.
Biological function summary	CD45RA plays a significant role in regulating T-cell receptor (TCR) signaling. By dephosphorylating specific phosphotyrosine residues on substrates CD45RA modulates signal transduction pathways important for T-cell activation and differentiation. This protein does not act alone but is part of a regulatory network that influences immune responses and cellular development through its interaction with other cell surface receptors and intracellular kinases.
Pathways	CD45RA modulation is integral to T-cell development and activation processes. It is an important participant in the T-cell receptor signaling pathway and the Jak-STAT signaling pathway. In these pathways CD45 interacts with proteins such as Lck and Fyn which are essential for propagating receptor-mediated signaling events in T-cells. Effective signal regulation through CD45RA is necessary to ensure appropriate immune responses.
Associated diseases and disorders	CD45RA has links to autoimmune diseases such as multiple sclerosis and rheumatoid arthritis. The aberrant expression or function of CD45RA on immune cells can lead to improper immune activation or suppression contributing to the pathology of these diseases. Crossover with other proteins such as CD45RO which is an isoform involved in memory T-cell function indicates a broader network of immune regulation where CD45RA maintains the balance between immune activation and tolerance.

Product promise

Tested

We have tested this species and application combination and it works. It is covered by our product promise.

Expected

We have not tested this specific species and application combination in-house, but expect it will work. It is covered by our product promise.

Predicted

This species and application combination has not been tested, but we predict it will work based on strong homology. However, this combination is not covered by our product promise.

Not recommended


We do not recommend this combination. It is not covered by our product promise.

We are dedicated to supporting your work with high quality reagents and we are here for you every step of the way should you need us.

In the unlikely event of one of our products not working as expected, you are covered by our product promise.

Full details and terms and conditions can be found here:
Terms & Conditions.

1 product image



Flow Cytometry - Alexa Fluor® 488 Anti-CD45RA antibody [MRC OX-33] (ab256263)

CD45RA Flow Cytometry staining using mouse Anti-CD45RA antibody

Lewis rat splenocytes stained with ab256263 (right) or mouse IgG1κ Alexa Fluor® 488 isotype (left). Lewis rat splenocytes were incubated for 30 min on ice in 1x PBS / 10 % rat serum to block FC receptors and non-specific protein-protein interaction followed by the antibody (ab256263) or mouse IgG1κ Alexa Fluor® 488 isotype (1x 10^6 in 100 µl at 0.2 µg/ml (1/2500)) for 30 min on ice. The cells were simultaneously stained with CD3.
Acquisition of >30,000 events were collected using a 50 mW Blue laser (488nm) and 530/30 bandpass filter. Events were gated on viable lymphocytes.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.